



B4GN2 Monoclonal Antibody

Catalog No	BYmab-05375
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	WB
Gene Name	B4GALNT2 GALGT2
Protein Name	Beta-1,4 N-acetylgalactosaminyltransferase 2 (EC 2.4.1) (Sd(a) beta-1,4-GalNAc transferase) (UDP-GalNAc:Neu5Aca2-3Galb-R b1,4-N-acetylgalactosaminyltransferase)
Immunogen	Synthesized peptide derived from part region of human protein
Specificity	B4GN2 Monoclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	62kD
Cell Pathway	Golgi apparatus membrane ; Single-pass type II membrane protein .
Tissue Specificity	Widely expressed. Highly expressed in colon and to a lesser extent in kidney, stomach, ileum and rectum.
Function	function:Involved in the synthesis of the Sd(a) antigen (Sia-alpha2,3-[GalNAc-beta1,4]Gal-beta1,4-GlcNAc), a carbohydrate determinant expressed on erythrocytes, the colonic mucosa and other tissues. Transfers a beta-1,4-linked GalNAc to the galactose residue of an alpha-2,3-sialylated chain.,online information:Beta-1,4 N-acetylgalactosaminyltransferase 2,online information:GlycoGene database,pathway:Protein modification; protein glycosylation.,similarity:Belongs to the glycosyltransferase 2 family.,tissue specificity:Widely expressed. Highly expressed in colon and to a lesser extent in kidney, stomach, ileum and rectum.,

Nanjing BYabscience technology Co.,Ltd



国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务,欢迎咨询



Background	beta-1,4-N-acetyl-galactosaminyltransferase 2(B4GALNT2) Homo sapiens B4GALNT2 catalyzes the last step in the biosynthesis of the human Sd(a) antigen through the addition of an N-acetylgalactosamine residue via a beta-1,4 linkage to a subterminal galactose residue substituted with an alpha-2,3-linked sialic acid. B4GALNT2 also catalyzes the last step in the biosynthesis of the Cad antigen (Montiel et al., 2003 [PubMed 12678917]).[supplied by OMIM, Mar 2008],
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.
Drodustalmosto	

Products Images

Nanjing BYabscience technology Co.,Ltd