



## CHAC1 Monoclonal Antibody

| by deglycination of the Notch receptor, which prevents receptor maturation and inhibits Notch signaling. This protein may also play a role in the unfolded protein response, and in regulation of glutathione levels and oxidative balance in the cell.   |                    |  |
|---|--------------------|--|
| ReactivityHuman;Rat;Mouse;ApplicationsWBGene NameCHAC1 BOTCHProtein NameCation transport regulator-like protein 1 (Blocks Notch protein) (Botch)ImmunogenSynthesized peptide derived from part region of human proteinSpecificityCHAC1 Monoclonal Antibody detects endogenous levels of protein.FormulationLiquid in PBS containing 50% glycerol, and 0.02% sodium azide.SourceMonoclonal, Mouse,IgGPurificationThe antibody was affinity-purified from mouse antiserum by<br>affinity-chromatography using epitope-specific immunogen.DilutionWB 1:500-2000Concentration1 mg/mlPurity≥90%Storage Stability-20°C/1 yearSynonymsUsing approxed in a specificityObserved Band24kDCell PathwayCytoplasm, cytosol . Golgi apparatus, trans-Golgi network.Tissue SpecificityLiver,Muscle,FunctionTimis gene encodes a member of the gamma-glutamylcyclotransferase family of<br>proteins. The encoded protein may alor a origin appa ar origin the response, and in regulation of glutathione levels and oxidave balance in the coden<br>response, and in regulation of glutathione levels and oxidave balance in the coden<br>to this gene encodes a member of the gamma-glutamylcyclotransferase family of<br>proteins. The encoded protein may also play ar origin the tracent in the cell.<br>Leverstein of this gene may indicate increased risk of cancer recurrence | Catalog No         | BYmab-05442  |
| ApplicationsWBGene NameCHAC1 BOTCHProtein NameCation transport regulator-like protein 1 (Blocks Notch protein) (Botch)ImmunogenSynthesized peptide derived from part region of human proteinSpecificityCHAC1 Monoclonal Antibody detects endogenous levels of protein.FormulationLiquid in PBS containing 50% glycerol, and 0.02% sodium azide.SourceMonoclonal, Mouse, IgGPurificationThe antibody was affinity-purified from mouse antiserum by<br>affinity-chromatography using epitope-specific immunogen.DilutionWB 1:500-2000Concentration1 mg/mlPurity≥90%Storage Stability-20°C/1 yearSynonymsCytoplasm, cytosol . Golgi apparatus, trans-Golgi network .Tissue SpecificityLiver,Muscle,Functionsimilarity:Belongs to the chaC family.,BackgroundThis gene encodes a member of the gamma-glutamylcyclotransferase family of<br>proteins. The encoded protein has been shown to promote neuronal differentiation<br>or by deglycination of the Notch receptor, which prevents receptor maturation and<br>inhibits Notch signaling. This protein may also play a role in the unfolded protein<br>response, and in regulation of glutathione levels and oxidative balance in the cell.   | Isotype            | lgG  |
| Gene Name   CHAC1 BOTCH     Protein Name   Cation transport regulator-like protein 1 (Blocks Notch protein) (Botch)     Immunogen   Synthesized peptide derived from part region of human protein     Specificity   CHAC1 Monoclonal Antibody detects endogenous levels of protein.     Formulation   Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.     Source   Monoclonal, Mouse,IgG     Purification   The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.     Dilution   WB 1:500-2000     Concentration   1 mg/ml     Purity   ≥90%     Storage Stability   -20°C/1 year     Synonyms   Cytoplasm, cytosol . Golgi apparatus, trans-Golgi network .     Tissue Specificity   Liver,Muscle,     Function   similarity:Belongs to the chaC family.,     Background   This gene encodes a member of the gamma-glutamylcyclotransferase family of proteins. The encoded protein has been shown to promote neuronal differentiation of hy deglycination of the Notch receptor, which prevents receptor maturation and inhibits Notch signaling. This protein may also play a role in the unfolded protein in the unfolded protein of glutathione levels and oxidative balance in the cell. Elevaled expression of this gene may indicate increased risk of cancer recurrence  | Reactivity         | Human;Rat;Mouse;   |
| Protein Name Cation transport regulator-like protein 1 (Blocks Notch protein) (Botch)   Immunogen Synthesized peptide derived from part region of human protein   Specificity CHAC1 Monoclonal Antibody detects endogenous levels of protein.   Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.   Source Monoclonal, Mouse,IgG   Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.   Dilution WB 1:500-2000   Concentration 1 mg/ml   Purity ≥90%   Storage Stability -20°C/1 year   Synonyms Cytoplasm, cytosol . Golgi apparatus, trans-Golgi network .   Tissue Specificity Liver,Muscle,   Function similarity:Belongs to the chaC family.,   Background This gene encodes a member of the gamma-glutamylcyclotransferase family of proteins. The encoded protein has been shown to promote neuronal differentiation in this first borts signaling. This protein may also play ar orde in the unfolded protein in the cell. Elevated expression of this gene may indicate increased risk of cancer recurrence  | Applications       | WB   |
| ImmunogenSynthesized peptide derived from part region of human proteinSpecificityCHAC1 Monoclonal Antibody detects endogenous levels of protein.FormulationLiquid in PBS containing 50% glycerol, and 0.02% sodium azide.SourceMonoclonal, Mouse,IgGPurificationThe antibody was affinity-purified from mouse antiserum by<br>affinity-chromatography using epitope-specific immunogen.DilutionWB 1:500-2000Concentration1 mg/mlPurity≥90%Storage Stability-20°C/1 yearSynonyms-20°C/1 yearObserved Band24kDCell PathwayCytoplasm, cytosol . Golgi apparatus, trans-Golgi network .Tissue SpecificityLiver,Muscle,Functionsimilarity:Belongs to the chaC family.,BackgroundThis gene encodes a member of the gamma-glutamylcyclotransferase family of<br>proteins. The encoded protein has been shown to promote neuronal differentiation<br>and in regulation of glutathione levels and oxidative balance in the cell.<br>Elevated expression of this gene encodes and in regulation of glutathione levels and oxidative balance in the cell.<br>Elevated expression of this gene may also play a role in the unfolded protein<br>inspite Notch receptor, which prevents receptor maturation and<br>inhibits Notch signaling. This protein may also play a role in the unfolded protein<br>inspite sont of signed encore recurrence  | Gene Name          | CHAC1 BOTCH  |
| Specificity CHAC1 Monoclonal Antibody detects endogenous levels of protein.   Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.   Source Monoclonal, Mouse,IgG   Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.   Dilution WB 1:500-2000   Concentration 1 mg/ml   Purity ≥90%   Storage Stability -20°C/1 year   Synonyms -20°C/1 year   Observed Band 24kD   Cell Pathway Cytoplasm, cytosol . Golgi apparatus, trans-Golgi network .   Tissue Specificity Liver,Muscle,   Function similarity:Belongs to the chaC family.,   Background This gene encodes a member of the gamma-glutamylcyclotransferase family of proteins. The encoded protein has been shown to promote neuronal differentiation by deglycination of the Notch receptor, which prevents receptor maturation and inhibits Notch signaling. This protein may also play a role in the unfolded protein leuroptical protein and sidative balance in the cell. Elevated expression of this gene may indicate increased risk of cancer recurrence   | Protein Name       | Cation transport regulator-like protein 1 (Blocks Notch protein) (Botch)   |
| Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.   Source Monoclonal, Mouse, IgG   Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.   Dilution WB 1:500-2000   Concentration 1 mg/ml   Purity ≥90%   Storage Stability -20°C/1 year   Synonyms -20°C/1 year   Observed Band 24kD   Cell Pathway Cytoplasm, cytosol . Golgi apparatus, trans-Golgi network .   Tissue Specificity Liver,Muscle,   Function similarity:Belongs to the chaC family.,   Background This gene encodes a member of the gamma-glutamylcyclotransferase family of proteins. The encoded protein has been shown to promote neuronal differentiation py deglycination of the Notch receptor, which prevents receptor maturation and inhibits Notch signaling. This protein may also play a role in the unfolded protein exponse, and in regulation of glutathione levels and oxidative balance in the cell. Elevated expression of this gene may indicate increased risk of cancer recurrence  | Immunogen          | Synthesized peptide derived from part region of human protein  |
| Source Monoclonal, Mouse, IgG   Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.   Dilution WB 1:500-2000   Concentration 1 mg/ml   Purity ≥90%   Storage Stability -20°C/1 year   Synonyms -20°C/1 year   Observed Band 24kD   Cell Pathway Cytoplasm, cytosol . Golgi apparatus, trans-Golgi network .   Tissue Specificity Liver,Muscle,   Function similarity:Belongs to the chaC family.,   Background This gene encodes a member of the gamma-glutamylcyclotransferase family of proteins. The encoded protein has been shown to promote neuronal differentiation in the Notch receptor, which prevents receptor maturation and inhibits Notch signaling. This protein may also play a role in the unfolded protein response, and in regulation of glutathione levels and oxidative balance in the cell. Elevelated expression of this gene may indicate increased risk of cancer recurrence   | Specificity        | CHAC1 Monoclonal Antibody detects endogenous levels of protein.  |
| Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.   Dilution WB 1:500-2000   Concentration 1 mg/ml   Purity ≥90%   Storage Stability -20°C/1 year   Synonyms -20°C/1 year   Observed Band 24kD   Cell Pathway Cytoplasm, cytosol . Golgi apparatus, trans-Golgi network .   Tissue Specificity Liver,Muscle,   Function similarity:Belongs to the chaC family.,   Background This gene encodes a member of the gamma-glutamylcyclotransferase family of proteins. The encoded protein has been shown to promote neuronal differentiation and inhibits Notch signaling. This protein may also play a role in the unfolded protein lesponse, and in regulation of glutathione levels and oxidative balance in the cell. Elevated expression of this gene may indicate increased risk of cancer recurrence   | Formulation        | Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.   |
| affinity-chromatography using epitope-specific immunogen.   Dilution WB 1:500-2000   Concentration 1 mg/ml   Purity ≥90%   Storage Stability -20°C/1 year   Synonyms -20°C/1 year   Observed Band 24kD   Cell Pathway Cytoplasm, cytosol . Golgi apparatus, trans-Golgi network .   Tissue Specificity Liver,Muscle,   Function similarity:Belongs to the chaC family.,   Background This gene encodes a member of the gamma-glutamylcyclotransferase family of proteins. The encoded protein has been shown to promote neuronal differentiation and inhibits Notch signaling. This protein may also play a role in the unfolded protein response, and in regulation of glutathione levels and rok of cancer recurrence   | Source             | Monoclonal, Mouse,IgG  |
| Concentration1 mg/mlPurity≥90%Storage Stability-20°C/1 yearSynonyms-20°C/1 yearObserved Band24kDCell PathwayCytoplasm, cytosol . Golgi apparatus, trans-Golgi network .Tissue SpecificityLiver,Muscle,Functionsimilarity:Belongs to the chaC family.,BackgroundThis gene encodes a member of the gamma-glutamylcyclotransferase family of proteins. The encoded protein has been shown to promote neuronal differentiation by deglycination of the Notch receptor, which prevents receptor maturation and inhibits Notch signaling. This gene may indicate increased risk of cancer recurrence  | Purification       |  |
| Purity ≥90%   Storage Stability -20°C/1 year   Synonyms -20°C/1 year   Observed Band 24kD   Cell Pathway Cytoplasm, cytosol . Golgi apparatus, trans-Golgi network .   Tissue Specificity Liver,Muscle,   Function similarity:Belongs to the chaC family.,   Background This gene encodes a member of the gamma-glutamylcyclotransferase family of proteins. The encoded protein has been shown to promote neuronal differentiation by deglycination of the Notch receptor, which prevents receptor maturation and inhibits Notch signaling. This protein may also play a role in the unfolded protein levels and oxidative balance in the cell. Elevated expression of this gene may indicate increased risk of cancer recurrence  | Dilution           | WB 1:500-2000  |
| Storage Stability -20°C/1 year   Synonyms -20°C/1 year   Observed Band 24kD   Cell Pathway Cytoplasm, cytosol . Golgi apparatus, trans-Golgi network .   Tissue Specificity Liver,Muscle,   Function similarity:Belongs to the chaC family.,   Background This gene encodes a member of the gamma-glutamylcyclotransferase family of proteins. The encoded protein has been shown to promote neuronal differentiation by deglycination of the Notch receptor, which prevents receptor maturation and inhibits Notch signaling. This protein may also play a role in the unfolded protein response, and in regulation of glutathione levels and oxidative balance in the cell. Elevated expression of this gene may indicate increased risk of cancer recurrence   | Concentration      | 1 mg/ml  |
| Synonyms   Observed Band 24kD   Cell Pathway Cytoplasm, cytosol . Golgi apparatus, trans-Golgi network .   Tissue Specificity Liver,Muscle,   Function similarity:Belongs to the chaC family.,   Background This gene encodes a member of the gamma-glutamylcyclotransferase family of proteins. The encoded protein has been shown to promote neuronal differentiation by deglycination of the Notch receptor, which prevents receptor maturation and inhibits Notch signaling. This protein may also play a role in the unfolded protein response, and in regulation of glutathione levels and oxidative balance in the cell. Elevated expression of this gene may indicate increased risk of cancer recurrence   | Purity             | ≥90%   |
| Observed Band24kDCell PathwayCytoplasm, cytosol . Golgi apparatus, trans-Golgi network .Tissue SpecificityLiver,Muscle,Functionsimilarity:Belongs to the chaC family.,BackgroundThis gene encodes a member of the gamma-glutamylcyclotransferase family of<br>proteins. The encoded protein has been shown to promote neuronal differentiation<br>by deglycination of the Notch receptor, which prevents receptor maturation and<br>inhibits Notch signaling. This protein may also play a role in the unfolded protein<br>Elevated expression of this gene may indicate increased risk of cancer recurrence  | Storage Stability  | -20°C/1 year   |
| Cell PathwayCytoplasm, cytosol . Golgi apparatus, trans-Golgi network .Tissue SpecificityLiver,Muscle,Functionsimilarity:Belongs to the chaC family.,BackgroundThis gene encodes a member of the gamma-glutamylcyclotransferase family of<br>proteins. The encoded protein has been shown to promote neuronal differentiation<br>by deglycination of the Notch receptor, which prevents receptor maturation and<br>inhibits Notch signaling. This protein may also play a role in the unfolded protein<br>response, and in regulation of glutathione levels and oxidative balance in the cell.<br>Elevated expression of this gene may indicate increased risk of cancer recurrence   | Synonyms           |  |
| Tissue SpecificityLiver,Muscle,Functionsimilarity:Belongs to the chaC family.,BackgroundThis gene encodes a member of the gamma-glutamylcyclotransferase family of<br>proteins. The encoded protein has been shown to promote neuronal differentiation<br>by deglycination of the Notch receptor, which prevents receptor maturation and<br>inhibits Notch signaling. This protein may also play a role in the unfolded protein<br>response, and in regulation of glutathione levels and oxidative balance in the cell.<br>Elevated expression of this gene may indicate increased risk of cancer recurrence  | Observed Band      | 24kD   |
| Functionsimilarity:Belongs to the chaC family.,BackgroundThis gene encodes a member of the gamma-glutamylcyclotransferase family of<br>proteins. The encoded protein has been shown to promote neuronal differentiation<br>by deglycination of the Notch receptor, which prevents receptor maturation and<br>inhibits Notch signaling. This protein may also play a role in the unfolded protein<br>response, and in regulation of glutathione levels and oxidative balance in the cell.<br>Elevated expression of this gene may indicate increased risk of cancer recurrence   | Cell Pathway       | Cytoplasm, cytosol . Golgi apparatus, trans-Golgi network .  |
| <b>Background</b> This gene encodes a member of the gamma-glutamylcyclotransferase family of proteins. The encoded protein has been shown to promote neuronal differentiation by deglycination of the Notch receptor, which prevents receptor maturation and inhibits Notch signaling. This protein may also play a role in the unfolded protein response, and in regulation of glutathione levels and oxidative balance in the cell. Elevated expression of this gene may indicate increased risk of cancer recurrence   | Tissue Specificity | Liver,Muscle,  |
| proteins. The encoded protein has been shown to promote neuronal differentiation<br>by deglycination of the Notch receptor, which prevents receptor maturation and<br>inhibits Notch signaling. This protein may also play a role in the unfolded protein<br>response, and in regulation of glutathione levels and oxidative balance in the cell.<br>Elevated expression of this gene may indicate increased risk of cancer recurrence  | Function           | similarity:Belongs to the chaC family.,  |
|   | Background         | proteins. The encoded protein has been shown to promote neuronal differentiation<br>by deglycination of the Notch receptor, which prevents receptor maturation and<br>inhibits Notch signaling. This protein may also play a role in the unfolded protein<br>response, and in regulation of glutathione levels and oxidative balance in the cell.<br>Elevated expression of this gene may indicate increased risk of cancer recurrence |

## Nanjing BYabscience technology Co.,Ltd






## **matters needing** Avoid repeated freezing and thawing! attention

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## **Products Images**



Western Blot analysis of various cells using CHAC1 Monoclonal Antibody

Nanjing BYabscience technology Co., Ltd