



CEP72 Monoclonal Antibody

Catalog No	BYmab-05451
Isotype	lgG
Reactivity	Human;Mouse
Applications	WB
Gene Name	CEP72 KIAA1519
Protein Name	Centrosomal protein of 72 kDa (Cep72)
Immunogen	Synthesized peptide derived from part region of human protein
Specificity	CEP72 Monoclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	71kD
Cell Pathway	71kD Cytoplasm, cytoskeleton, microtubule organizing center, centrosome . Cytoplasm, cytoskeleton, microtubule organizing center, centrosome, centriolar satellite . Localizes to the centrosome and centrosome-surrounding particles throughout the cell cycle. These particles disappear after microtubules are depolymerized using nocodazole, suggesting that CEP72-associating particles localize in a microtubule- dependent manner.
	Cytoplasm, cytoskeleton, microtubule organizing center, centrosome . Cytoplasm, cytoskeleton, microtubule organizing center, centrosome, centriolar satellite . Localizes to the centrosome and centrosome-surrounding particles throughout the cell cycle. These particles disappear after microtubules are depolymerized using nocodazole, suggesting that CEP72-associating particles localize in a
Cell Pathway	Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome, centriolar satellite. Localizes to the centrosome and centrosome-surrounding particles throughout the cell cycle. These particles disappear after microtubules are depolymerized using nocodazole, suggesting that CEP72-associating particles localize in a microtubule- dependent manner.
Cell Pathway Tissue Specificity	Cytoplasm, cytoskeleton, microtubule organizing center, centrosome . Cytoplasm, cytoskeleton, microtubule organizing center, centrosome, centriolar satellite . Localizes to the centrosome and centrosome-surrounding particles throughout the cell cycle. These particles disappear after microtubules are depolymerized using nocodazole, suggesting that CEP72-associating particles localize in a microtubule- dependent manner. Brain,Epithelium,Lung,Placenta,Teratocarcinoma,

Nanjing BYabscience technology Co.,Ltd



国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务,欢迎咨询



matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images

Nanjing BYabscience technology Co.,Ltd