



## **CNTFR Monoclonal Antibody**

Catalog No	BYmab-05471
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	CNTFR
Protein Name	Ciliary neurotrophic factor receptor subunit alpha (CNTF receptor subunit alpha) (CNTFR-alpha)
Immunogen	Synthesized peptide derived from part region of human protein
Specificity	CNTFR Monoclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Synonyms Observed Band	40kD
•	40kD Cell membrane; Lipid-anchor, GPI-anchor.
Observed Band	
Observed Band Cell Pathway	Cell membrane; Lipid-anchor, GPI-anchor.  Nervous system and skeletal muscle.  domain:The WSXWS motif appears to be necessary for proper protein folding and thereby efficient intracellular transport and cell-surface receptor binding.,function:Binds to CNTF. The alpha chain provides the receptor specificity.,similarity:Belongs to the type I cytokine receptor family. Type 3 subfamilysimilarity:Contains 1 Iq-like C2-type (immunoglobulin-like)
Observed Band Cell Pathway Tissue Specificity	Cell membrane; Lipid-anchor, GPI-anchor.  Nervous system and skeletal muscle.  domain:The WSXWS motif appears to be necessary for proper protein folding and thereby efficient intracellular transport and cell-surface receptor binding.,function:Binds to CNTF. The alpha chain provides the receptor specificity.,similarity:Belongs to the type I cytokine receptor family. Type 3 subfamily.,similarity:Contains 1 Ig-like C2-type (immunoglobulin-like) domain.,similarity:Contains 2 fibronectin type-III domains.,subunit:Heterotrimer of the alpha chain, LIFR and IL6ST.,tissue specificity:Nervous system and skeletal

Nanjing BYabscience technology Co.,Ltd



国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务,欢迎咨询



differentiation and gene expression. Binding of ciliary neurotrophic factor to the encoded protein recruits the transmembrane components of the receptor, gp130 and leukemia inhibitory factor receptor, facilitating signal transduction. Single nucleotide polymorphisms in this gene may be associated with variations in muscle strength, as well as early onset of eating disorders. Alternatively spliced transcript variants have been observed for this gene. [provided by RefSeq, May 2011],
Avoid repeated freezing and thawing!


Usage suggestions

matters needing

attention

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## **Products Images**



Western Blot analysis of various cells using CNTFR Monoclonal Antibody

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658