



## **DPOA2** Monoclonal Antibody

Catalog No	BYmab-05519
Isotype	lgG
Reactivity	Human;Rat;Mouse;
Applications	WB
Gene Name	POLA2
Protein Name	DNA polymerase alpha subunit B (DNA polymerase alpha 70 kDa subunit)
Immunogen	Synthesized peptide derived from part region of human protein
Specificity	DPOA2 Monoclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	65kD
Cell Pathway	Nucleus.
Tissue Specificity	Cervix epithelium,Epithelium,Lung,
Function	domain:The N-terminal 240 amino acids are sufficient to mediate complex formation.,function:May play an essential role at the early stage of chromosomal DNA replication by coupling the polymerase alpha/primase complex to the cellular replication machinery.,PTM:Phosphorylated in a cell cycle-dependent manner, in G2/M phase.,similarity:Belongs to the DNA polymerase alpha subunit B family.,subunit:DNA polymerase alpha:primase is a four subunit enzyme complex, which is assembled throughout the cell cycle, and consists of the two DNA polymerase subunits A and B, and the DNA primase large and small subunits. Subunit B binds to subunit A.,
Background	domain:The N-terminal 240 amino acids are sufficient to mediate complex formation.,function:May play an essential role at the early stage of chromosomal
	Nanjing BYabscience technology Co.,Ltd



国内优质抗体供应商

Шk 精准的 WB 检测服务 24H 在线服务,欢迎咨询

	——————————————————————————————————————	
	DNA replication by coupling the polymerase alpha/primase complex to the cellular replication machinery.,PTM:Phosphorylated in a cell cycle-dependent manner, in G2/M phase.,similarity:Belongs to the DNA polymerase alpha subunit B family.,subunit:DNA polymerase alpha:primase is a four subunit enzyme complex, which is assembled throughout the cell cycle, and consists of the two DNA polymerase subunits A and B, and the DNA primase large and small subunits. Subunit B binds to subunit A.,	
matters needing attention	Avoid repeated freezing and thawing!	
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.	

**Products Images** 

Nanjing BYabscience technology Co.,Ltd