



HECW1 Monoclonal Antibody

| | |
|---------------------------|---|
| Catalog No | BYmab-05552 |
| Isotype | IgG |
| Reactivity | Human;Rat;Mouse; |
| Applications | WB |
| Gene Name | HECW1 KIAA0322 NEDL1 |
| Protein Name | E3 ubiquitin-protein ligase HECW1 (EC 6.3.2.-) (HECT, C2 and WW domain-containing protein 1) (NEDD4-like E3 ubiquitin-protein ligase 1) (hNEDL1) |
| Immunogen | Synthesized peptide derived from part region of human protein. AA range: 1-67 |
| Specificity | HECW1 Monoclonal Antibody detects endogenous levels of protein. |
| Formulation | Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. |
| Source | Monoclonal, Mouse,IgG |
| Purification | The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. |
| Dilution | WB 1:500-2000 |
| Concentration | 1 mg/ml |
| Purity | ≥90% |
| Storage Stability | -20°C/1 year |
| Synonyms | |
| Observed Band | 176kD |
| Cell Pathway | Cytoplasm . |
| Tissue Specificity | Predominantly expressed in neurons of adult and fetal brain. Weakly expressed in the kidney. |
| Function | function:E3 ubiquitin-protein ligase that mediates ubiquitination and subsequent degradation of DVL1. Also targets the mutant SOD1 protein involved in familial amyotrophic lateral sclerosis (FALS). Forms cytotoxic aggregates with DVL1, SSR3 and mutant SOD1 that lead to motor neuron death in FALS.,pathway:Protein modification; protein ubiquitination.,similarity:Contains 1 C2 domain.,similarity:Contains 1 HECT (E6AP-type E3 ubiquitin-protein ligase) domain.,similarity:Contains 2 WW domains.,subunit:Interacts with DVL1 and SSR3. Also interacts with mutant SOD1.,tissue specificity:Predominantly expressed in neurons of adult and fetal brain. Weakly expressed in the kidney., |
| Background | function:E3 ubiquitin-protein ligase that mediates ubiquitination and subsequent degradation of DVL1. Also targets the mutant SOD1 protein involved in familial |

Nanjing BYabscience technology Co.,Ltd



amyotrophic lateral sclerosis (FALS). Forms cytotoxic aggregates with DVL1, SSR3 and mutant SOD1 that lead to motor neuron death in FALS.,pathway:Protein modification; protein ubiquitination.,similarity:Contains 1 C2 domain.,similarity:Contains 1 HECT (E6AP-type E3 ubiquitin-protein ligase) domain.,similarity:Contains 2 WW domains.,subunit:Interacts with DVL1 and SSR3. Also interacts with mutant SOD1.,tissue specificity:Predominantly expressed in neurons of adult and fetal brain. Weakly expressed in the kidney.,

matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com

官方热线: 025-5229-8998

监督电话: 15950492658