




GCP2 Monoclonal Antibody

mRNA decay,function:Gamma-tubulin complex is necessary for microtubule nucleation at the centrosome.,similarity:Belongs to the GCP family.,subunit:Gamma-tubulin complex is composed of gamma-tubulin, GCP2, GCP3, GCP4, GCP5 and GCP6.,tissue specificity:Ubiquitously expressed., Background alternative products:Additional isoforms may exist. May be produced at very low		
Reactivity Human;Mouse Applications WB Gene Name TUBGCP2 GCP2 Protein Name Gamma-tubulin complex component 2 (GCP-2) (hGCP2) (Gamma-ring complex protein 103 kDa) (h103 p) (hGrip103) (Spindle pole body protein Spc97 homolog) (hSpc97) Immunogen Synthesized peptide derived from part region of human protein Specificity GCP2 Monoclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 99kD Cell Pathway Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Tissue Specificity Ubiquitously expressed. Function alternative products: Additional isoforms may exist. May be produced at very low levels due to a premature stop codon in the mRNA, leading to nonsense-mediated mRNA decay function: Gamma-tubulin complex is necessary for microtubule rucleation at the centrosome, similarity, Be	Catalog No	BYmab-05596
Applications Gene Name TUBGCP2 GCP2 Protein Name Gamma-tubulin complex component 2 (GCP-2) (hGCP2) (Gamma-ring complex protein 103 kDa) (h103p) (hGrip103) (Spindle pole body protein Spc97 homolog) (hSpc97) Immunogen Synthesized peptide derived from part region of human protein Specificity GCP2 Monoclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 99kD Cell Pathway Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Tissue Specificity Ubiquitously expressed. Function alternative products: Additional isoforms may exist. May be produced at very low levels due to a premature stop codon in the mRNA, leading to nonsense-mediated mRNA decay function: Gamma-tubulin complex is necessary for microtubule nucleation at the centrosome. similarity. Belongs to the GCP family, subunit. Gamma-tubulin complex is composed of gamma-tubulin, GCP2, GCP3, GCP4, GCP5 and GCP6, lissue specificity. Way be produced at very low levels due to a premature stop codon in the mRNA, leading to nonsense-mediated mRNA decay, function: Gamma-tubulin complex is composed of gamma-tubulin GCP2. GCP3, GCP4, GCP5 and GCP6, lissue specificity. Ubiquitously expressed., alternative products: Additional isoforms may exist. May be produced at very low levels due to a premature stop codon in the mRNA, leading to nonsense-mediated mRNA decay, function: Gamma-tubulin complex is composed of gamma-tubulin decay function: Gamma-tubulin complex is composed.	Isotype	IgG
Gene Name TUBGCP2 GCP2 Protein Name Gamma-tubullin complex component 2 (GCP-2) (hGCP2) (Gamma-ring complex protein 103 kDa) (h103p) (hGrip103) (Spindle pole body protein Spc97 homolog) (hSpc97) Immunogen Synthesized peptide derived from part region of human protein Specificity GCP2 Monoclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Monoclonal, Mouse, lgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 99kD Cell Pathway Cytoplasm, cytoskeleton, microtubule organizing center, centrosome . Tissue Specificity Ubiquitously expressed. Function alternative products:Additional isoforms may exist. May be produced at very low levels due to a premature stop codon in the mRNA, leading to nonsense-mediated mRNA decay, function:Gamma-tubulin complex is necessary for microtubule nucleation at the centrosome, similarity:Belongs to the GCP family, subunit:Gamma-tubulin complex is composed of gamma-tubulin, GCP2, GCP3, GCP4, GCP5 and GCP6, tissue specificity:Ubiquitou	Reactivity	Human;Mouse
Protein Name Gamma-tubulin complex component 2 (GCP-2) (hGCP2) (Gamma-ring complex protein 103 kDa) (h103p) (hGrip103) (Spindle pole body protein Spc97 homolog) (hSpc97) Immunogen Synthesized peptide derived from part region of human protein Specificity GCP2 Monoclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Monoclonal, Mouse, lgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 99kD Cell Pathway Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Tissue Specificity Ubiquitously expressed. alternative products: Additional isoforms may exist. May be produced at very low levels due to a premature stop codon in the mRNA, leading to nonsense-mediated mRNA decay, function: Gamma-tubulin complex is necessary for microtubule uncleation at the centrosome, similarity: Belongs to the GCP family, subunit: Gamma-tubulin complex is composed of gamma-tubulin, GCP2, GCP3, GCP4, GCP5 and GCP6, tissue specificity: Ubiquitously expressed. Background	Applications	WB
Immunogen Synthesized peptide derived from part region of human protein	Gene Name	TUBGCP2 GCP2
Specificity GCP2 Monoclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 99kD Cell Pathway Cytoplasm, cytoskeleton, microtubule organizing center, centrosome . Tissue Specificity Ubiquitously expressed. Function alternative products:Additional isoforms may exist. May be produced at very low levels due to a premature stop codon in the mRNA, leading to nonsense-mediated mRNA decay, function:Gamma-tubulin complex is composed of gamma-tubulin, GCP2, GCP3, GCP4, GCP5 and GCP6, tissus specificity:Ubiquitously expressed. Background alternative products:Additional isoforms may exist. May be produced at very low levels due to a premature stop codon in the mRNA, leading to nonsense-mediated mRNA decay, function:Gamma-tubulin complex is necessary for microtubule on the products: and the complex is necessary for microtubule on the products of the gamma-tubulin complex is necessary for microtubule on the products and the products are products and the product of the mRNA, leading to n	Protein Name	Gamma-tubulin complex component 2 (GCP-2) (hGCP2) (Gamma-ring complex protein 103 kDa) (h103p) (hGrip103) (Spindle pole body protein Spc97 homolog) (hSpc97)
Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 99kD Cell Pathway Cytoplasm, cytoskeleton, microtubule organizing center, centrosome . Tissue Specificity Ubiquitously expressed. Function alternative products:Additional isoforms may exist. May be produced at very low levels due to a premature stop codon in the mRNA, leading to nonsense-mediated mRNA decay, function:Gamma-tubulin complex is necessary for microtubule nucleation at the centrosome.similarity:Belongs to the GCP familysubunit:Gamma-tubulin complex is composed of gamma-tubulin, GCP2, GCP3, GCP4, GCP5 and GCP6.,tissue specificity:Ubiquitously expressed., Background alternative products:Additional isoforms may exist. May be produced at very low levels due to a premature stop codon in the mRNA, leading to nonsense-mediated mRNA decay, function:Gamma-tubulin complex is composed of pamma-tubulin, GCP2, GCP3, GCP4, GCP5 and GCP6.,tissue specificity:Ubiquitously expressed., Background	Immunogen	Synthesized peptide derived from part region of human protein
Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 99kD Cell Pathway Cytoplasm, cytoskeleton, microtubule organizing center, centrosome . Tissue Specificity Ubiquitously expressed. Function alternative products:Additional isoforms may exist. May be produced at very low levels due to a premature stop codon in the mRNA, leading to nonsense-mediated mRNA decay, function:Gamma-tubulin complex is composed of gamma-tubulin, GCP2, GCP3, GCP4, GCP5 and GCP6., tissue specificity:Ubiquitously expressed., Background alternative products:Additional isoforms may exist. May be produced at very low levels due to a premature stop codon in the mRNA, leading to nonsense-mediated mRNA decay, function:Gamma-tubulin complex is composed of gamma-tubulin GCP2, GCP3, GCP4, GCP5 and GCP6., tissue specificity:Ubiquitously expressed.	Specificity	GCP2 Monoclonal Antibody detects endogenous levels of protein.
Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 99kD Cell Pathway Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Tissue Specificity Ubiquitously expressed. Function alternative products: Additional isoforms may exist. May be produced at very low levels due to a premature stop codon in the mRNA, leading to nonsense-mediated mRNA decay, function:Gamma-tubulin complex is necessary for microtubule nucleation at the centrosomesimilarity:Belongs to the GCP familiy, subunit:Gamma-tubulin complex is composed of gamma-tubulin, GCP2, GCP3, GCP4, GCP5 and GCP6, tissue specificity:Ubiquitously expressed. Background alternative products:Additional isoforms may exist. May be produced at very low levels due to a premature stop codon in the mRNA, leading to nonsense-mediated mRNA decay, function:Gamma-tubulin complex is necessary for microtubule nucleation at the centrosomegamma-tubulin complex is necessary for microtubule nucleation at the centrosome. Supplementation of gamma-tubulin complex is necessary for microtubule nonsense-mediated mRNA decay, function:Gamma-tubulin complex is necessary for microtubule nonsense-mediated mRNA decay, function:Gamma-tubulin complex is necessary for microtubule nonsense-mediated mRNA decay, function:Gamma-tubulin complex is necessary for microtubule nonsense-mediated mRNA decay, function:Gamma-tubulin complex is necessary for microtubule nonsense-mediated mRNA decay, function:Gamma-tubulin complex is necessary for microtubule nonsense-mediated mRNA decay, function:Gamma-tubulin complex is necessary for microtubule nonsense-mediated mRNA decay, function:Gamma-tubulin complex is necessary for microtubule mRNA decay, function:Gamma-tubulin complex is necessary for microtubule	Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 99kD Cell Pathway Cytoplasm, cytoskeleton, microtubule organizing center, centrosome . Tissue Specificity Ubiquitously expressed. Function alternative products:Additional isoforms may exist. May be produced at very low levels due to a premature stop codon in the mRNA, leading to nonsense-mediated mRNA decay, function:Gamma-tubulin complex is necessary for microtubule nucleation at the centrosome. similarity:Belongs to the GCP family., subunit:Gamma-tubulin complex is composed of gamma-tubulin, GCP2, GCP3, GCP4, GCP5 and GCP6, tissue specificity:Ubiquitously expressed., Background alternative products:Additional isoforms may exist. May be produced at very low levels due to a premature stop codon in the mRNA, leading to nonsense-mediated mRNA decay, function:Gamma-tubulin complex is necessary for microtubule nonsense-mediated mRNA decay, function:Gamma-tubulin complex is necessary for microtubule	Source	Monoclonal, Mouse,IgG
Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 99kD Cell Pathway Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Tissue Specificity Ubiquitously expressed. Function alternative products:Additional isoforms may exist. May be produced at very low levels due to a premature stop codon in the mRNA, leading to nonsense-mediated mRNA decay, function:Gamma-tubulin complex is necessary for microtubule nucleation at the centrosome.,similarity:Belongs to the GCP family.,subunit:Gamma-tubulin complex is composed of gamma-tubulin, GCP2, GCP3, GCP4, GCP5 and GCP6.,tissue specificity:Ubiquitously expressed., Background alternative products:Additional isoforms may exist. May be produced at very low levels due to a premature stop codon in the mRNA, leading to nonsense-mediated mRNA decay,function:Gamma-tubulin complex is necessary for microtubule revel due to a premature stop codon in the mRNA, leading to nonsense-mediated mRNA decay, function:Gamma-tubulin complex is necessary for microtubule	Purification	
Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 99kD Cell Pathway Cytoplasm, cytoskeleton, microtubule organizing center, centrosome . Tissue Specificity Ubiquitously expressed. Function alternative products:Additional isoforms may exist. May be produced at very low levels due to a premature stop codon in the mRNA, leading to nonsense-mediated mRNA decay,function:Gamma-tubulin complex is necessary for microtubule nucleation at the centrosome.,similarity:Belongs to the GCP family.,subunit:Gamma-tubulin complex is composed of gamma-tubulin, GCP2, GCP3, GCP4, GCP5 and GCP6.,tissue specificity:Ubiquitously expressed., Background alternative products:Additional isoforms may exist. May be produced at very low levels due to a premature stop codon in the mRNA, leading to nonsense-mediated mRNA decay,function:Gamma-tubulin complex is necessary for microtubule	Dilution	WB 1:500-2000
Storage Stability -20°C/1 year Synonyms Observed Band 99kD Cell Pathway Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Tissue Specificity Ubiquitously expressed. Function alternative products: Additional isoforms may exist. May be produced at very low levels due to a premature stop codon in the mRNA, leading to nonsense-mediated mRNA decay, function: Gamma-tubulin complex is necessary for microtubule nucleation at the centrosome., similarity: Belongs to the GCP family., subunit: Gamma-tubulin complex is composed of gamma-tubulin, GCP2, GCP3, GCP4, GCP5 and GCP6., tissue specificity: Ubiquitously expressed., Background alternative products: Additional isoforms may exist. May be produced at very low levels due to a premature stop codon in the mRNA, leading to nonsense-mediated mRNA decay, function: Gamma-tubulin complex is necessary for microtubule	Concentration	1 mg/ml
Synonyms Observed Band 99kD Cell Pathway Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Tissue Specificity Ubiquitously expressed. Function alternative products:Additional isoforms may exist. May be produced at very low levels due to a premature stop codon in the mRNA, leading to nonsense-mediated mRNA decay, function:Gamma-tubulin complex is necessary for microtubule nucleation at the centrosome., similarity:Belongs to the GCP family., subunit:Gamma-tubulin complex is composed of gamma-tubulin, GCP2, GCP3, GCP4, GCP5 and GCP6.,tissue specificity:Ubiquitously expressed., Background alternative products:Additional isoforms may exist. May be produced at very low levels due to a premature stop codon in the mRNA, leading to nonsense-mediated mRNA decay, function:Gamma-tubulin complex is necessary for microtubule	Purity	≥90%
Observed Band Cell Pathway Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Tissue Specificity Ubiquitously expressed. Function alternative products:Additional isoforms may exist. May be produced at very low levels due to a premature stop codon in the mRNA, leading to nonsense-mediated mRNA decay,function:Gamma-tubulin complex is necessary for microtubule nucleation at the centrosome.,similarity:Belongs to the GCP family.,subunit:Gamma-tubulin complex is composed of gamma-tubulin, GCP2, GCP3, GCP4, GCP5 and GCP6.,tissue specificity:Ubiquitously expressed., Background alternative products:Additional isoforms may exist. May be produced at very low levels due to a premature stop codon in the mRNA, leading to nonsense-mediated mRNA decay,function:Gamma-tubulin complex is necessary for microtubule	Storage Stability	-20°C/1 year
Cell Pathway Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Ubiquitously expressed. Function alternative products:Additional isoforms may exist. May be produced at very low levels due to a premature stop codon in the mRNA, leading to nonsense-mediated mRNA decay,function:Gamma-tubulin complex is necessary for microtubule nucleation at the centrosome.,similarity:Belongs to the GCP family.,subunit:Gamma-tubulin complex is composed of gamma-tubulin, GCP2, GCP3, GCP4, GCP5 and GCP6.,tissue specificity:Ubiquitously expressed., Background alternative products:Additional isoforms may exist. May be produced at very low levels due to a premature stop codon in the mRNA, leading to nonsense-mediated mRNA decay,function:Gamma-tubulin complex is necessary for microtubule	Synonyms	
Tissue Specificity Ubiquitously expressed. Function alternative products:Additional isoforms may exist. May be produced at very low levels due to a premature stop codon in the mRNA, leading to nonsense-mediated mRNA decay,function:Gamma-tubulin complex is necessary for microtubule nucleation at the centrosome.,similarity:Belongs to the GCP family.,subunit:Gamma-tubulin complex is composed of gamma-tubulin, GCP2, GCP3, GCP4, GCP5 and GCP6.,tissue specificity:Ubiquitously expressed., Background alternative products:Additional isoforms may exist. May be produced at very low levels due to a premature stop codon in the mRNA, leading to nonsense-mediated mRNA decay,function:Gamma-tubulin complex is necessary for microtubule	Observed Band	99kD
Function alternative products:Additional isoforms may exist. May be produced at very low levels due to a premature stop codon in the mRNA, leading to nonsense-mediated mRNA decay,function:Gamma-tubulin complex is necessary for microtubule nucleation at the centrosome.,similarity:Belongs to the GCP family.,subunit:Gamma-tubulin complex is composed of gamma-tubulin, GCP2, GCP3, GCP4, GCP5 and GCP6.,tissue specificity:Ubiquitously expressed., Background alternative products:Additional isoforms may exist. May be produced at very low levels due to a premature stop codon in the mRNA, leading to nonsense-mediated mRNA decay,function:Gamma-tubulin complex is necessary for microtubule	Cell Pathway	Cytoplasm, cytoskeleton, microtubule organizing center, centrosome .
levels due to a premature stop codon in the mRNA, leading to nonsense-mediated mRNA decay, function: Gamma-tubulin complex is necessary for microtubule nucleation at the centrosome., similarity: Belongs to the GCP family., subunit: Gamma-tubulin complex is composed of gamma-tubulin, GCP2, GCP3, GCP4, GCP5 and GCP6., tissue specificity: Ubiquitously expressed., Background alternative products: Additional isoforms may exist. May be produced at very low levels due to a premature stop codon in the mRNA, leading to nonsense-mediated mRNA decay, function: Gamma-tubulin complex is necessary for microtubule	Tissue Specificity	Ubiquitously expressed.
levels due to a premature stop codon in the mRNA, leading to nonsense-mediated mRNA decay,function:Gamma-tubulin complex is necessary for microtubule	Function	levels due to a premature stop codon in the mRNA, leading to nonsense-mediated mRNA decay, function: Gamma-tubulin complex is necessary for microtubule nucleation at the centrosome., similarity: Belongs to the GCP family., subunit: Gamma-tubulin complex is composed of gamma-tubulin, GCP2,
	Background	levels due to a premature stop codon in the mRNA, leading to nonsense-mediated mRNA decay, function: Gamma-tubulin complex is necessary for microtubule

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658







	family.,subunit:Gamma-tubulin complex is composed of gamma-tubulin, GCP2, GCP3, GCP4, GCP5 and GCP6.,tissue specificity:Ubiquitously expressed.,
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658