



RGPA1 Monoclonal Antibody

Isotype IgG		
Reactivity Human;Mouse;Rat Applications WB Gene Name RALGAPA1 GARNL1 KIAA0884 TULIP1 Protein Name Ral GTPase-activating protein subunit alpha-1 (GAP-related-interacting partner to £12) (GRIPE) (GTPase-activating Rap/Ran-GAP domain-like 1) (Tuberin-like protein 1) (p240) Immunogen Synthesized peptide derived from part region of human protein Specificity RGPA1 Monoclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 223kD Cell Pathway Cytoplasm . Nucleus . Translocated to the nucleus, when associated with TCF3/E12 Tissue Specificity Widely expressed. Function regulation of small GTPase mediated signal transduction, This gene encodes a major subunit of the RAL-GTPase activating protein. A similar protein in mouse binds E12, a transcriptional regulator of immunoglobulin genes. The mouse protein also functions in skeletal muscle by binding to the regulatory 14-3-3 proteins upon stimulation with insulin or muscle contraction. A pseudogene of this gene has been identified on chromosome 9. [provided by speudogene of this gene has been identified on chromosome 9. [provided by speudogene of this gene has been identified on chromosome 9. [provided by pseudogene of this gene has been identified on chromosome 9.] provided by contraction.	Catalog No	BYmab-05613
Applications WB Gene Name RALGAPA1 GARNL1 KIAA0884 TULIP1 Protein Name Ral GTPase-activating protein subunit alpha-1 (GAP-related-interacting partner to E12) (GRIPE) (GTPase-activating Rap/Ran-GAP domain-like 1) (Tuberin-like protein 1) (p240) Immunogen Synthesized peptide derived from part region of human protein Specificity RGPA1 Monoclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Monoclonal, Mouse, lgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 223kD Cell Pathway Cytoplasm . Nucleus . Translocated to the nucleus, when associated with TCF3/E12. Tissue Specificity Widely expressed. Function regulation of small GTPase mediated signal transduction, This gene encodes a major subunit of the RAL-GTPase activating protein. A similar protein in mouse binds E12, a transcriptional regulator of immunoglobulin species. The mouse protein also functions in skeletal muscle by binding to the regulatory 14-3-3 proteins upon stimulation with insulin or muscle by binding to the regulatory 14-3-3 proteins upon stimulation with insulin or muscle by provided by pseudogene of this gene has been identified on chromosome 9. [provided by	Isotype	IgG
Gene Name RALGAPA1 GARNL1 KIAA0884 TULIP1 Protein Name Ral GTPase-activating protein subunit alpha-1 (GAP-related-interacting partner to E12) (GRIPE) (GTPase-activating Rap/Ran-GAP domain-like 1) (Tuberin-like protein 1) (p240) Immunogen Synthesized peptide derived from part region of human protein Specificity RGPA1 Monoclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Monoclonal, Mouse,lgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 223kD Cell Pathway Cytoplasm . Nucleus . Translocated to the nucleus, when associated with TCF3/E12 Tissue Specificity Widely expressed. Function regulation of small GTPase mediated signal transduction, Background This gene encodes a major subunit of the RAL-GTPase activating protein. A similar protein in mouse binds E12, a transcriptional regulator of immunoglobulin genes. The mouse protein also functions in skeletal muscle by binding to the regulatory 14-3-3 proteins upon stimulation with insulin or muscle contraction. A pseudogene of this gene has been identified on chromosome 9. [provided by pseudogene of this gene has been identified on chromosome 9. [provided by pseudogene of this gene has been identified on chromosome 9. [provided by	Reactivity	Human;Mouse;Rat
Protein Name Ral GTPase-activating protein subunit alpha-1 (GAP-related-interacting partner to E12) (GRIPE) (GTPase-activating Rap/Ran-GAP domain-like 1) (Tuberin-like protein 1) (p240) Immunogen Synthesized peptide derived from part region of human protein Specificity RGPA1 Monoclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Monoclonal, Mouse, lgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity 290% Storage Stability -20°C/1 year Synonyms Observed Band 223kD Cell Pathway Cytoplasm . Nucleus . Translocated to the nucleus, when associated with TCF3/E12. Tissue Specificity Widely expressed. Function regulation of small GTPase mediated signal transduction, Background This gene encodes a major subunit of the RAL-GTPase activating protein. A similar protein in mouse binds E12, a transcriptional regulator of immunoglobulin genes. The mouse protein also functions in skeletal muscle by binding to the regulatory 14-3-3 proteins upon stimulation with insulin or muscle contraction. A pseudogene of this gene has been identified on chromosome 9. [provided by pseudogene of this gene has been identified on chromosome 9. [provided by pseudogene of this gene has been identified on chromosome 9. [provided by	Applications	WB
Immunogen Synthesized peptide derived from part region of human protein Specificity RGPA1 Monoclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 223kD Cell Pathway Cytoplasm · Nucleus · Translocated to the nucleus, when associated with TCF3/E12. Tissue Specificity Widely expressed. Function regulation of small GTPase mediated signal transduction, Background This gene encodes a major subunit of the RAL-GTPase activating protein. A similar protein in mouse binds E12, a transcriptional regulator of immunoglobulin genes. The mouse protein also functions in skeletal muscle by binding to the regulatory 14-3-3 proteins upon stimulation with insulin or muscle contraction. A pseudogene of this gene has been identified on chromosome 9. [provided by provided by	Gene Name	RALGAPA1 GARNL1 KIAA0884 TULIP1
Specificity RGPA1 Monoclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 223kD Cell Pathway Cytoplasm Nucleus . Translocated to the nucleus, when associated with TCF3/E12 . Tissue Specificity Widely expressed. Function regulation of small GTPase mediated signal transduction, Background This gene encodes a major subunit of the RAL-GTPase activating protein. A similar protein in mouse binds E12, a transcriptional regulator of immunoglobulin genes. The mouse protein also functions in skeletal muscle by binding to the regulatory 14-3-3 proteins upon stimulation with insulin or muscle contraction. A pseudogene of this gene has been identified on chromosome 9. [provided by	Protein Name	Ral GTPase-activating protein subunit alpha-1 (GAP-related-interacting partner to E12) (GRIPE) (GTPase-activating Rap/Ran-GAP domain-like 1) (Tuberin-like protein 1) (p240)
Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Cytoplasm . Nucleus . Translocated to the nucleus, when associated with TCF3/E12. Tissue Specificity Widely expressed. Function regulation of small GTPase mediated signal transduction, Background This gene encodes a major subunit of the RAL-GTPase activating protein. A similar protein in mouse binds E12, a transcriptional regulator of immunoglobulin genes. The mouse protein also functions in skeletal muscle by binding to the regulatory 14-3-3 proteins upon stimulation with insulin or muscle contraction. A pseudogene of this gene has been identified on chromosome 9. [provided by	Immunogen	Synthesized peptide derived from part region of human protein
Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Cobserved Band Cell Pathway Cytoplasm . Nucleus . Translocated to the nucleus, when associated with TCF3/E12. Tissue Specificity Widely expressed. Function regulation of small GTPase mediated signal transduction, Background This gene encodes a major subunit of the RAL-GTPase activating protein. A similar protein in mouse binds E12, a transcriptional regulator of immunoglobulin genes. The mouse protein also functions in skeletal muscle by binding to the regulatory 14-3-3 proteins upon stimulation with insulin or muscle contraction. A pseudogene of this gene has been identified on chromosome 9. [provided by	Specificity	RGPA1 Monoclonal Antibody detects endogenous levels of protein.
Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 223kD Cell Pathway Cytoplasm . Nucleus . Translocated to the nucleus, when associated with TCF3/E12 Tissue Specificity Widely expressed. Function regulation of small GTPase mediated signal transduction, Background This gene encodes a major subunit of the RAL-GTPase activating protein. A similar protein in mouse binds E12, a transcriptional regulator of immunoglobulin genes. The mouse protein also functions in skeletal muscle by binding to the regulatory 14-3-3 proteins upon stimulation with insulin or muscle contraction. A pseudogene of this gene has been identified on chromosome 9. [provided by	Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 223kD Cell Pathway Cytoplasm . Nucleus . Translocated to the nucleus, when associated with TCF3/E12 Tissue Specificity Widely expressed. Function regulation of small GTPase mediated signal transduction, Background This gene encodes a major subunit of the RAL-GTPase activating protein. A similar protein in mouse binds E12, a transcriptional regulator of immunoglobulin genes. The mouse protein also functions in skeletal muscle by binding to the regulatory 14-3-3 proteins upon stimulation with insulin or muscle contraction. A pseudogene of this gene has been identified on chromosome 9. [provided by	Source	Monoclonal, Mouse,IgG
Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms	Purification	· · · · · · · · · · · · · · · · · · ·
Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 223kD Cell Pathway Cytoplasm . Nucleus . Translocated to the nucleus, when associated with TCF3/E12 Tissue Specificity Widely expressed. Function regulation of small GTPase mediated signal transduction, Background This gene encodes a major subunit of the RAL-GTPase activating protein. A similar protein in mouse binds E12, a transcriptional regulator of immunoglobulin genes. The mouse protein also functions in skeletal muscle by binding to the regulatory 14-3-3 proteins upon stimulation with insulin or muscle contraction. A pseudogene of this gene has been identified on chromosome 9. [provided by	Dilution	WB 1:500-2000
Storage Stability Synonyms Observed Band Cell Pathway Cytoplasm . Nucleus . Translocated to the nucleus, when associated with TCF3/E12 Tissue Specificity Widely expressed. Function regulation of small GTPase mediated signal transduction, Background This gene encodes a major subunit of the RAL-GTPase activating protein. A similar protein in mouse binds E12, a transcriptional regulator of immunoglobulin genes. The mouse protein also functions in skeletal muscle by binding to the regulatory 14-3-3 proteins upon stimulation with insulin or muscle contraction. A pseudogene of this gene has been identified on chromosome 9. [provided by	Concentration	1 mg/ml
Synonyms Observed Band 223kD Cell Pathway Cytoplasm . Nucleus . Translocated to the nucleus, when associated with TCF3/E12 Tissue Specificity Widely expressed. Function regulation of small GTPase mediated signal transduction, Background This gene encodes a major subunit of the RAL-GTPase activating protein. A similar protein in mouse binds E12, a transcriptional regulator of immunoglobulin genes. The mouse protein also functions in skeletal muscle by binding to the regulatory 14-3-3 proteins upon stimulation with insulin or muscle contraction. A pseudogene of this gene has been identified on chromosome 9. [provided by	Purity	≥90%
Observed Band 223kD Cell Pathway Cytoplasm . Nucleus . Translocated to the nucleus, when associated with TCF3/E12 Tissue Specificity Widely expressed. Function regulation of small GTPase mediated signal transduction, Background This gene encodes a major subunit of the RAL-GTPase activating protein. A similar protein in mouse binds E12, a transcriptional regulator of immunoglobulin genes. The mouse protein also functions in skeletal muscle by binding to the regulatory 14-3-3 proteins upon stimulation with insulin or muscle contraction. A pseudogene of this gene has been identified on chromosome 9. [provided by	Storage Stability	-20°C/1 year
Cell Pathway Cytoplasm . Nucleus . Translocated to the nucleus, when associated with TCF3/E12 Tissue Specificity Widely expressed. Function regulation of small GTPase mediated signal transduction, This gene encodes a major subunit of the RAL-GTPase activating protein. A similar protein in mouse binds E12, a transcriptional regulator of immunoglobulin genes. The mouse protein also functions in skeletal muscle by binding to the regulatory 14-3-3 proteins upon stimulation with insulin or muscle contraction. A pseudogene of this gene has been identified on chromosome 9. [provided by	Synonyms	
TCF3/E12 Tissue Specificity Widely expressed. Function regulation of small GTPase mediated signal transduction, This gene encodes a major subunit of the RAL-GTPase activating protein. A similar protein in mouse binds E12, a transcriptional regulator of immunoglobulin genes. The mouse protein also functions in skeletal muscle by binding to the regulatory 14-3-3 proteins upon stimulation with insulin or muscle contraction. A pseudogene of this gene has been identified on chromosome 9. [provided by	Observed Band	223kD
Function regulation of small GTPase mediated signal transduction, Background This gene encodes a major subunit of the RAL-GTPase activating protein. A similar protein in mouse binds E12, a transcriptional regulator of immunoglobulin genes. The mouse protein also functions in skeletal muscle by binding to the regulatory 14-3-3 proteins upon stimulation with insulin or muscle contraction. A pseudogene of this gene has been identified on chromosome 9. [provided by	Cell Pathway	Cytoplasm . Nucleus . Translocated to the nucleus, when associated with TCF3/E12
This gene encodes a major subunit of the RAL-GTPase activating protein. A similar protein in mouse binds E12, a transcriptional regulator of immunoglobulin genes. The mouse protein also functions in skeletal muscle by binding to the regulatory 14-3-3 proteins upon stimulation with insulin or muscle contraction. A pseudogene of this gene has been identified on chromosome 9. [provided by	Tissue Specificity	Widely expressed.
similar protein in mouse binds E12, a transcriptional regulator of immunoglobulin genes. The mouse protein also functions in skeletal muscle by binding to the regulatory 14-3-3 proteins upon stimulation with insulin or muscle contraction. A pseudogene of this gene has been identified on chromosome 9. [provided by	Function	regulation of small GTPase mediated signal transduction,
	Background	similar protein in mouse binds E12, a transcriptional regulator of immunoglobulin genes. The mouse protein also functions in skeletal muscle by binding to the regulatory 14-3-3 proteins upon stimulation with insulin or muscle contraction. A pseudogene of this gene has been identified on chromosome 9. [provided by

Nanjing BYabscience technology Co.,Ltd



国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务, 欢迎咨询



matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images

Nanjing BYabscience technology Co.,Ltd