



TGIF2 Monoclonal Antibody

Catalog No	BYmab-05669
Isotype	IgG
Reactivity	Human;Mouse
Applications	WB
Gene Name	TGIF2
Protein Name	Homeobox protein TGIF2 (5'-TG-3'-interacting factor 2) (TGF-beta-induced transcription factor 2) (TGFB-induced factor 2)
Immunogen	Synthesized peptide derived from part region of human protein
Specificity	TGIF2 Monoclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	26kD
Cell Pathway	Nucleus. Excluded from nucleoli.
Tissue Specificity	Widely expressed. Highly expressed in heart, kidney and testis. Weakly expressed in brain and prostate.
Function	function:Transcriptional repressor, which probably repress transcription by binding directly the 5'-CTGTCAA-3' DNA sequence or by interacting with TGF-beta activated SMAD proteins. Probably represses transcription via the recruitment of histone deacetylase proteins.,miscellaneous:TGIF2 is amplified and overexpressed in several ovarian cancer cell lines.,PTM:The C-terminal part is phosphorylated in response to EGF signaling by the Ras/MAPK pathway.,similarity:Belongs to the TALE/TGIF homeobox family.,similarity:Contains 1 homeobox DNA-binding domain.,subcellular location:Excluded from nucleoli.,subunit:Interacts with the transcriptional modulator SMAD3 and the histone deacetylase HDAC1.,tissue specificity:Widely expressed. Highly expressed in heart, kidney and testis. Weakly expressed in

Nanjing BYabscience technology Co.,Ltd



brain and prostate.,

Background

The protein encoded by this gene is a DNA-binding homeobox protein and a transcriptional repressor, which appears to repress transcription by recruiting histone deacetylases to TGF beta-responsive genes. This gene is amplified and over-expressed in some ovarian cancers. Alternative splicing results in multiple transcript variants. A related pseudogene has been identified on chromosome 1. Read-through transcription also exists between this gene and the neighboring downstream C20orf24 (chromosome 20 open reading frame 24) gene. [provided by RefSeq, Dec 2010],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using TGIF2 Monoclonal Antibody

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com

官方热线: 025-5229-8998

监督电话: 15950492658