



# HAP1 Monoclonal Antibody

<b>Catalog No</b>	BYmab-05673
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human;Rat;Mouse
<b>Applications</b>	WB
<b>Gene Name</b>	HAP1 HAP2 HLP1
<b>Protein Name</b>	Huntingtin-associated protein 1 (HAP-1) (Neuroan 1)
<b>Immunogen</b>	Synthesized peptide derived from part region of human protein
<b>Specificity</b>	HAP1 Monoclonal Antibody detects endogenous levels of protein.
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
<b>Source</b>	Monoclonal, Mouse,IgG
<b>Purification</b>	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Dilution</b>	WB 1:500-2000
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	
<b>Observed Band</b>	73kD
<b>Cell Pathway</b>	Cytoplasm . Cell projection, axon . Cell junction, synapse, presynapse . Cytoplasm, cytoskeleton . Cell projection, dendritic spine . Cell projection, dendrite . Lysosome . Endoplasmic reticulum . Mitochondrion . Nucleus . Cytoplasmic vesicle, autophagosome . Early endosome . Cell projection, growth cone . Cell projection, neuron projection . Cytoplasmic vesicle, secretory vesicle, synaptic vesicle . Localizes to large nonmembrane-bound cytoplasmic bodies found in various types of neurons, called stigmoid bodies (STBs). Localization to neuronal processes and neurite tips is decreased by YWHAZ. In the nucleus localizes to nuclear rods. .
<b>Tissue Specificity</b>	Predominantly expressed in brain. Selectively expressed in neurons.
<b>Function</b>	alternative products:Additional isoforms seem to exist,caution:The sequence shown here is derived from an Ensembl automatic analysis pipeline and should be considered as preliminary data.,function:Associates specifically with huntingtin. This binding is enhanced by an expanded polyglutamine repeat.,similarity:Contains 1 HAP1 N-terminal domain.,tissue

**Nanjing BYabscience technology Co.,Ltd**



specificity:Predominantly expressed in brain. Selectively expressed in neurons.,

**Background**

Huntington's disease (HD), a neurodegenerative disorder characterized by loss of striatal neurons, is caused by an expansion of a polyglutamine tract in the HD protein huntingtin. This gene encodes a protein that interacts with huntingtin, with two cytoskeletal proteins (dynactin and pericentriolar autoantigen protein 1), and with a hepatocyte growth factor-regulated tyrosine kinase substrate. The interactions with cytoskeletal proteins and a kinase substrate suggest a role for this protein in vesicular trafficking or organelle transport. Several alternatively spliced transcript variants encoding different isoforms have been described for this gene. [provided by RefSeq, Jul 2008],

**matters needing attention**

Avoid repeated freezing and thawing!

**Usage suggestions**

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## Products Images

Nanjing BYabscience technology Co.,Ltd

网址: [www.njbybio.com](http://www.njbybio.com)

官方热线: 025-5229-8998

监督电话: 15950492658