



UBP54 Monoclonal Antibody

| | |
|---------------------------|---|
| Catalog No | BYmab-05679 |
| Isotype | IgG |
| Reactivity | Human;Rat;Mouse |
| Applications | WB |
| Gene Name | USP54 C10orf29 |
| Protein Name | Inactive ubiquitin carboxyl-terminal hydrolase 54 (Inactive ubiquitin-specific peptidase 54) |
| Immunogen | Synthesized peptide derived from part region of human protein |
| Specificity | UBP54 Monoclonal Antibody detects endogenous levels of protein. |
| Formulation | Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. |
| Source | Monoclonal, Mouse,IgG |
| Purification | The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. |
| Dilution | WB 1:500-2000 |
| Concentration | 1 mg/ml |
| Purity | ≥90% |
| Storage Stability | -20°C/1 year |
| Synonyms | |
| Observed Band | 185kD |
| Cell Pathway | |
| Tissue Specificity | Weakly expressed in a few tissues. |
| Function | caution:Although the active site residues are conserved, it lacks the conserved His residue which is normally found 9 residues before the catalytic His.,function:Has no peptidase activity.,similarity:Belongs to the peptidase C19 family.,tissue specificity:Weakly expressed in a few tissues., |
| Background | caution:Although the active site residues are conserved, it lacks the conserved His residue which is normally found 9 residues before the catalytic His.,function:Has no peptidase activity.,similarity:Belongs to the peptidase C19 family.,tissue specificity:Weakly expressed in a few tissues., |
| matters needing attention | Avoid repeated freezing and thawing! |


Nanjing BYabscience technology Co.,Ltd



Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using UBP54 Monoclonal Antibody