



## MYO3B Monoclonal Antibody

2 IQ domains.,similarity:In the N-terminal section; belongs to the protein kinase superfamily. STE Ser/Thr protein kinase family.,tissue specificity:Expressed in retina, kidney and testis.,  Background  This gene encodes one of the class III myosins. Myosins are ATPases, activated by actin, that move along actin filaments in the cell. This class of myosins are characterized by an amino-terminal kinase domain and shown to be present in		
Reactivity         Human;Rat;Mouse;           Applications         WB           Gene Name         MYO3B           Protein Name         Myosin-IIIb (EC 2.7.11.1)           Immunogen         Synthesized peptide derived from human protein . at AA range: 1030-1110           Specificity         MYO3B Monoclonal Antibody detects endogenous levels of protein.           Formulation         Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.           Source         Monoclonal, Mouse, IgG           Purification         The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.           Dilution         WB 1:500-2000           Concentration         1 mg/ml           Purity         ≥90%           Storage Stability         -20°C/1 year           Synonyms         Observed Band         147kD           Cell Pathway         Cytoplasm, cytoskeleton. Cell projection, stereocilium.           Tissue Specificity         Expressed in retina, kidney and testis.           Function         catalytic activity. ATP + a protein = ADP + a phosphoprotein, function:Probable actin-based motor with a protein kinase activity, similarity. Contains: 1 protein kinase driving islarity. Contains: 2 [10 domains, similarity. 10 the N-terminal section; belongs to the protein kinase superfamily, ISTE Ser/Th protein kinase family., tissue specificity:Expressed in retina, kidney and testis.	Catalog No	BYmab-05779
Applications WB Gene Name MYO3B Protein Name Myosin-Illb (EC 2.7.11.1) Immunogen Synthesized peptide derived from human protein . at AA range: 1030-1110 Specificity MYO3B Monoclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity 290% Storage Stability -20°C/1 year Synonyms Observed Band 147kD Cell Pathway Cytoplasm, cytoskeleton. Cell projection, stereocilium. Tissue Specificity Expressed in retina, kidney and testis. Function catalytic activity:ATP + a protein = ADP + a phosphoprotein function:Probable actin-based motor with a protein kinase activity, similarity. Contains 1 protein kinase domain. similarity. STE Ser/Th protein kinase activity, issue specificity: Expressed in retina, kidney and testis. This gene encodes one of the class III myosins are ATPases, activated by actin, that move along actin filaments in the cell. This class of myosins are characterized by an amino-terminal kinase domain and shown to be present in photoreceptors. Alternative splicing results in multiple transcrivations beneding results in multiple transcrivations of the protein conditions are conditional conditions.	Isotype	IgG
Gene Name MyO3B  Protein Name Myosin-IIIb (EC 2.7.11.1)  Immunogen Synthesized peptide derived from human protein . at AA range: 1030-1110  Specificity MYO3B Monoclonal Antibody detects endogenous levels of protein.  Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.  Source Monoclonal, Mouse, IgG  Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-2000  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms  Observed Band 147kD  Cell Pathway Cytoplasm, cytoskeleton. Cell projection, stereocilium.  Tissue Specificity Expressed in retina, kidney and testis.  Function catalytic activity.ATP + a protein = ADP + a phosphoprotein.,function:Probable actin-based motor with a protein kinase activity, similarity:Contains 1 myosin head-like domain, similarity:Contains 1 protein kinase domain, similarity:Contains 2 IQ domains, similarity:In the N-terminal section; belongs to the protein kinase superfamily. STE Ser? Thr protein kinase family, tissue specificity:Expressed in retina, kidney and testis.  Background This gene encodes one of the class III myosis me ATPases, activated by actin, that move along actin filaments in the cell. This class of myosins are characterized by an amino-terminal kinase domain and shown to be present in photoreceptors. Alternative splicing results in multiple transcript variants encoding	Reactivity	Human;Rat;Mouse;
Protein Name         Myosin-Illb (EC 2.7.11.1)           Immunogen         Synthesized peptide derived from human protein . at AA range: 1030-1110           Specificity         MYO3B Monoclonal Antibody detects endogenous levels of protein.           Formulation         Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.           Source         Monoclonal, Mouse, lgG           Purification         The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.           Dilution         WB 1:500-2000           Concentration         1 mg/ml           Purity         ≥90%           Storage Stability         -20°C/1 year           Synonyms         Cytoplasm, cytoskeleton. Cell projection, stereocilium.           Tissue Specificity         Expressed in retina, kidney and testis.           Function         catalytic activity:ATP + a protein = ADP + a phosphoprotein, function:Probable actin-based motor with a protein kinase activity, similarity:Contains 1 myosin head-like domain, similarity:Contains 1 myosin head-like domain, similarity:Contains 1 protein kinase activity, similarity:Contains 1 protein kinase activity, similarity:Contains 1 protein kinase superfamily. STE Ser/Thr protein kinase family, tissue specificity:Expressed in retina, kidney and testis.           Background         This gene encodes one of the class Ill myosins. Myosins are ATPases, activated by actin, that move along actin filaments in the cell. This class of myosins are characterized by an amino-termi	Applications	WB
Immunogen         Synthesized peptide derived from human protein . at AA range: 1030-1110           Specificity         MYO3B Monoclonal Antibody detects endogenous levels of protein.           Formulation         Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.           Source         Monoclonal, Mouse, IgG           Purification         The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.           Dilution         WB 1:500-2000           Concentration         1 mg/ml           Purity         ≥90%           Storage Stability         -20°C/1 year           Synonyms         Observed Band         147kD           Cell Pathway         Cytoplasm, cytoskeleton. Cell projection, stereocilium.           Tissue Specificity         Expressed in retina, kidney and testis.           Function         catalytic activity:ATP + a protein = ADP + a phosphoprotein "function:Probable actin-based motor with a protein kinase activity, similarity:Contains 1 myosin head-like domain, similarity:Contains 1 protein kinase adomain, similarity:Contains 2 IQ domains, similarity:Contains 1 protein kinase adomain, similarity:Contains 2 IQ domains, similarity:Contains 1 protein kinase sedomain, similarity:Contains 2 iprotein kinase activity, similarity:Contains 2 iprotein kinase activity, similarity:Contains 2 iprotein kinase activity. STE Ser/Thr protein kinase activity. Stessue specificity: Expressed in retina, kidney and testis.           Background         This class of myo	Gene Name	MYO3B
Specificity         MYO3B Monoclonal Antibody detects endogenous levels of protein.           Formulation         Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.           Source         Monoclonal, Mouse, IgG           Purification         The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.           Dilution         WB 1:500-2000           Concentration         1 mg/ml           Purity         ≥90%           Storage Stability         -20°C/1 year           Synonyms           Observed Band         147kD           Cell Pathway         Cytoplasm, cytoskeleton. Cell projection, stereocilium.           Tissue Specificity         Expressed in retina, kidney and testis.           Function         catalytic activity:ATP + a protein = ADP + a phosphoprotein, function:Probable actin-based motor with a protein kinase activity, similarity:Contains 1 myosin head-like domain, similarity:Contains 1 protein kinase domain, similarity:Contains 2 IQ domains, similarity:Contains 1 protein kinase domain, similarity:STE Ser/Thr protein kinase family, tissue specificity:Expressed in retina, kidney and testis.           Background         This gene encodes one of the class III myosins. Myosins are ATPases, activated by actin, that move along actin filaments in the cell. This class of myosins are characterized by an amino-terminal kinase domain and shown to be present in photoreceptors. Alternative splicing results in multiple transcript variants encoding	Protein Name	Myosin-IIIb (EC 2.7.11.1)
Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.  Source Monoclonal, Mouse, IgG  Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-2000 Concentration 1 mg/ml  Purity ≥90% Storage Stability -20°C/1 year  Synonyms  Observed Band 147kD Cell Pathway Cytoplasm, cytoskeleton. Cell projection, stereocilium.  Tissue Specificity Expressed in retina, kidney and testis.  Function catalytic activity:ATP + a protein = ADP + a phosphoprotein.,function:Probable actin-based motor with a protein kinase activity.,similarity:Contains 1 myosin head-like domain.,similarity:Contains 2 IQ domains, similarity:Contains 1 protein kinase domain. similarity:Contains 2 IQ domains, similarity:Contains 1 protein kinase domain. similarity:Contains 1 protein kinase family.,tissue specificity:Expressed in retina, kidney and testis.,  Background This gene encodes one of the class III myosins. Myosins are ATPases, activated by actin, that move along actin filaments in the cell. This class of myosins are characterized by an amino-terminal kinase domain and shown to be present in photoreceptors. Alternative splicing results in multiple transcript variants encoding	Immunogen	Synthesized peptide derived from human protein . at AA range: 1030-1110
Source       Monoclonal, Mouse, IgG         Purification       The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.         Dilution       WB 1:500-2000         Concentration       1 mg/ml         Purity       ≥90%         Storage Stability       -20°C/1 year         Synonyms       Observed Band       147kD         Cell Pathway       Cytoplasm, cytoskeleton. Cell projection, stereocilium.         Tissue Specificity       Expressed in retina, kidney and testis.         Function       catalytic activity:ATP + a protein = ADP + a phosphoprotein.,function:Probable actin-based motor with a protein kinase activity, similarity:Contains 1 myosin head-like domain.,similarity:Contains 1 protein kinase domain. similarity:Contains 2 IQ domains, similarity:Contains 1 protein kinase domain. similarity:Expressed in retina, kidney and testis         Background       This gene encodes one of the class III myosins. Myosins are ATPases, activated by actin, that move along actin filaments in the cell. This class of myosins are characterized by an amino-terminal kinase domain and shown to be present in photoreceptors. Alternative splicing results in multiple transcript variants encoding	Specificity	MYO3B Monoclonal Antibody detects endogenous levels of protein.
Purification  The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution  WB 1:500-2000  Concentration  1 mg/ml  Purity  ≥90%  Storage Stability  -20°C/1 year  Synonyms  Observed Band  147kD  Cell Pathway  Cytoplasm, cytoskeleton. Cell projection, stereocilium.  Tissue Specificity  Expressed in retina, kidney and testis.  Function  catalytic activity:ATP + a protein = ADP + a phosphoprotein.,function:Probable actin-based motor with a protein kinase activity, similarity:Contains 1 myosin head-like domain., similarity:Contains 1 protein kinase activity in the N-terminal section; belongs to the protein kinase superfamily. STE Ser/Thr protein kinase family.,tissue specificity:Expressed in retina, kidney and testis.  Background  This gene encodes one of the class III myosins. Myosins are ATPases, activated by actin, that move along actin filaments in the cell. This class of myosins are characterized by an amino-terminal kinase domain and shown to be present in photoreceptors. Alternative splicing results in multiple transcript variants encoding	Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-2000  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms  Observed Band 147kD  Cell Pathway Cytoplasm, cytoskeleton. Cell projection, stereocilium.  Tissue Specificity Expressed in retina, kidney and testis.  Function catalytic activity:ATP + a protein = ADP + a phosphoprotein, function:Probable actin-based motor with a protein kinase activitysimilarity:Contains 1 myosin head-like domain.,similarity:Contains 1 protein kinase domain.,similarity:Contains 2 IQ domains. similarity:In the N-terminal section; belongs to the protein kinase superfamily. STE Ser/Thr protein kinase family.,tissue specificity:Expressed in retina, kidney and testis.  Background This gene encodes one of the class III myosins. Myosins are ATPases, activated by actin, that move along actin filaments in the cell. This class of myosins are characterized by an amino-terminal kinase domain and shown to be present in photoreceptors. Alternative splicing results in multiple transcript variants encoding	Source	Monoclonal, Mouse,IgG
Concentration       1 mg/ml         Purity       ≥90%         Storage Stability       -20°C/1 year         Synonyms         Observed Band       147kD         Cell Pathway       Cytoplasm, cytoskeleton. Cell projection, stereocilium .         Tissue Specificity       Expressed in retina, kidney and testis.         Function       catalytic activity:ATP + a protein = ADP + a phosphoprotein.,function:Probable actin-based motor with a protein kinase activity.,similarity:Contains 1 myosin head-like domain.,similarity:Contains 1 protein kinase domain.,similarity:Contains 2 IQ domains.,similarity:In the N-terminal section; belongs to the protein kinase superfamily. STE Ser/Thr protein kinase family.,tissue specificity:Expressed in retina, kidney and testis.,         Background       This gene encodes one of the class III myosins. Myosins are ATPases, activated by actin, that move along actin filaments in the cell. This class of myosins are characterized by an amino-terminal kinase domain and shown to be present in photoreceptors. Alternative splicing results in multiple transcript variants encoding	Purification	
Purity ≥90%  Storage Stability -20°C/1 year  Synonyms  Observed Band 147kD  Cell Pathway Cytoplasm, cytoskeleton. Cell projection, stereocilium.  Tissue Specificity Expressed in retina, kidney and testis.  Function catalytic activity:ATP + a protein = ADP + a phosphoproteinfunction:Probable actin-based motor with a protein kinase activitysimilarity:Contains 1 myosin head-like domain.,similarity:Contains 1 protein kinase domain.,similarity:Contains 2 IQ domains.s,ismilarity:In the N-terminal section; belongs to the protein kinase superfamily. STE Ser/Thr protein kinase family.,tissue specificity:Expressed in retina, kidney and testis.,  Background This gene encodes one of the class III myosins. Myosins are ATPases, activated by actin, that move along actin filaments in the cell. This class of myosins are characterized by an amino-terminal kinase domain and shown to be present in photoreceptors. Alternative splicing results in multiple transcript variants encoding	Dilution	WB 1:500-2000
Synonyms  Observed Band 147kD  Cell Pathway Cytoplasm, cytoskeleton. Cell projection, stereocilium.  Tissue Specificity Expressed in retina, kidney and testis.  Function catalytic activity:ATP + a protein = ADP + a phosphoprotein.,function:Probable actin-based motor with a protein kinase activity.,similarity:Contains 1 myosin head-like domain.,similarity:Contains 1 protein kinase domain.,similarity:Contains 2 IQ domains.,similarity:In the N-terminal section; belongs to the protein kinase superfamily. STE Ser/Thr protein kinase family.,tissue specificity:Expressed in retina, kidney and testis.,  Background This gene encodes one of the class III myosins. Myosins are ATPases, activated by actin, that move along actin filaments in the cell. This class of myosins are characterized by an amino-terminal kinase domain and shown to be present in photoreceptors. Alternative splicing results in multiple transcript variants encoding	Concentration	1 mg/ml
Synonyms  Observed Band  147kD  Cell Pathway  Cytoplasm, cytoskeleton. Cell projection, stereocilium.  Tissue Specificity  Expressed in retina, kidney and testis.  Function  catalytic activity:ATP + a protein = ADP + a phosphoprotein.,function:Probable actin-based motor with a protein kinase activity.,similarity:Contains 1 myosin head-like domain.,similarity:Contains 1 protein kinase domain.,similarity:Contains 2 IQ domains.,similarity:In the N-terminal section; belongs to the protein kinase superfamily. STE Ser/Thr protein kinase family.,tissue specificity:Expressed in retina, kidney and testis.,  Background  This gene encodes one of the class III myosins. Myosins are ATPases, activated by actin, that move along actin filaments in the cell. This class of myosins are characterized by an amino-terminal kinase domain and shown to be present in photoreceptors. Alternative splicing results in multiple transcript variants encoding	Purity	≥90%
Cell Pathway  Cytoplasm, cytoskeleton. Cell projection, stereocilium.  Expressed in retina, kidney and testis.  Function  catalytic activity:ATP + a protein = ADP + a phosphoprotein.,function:Probable actin-based motor with a protein kinase activity.,similarity:Contains 1 myosin head-like domain.,similarity:Contains 1 protein kinase domain.,similarity:Contains 2 IQ domains.,similarity:In the N-terminal section; belongs to the protein kinase superfamily. STE Ser/Thr protein kinase family.,tissue specificity:Expressed in retina, kidney and testis.,  Background  This gene encodes one of the class III myosins. Myosins are ATPases, activated by actin, that move along actin filaments in the cell. This class of myosins are characterized by an amino-terminal kinase domain and shown to be present in photoreceptors. Alternative splicing results in multiple transcript variants encoding	Storage Stability	-20°C/1 year
Cell Pathway  Cytoplasm, cytoskeleton. Cell projection, stereocilium.  Expressed in retina, kidney and testis.  Function  catalytic activity:ATP + a protein = ADP + a phosphoprotein.,function:Probable actin-based motor with a protein kinase activity.,similarity:Contains 1 myosin head-like domain.,similarity:Contains 1 protein kinase domain.,similarity:Contains 2 IQ domains.,similarity:In the N-terminal section; belongs to the protein kinase superfamily. STE Ser/Thr protein kinase family.,tissue specificity:Expressed in retina, kidney and testis.,  Background  This gene encodes one of the class III myosins. Myosins are ATPases, activated by actin, that move along actin filaments in the cell. This class of myosins are characterized by an amino-terminal kinase domain and shown to be present in photoreceptors. Alternative splicing results in multiple transcript variants encoding	Synonyms	
Tissue Specificity  Expressed in retina, kidney and testis.  Function  catalytic activity:ATP + a protein = ADP + a phosphoprotein.,function:Probable actin-based motor with a protein kinase activity.,similarity:Contains 1 myosin head-like domain.,similarity:Contains 1 protein kinase domain.,similarity:Contains 2 IQ domains.,similarity:In the N-terminal section; belongs to the protein kinase superfamily. STE Ser/Thr protein kinase family.,tissue specificity:Expressed in retina, kidney and testis.,  Background  This gene encodes one of the class III myosins. Myosins are ATPases, activated by actin, that move along actin filaments in the cell. This class of myosins are characterized by an amino-terminal kinase domain and shown to be present in photoreceptors. Alternative splicing results in multiple transcript variants encoding	Observed Band	147kD
catalytic activity:ATP + a protein = ADP + a phosphoprotein.,function:Probable actin-based motor with a protein kinase activity.,similarity:Contains 1 myosin head-like domain.,similarity:Contains 1 protein kinase domain.,similarity:Contains 2 IQ domains.,similarity:In the N-terminal section; belongs to the protein kinase superfamily. STE Ser/Thr protein kinase family.,tissue specificity:Expressed in retina, kidney and testis.,  Background  This gene encodes one of the class III myosins. Myosins are ATPases, activated by actin, that move along actin filaments in the cell. This class of myosins are characterized by an amino-terminal kinase domain and shown to be present in photoreceptors. Alternative splicing results in multiple transcript variants encoding	Cell Pathway	Cytoplasm, cytoskeleton. Cell projection, stereocilium.
actin-based motor with a protein kinase activity.,similarity:Contains 1 myosin head-like domain.,similarity:Contains 1 protein kinase domain.,similarity:Contains 2 IQ domains.,similarity:In the N-terminal section; belongs to the protein kinase superfamily. STE Ser/Thr protein kinase family.,tissue specificity:Expressed in retina, kidney and testis.,  Background  This gene encodes one of the class III myosins. Myosins are ATPases, activated by actin, that move along actin filaments in the cell. This class of myosins are characterized by an amino-terminal kinase domain and shown to be present in photoreceptors. Alternative splicing results in multiple transcript variants encoding	Tissue Specificity	Expressed in retina, kidney and testis.
by actin, that move along actin filaments in the cell. This class of myosins are characterized by an amino-terminal kinase domain and shown to be present in photoreceptors. Alternative splicing results in multiple transcript variants encoding	Function	actin-based motor with a protein kinase activity.,similarity:Contains 1 myosin head-like domain.,similarity:Contains 1 protein kinase domain.,similarity:Contains 2 IQ domains.,similarity:In the N-terminal section; belongs to the protein kinase superfamily. STE Ser/Thr protein kinase family.,tissue specificity:Expressed in
	Background	characterized by an amino-terminal kinase domain and shown to be present in photoreceptors. Alternative splicing results in multiple transcript variants encoding

Nanjing BYabscience technology Co.,Ltd



国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务, 欢迎咨询



matters needing attention

Avoid repeated freezing and thawing!

**Usage suggestions** 

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## **Products Images**

Nanjing BYabscience technology Co.,Ltd