




## NDUB3 Monoclonal Antibody

Catalog No         BYmab-05799           Isotype         IgG           Reactivity         Human;Mouse           Applications         WB           Gene Name         NDUFB3           Protein Name         NADH dehydrogenase [ubiquinone] 1 beta subcomplex subunit 3 (Complex I-B12) (CI-B12) (NADH-ubiquinone oxidoreductase B12 subunit)           Immunogen         Synthesized peptide derived from human protein . at AA range: 40-120           Specificity         NDUB3 Monoclonal Antibody detects endogenous levels of protein.           Formulation         Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.           Source         Monoclonal, Mouse, IgG           Purification         The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.           Dilution         WB 1:500-2000           Concentration         1 mg/ml           Purity         ≥90%           Storage Stability         -20°C/1 year           Synonyms         Observed Band         10kD           Cell Pathway         Mitochondrion inner membrane; Single-pass membrane protein; Matrix side .           Tissue Specificity         Heart, Lung,           Function         function: Accessory subunit of the mitochondrial membrane respiratory chain. NADH dehydrogenase (Complex I), that is believed to be not involved in catalysis. Complex		
Applications  WB  Gene Name  NDUFB3  Protein Name  NADH dehydrogenase [ubiquinone] 1 beta subcomplex subunit 3 (Complex I-B12) (CI-B12) (NADH-ubiquinone oxidoreductase B12 subunit)  Immunogen  Synthesized peptide derived from human protein . at AA range: 40-120  Specificity  NDUB3 Monoclonal Antibody detects endogenous levels of protein.  Formulation  Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.  Source  Monoclonal, Mouse, IgG  Purification  The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution  WB 1:500-2000  Concentration  1 mg/ml  Purity  ≥90%  Storage Stability  -20°C/1 year  Synonyms  Observed Band  10kD  Cell Pathway  Mitochondrion inner membrane ; Single-pass membrane protein ; Matrix side .  Tissue Specificity  Heart, Lung,  function: Accessory subunit of the mitochondrial membrane respiratory chain NADH dehydrogenase (Complex I), that is believed to be not involved in catalysis. Complex I functions in the transfer of electrons from NADH or respiratory chain in The immediate electron acceptor for the enzyme is believed to be ubiquiprione. similarity Belongs to the complex I NDUFB3 subunit family, subunit: Complex I is composed of 45 different subunits.  This gene encodes an accessory subunit of the mitochondrial membrane respiratory chain in the respiratory chain in NADH dehydrogenase (Complex I) is the first enzyme in the electron transport chain of milochondria. This protein localizes to the inner membrane of the mitochondrian as single-pass membrane protein. Mutations in the electron transport chain of milochondria. This protein localizes to the inner membrane of the mitochondrian as a single-pass membrane protein. Mutations in the tensfer as a single-pass membrane protein. Mutations in the electron transport chain of milochondria. This protein localizes to the inner membrane of the mitochondrian as a single-pass membrane protein. Mutations in	Catalog No	BYmab-05799
Applications WB Gene Name NDUFB3 Protein Name   NADH dehydrogenase [ubiquinone] 1 beta subcomplex subunit 3 (Complex I-B12) (CI-B12) (NADH-lubiquinone oxidoreductase B12 subunit) Immunogen   Synthesized peptide derived from human protein . at AA range. 40-120 Specificity   NDUB3 Monoclonal Antibody detects endogenous levels of protein.  Formulation   Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.  Source   Monoclonal, Mouse, IgG  Purification   The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution   WB 1:500-2000  Concentration   1 mg/ml  Purity   ≥90%  Storage Stability   -20°C/1 year  Synonyms  Observed Band   10kD  Cell Pathway   Mitochondrion inner membrane ; Single-pass membrane protein ; Matrix side .  Tissue Specificity   Heart, Lung,  Function   function:Accessory subunit of the mitochondrial membrane respiratory chain NADH dehydrogenase (Complex I), that is believed to be not involved in catalysis. Complex I functions in the transfer of electrons from NADH to the respiratory chain in The immediate electron acceptor for the enzyme is believed to be ubiquinone, similarity-Belongs to the complex I NDUFBS subunit family, subunit:Complex I is composed of 45 different subunits.  Background   This gene encodes an accessory subunit of the mitochondrial membrane respiratory chain in the feasing as a single-pass membrane protein. Mutations in the feasing as a single-pass membrane protein. Mutations in the telectron transport chain of mitochondria. This protein localizes to the inner membrane of the mitochondrian as a single-pass membrane protein. Mutations in	Isotype	IgG
Gene Name         NDUFB3           Protein Name         NADH dehydrogenase [ubiquinone] 1 beta subcomplex subunit 3 (Complex I-B12) (CI-B12) (NADH-ubiquinone oxidoreductase B12 subunit)           Immunogen         Synthesized peptide derived from human protein . at AA range: 40-120           Specificity         NDUB3 Monoclonal Antibody detects endogenous levels of protein.           Formulation         Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.           Source         Monoclonal, Mouse, IgG           Purification         The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.           Dilution         WB 1:500-2000           Concentration         1 mg/ml           Purity         ≥90%           Storage Stability         -20°C/1 year           Synonyms         Observed Band         10kD           Cell Pathway         Mitochondrion inner membrane; Single-pass membrane protein; Matrix side .           Function         function:Accessory subunit of the mitochondrial membrane respiratory chain NADH dehydrogenase (Complex I), that is believed to be not involved in catalysis. Complex I functions in the transfer of electrons from NADH to the respiratory chain. The immediate electron acceptor for the enzyme is believed to be ubiquinonesimilarity:Belongs to the complex I NDUFB3 subunit family, subunit family, subunit Complex I is composed of 45 different subunits.,           Background         This gene encodes an	Reactivity	Human;Mouse
Protein Name  NADH dehydrogenase [ubiquinone] 1 beta subcomplex subunit 3 (Complex I-B12) (CI-B12) (NADH-ubiquinone oxidoreductase B12 subunit)  Immunogen  Synthesized peptide derived from human protein . at AA range: 40-120  Specificity  NDUB3 Monoclonal Antibody detects endogenous levels of protein.  Formulation  Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.  Source  Monoclonal, Mouse, IgG  Purification  The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution  WB 1:500-2000  Concentration  1 mg/ml  Purity  ≥90%  Storage Stability  -20°C/1 year  Synonyms  Observed Band  10kD  Cell Pathway  Mitochondrion inner membrane; Single-pass membrane protein; Matrix side .  Tissue Specificity  Heart, Lung,  Function  function: Accessory subunit of the mitochondrial membrane respiratory chain NADH dehydrogenase (Complex I), that is believed to be not involved in catalysis. Complex I functions in the transfer of electrons from NADH to the respiratory chain. The immediate electron acceptor for the enzyme is believed to be ubiquinone, similarity: Belongs to the complex I NDUFBS subunit family, subunit: Complex I is composed of 45 different subunits.  Background  This gene encodes an accessory subunit of the mitochondrial. This protein localizes to the inner membrane of the mitochondria. This protein localizes to the inner membrane of the mitochondrian as a single-pass membrane protein. Mutations in	Applications	WB
Immunogen         Synthesized peptide derived from human protein . at AA range: 40-120           Specificity         NDUB3 Monoclonal Antibody detects endogenous levels of protein.           Formulation         Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.           Source         Monoclonal, Mouse, IgG           Purification         The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.           Dilution         WB 1:500-2000           Concentration         1 mg/ml           Purity         ≥90%           Storage Stability         -20°C/1 year           Synonyms           Observed Band         10kD           Cell Pathway         Mitochondrion inner membrane; Single-pass membrane protein; Matrix side .           Tissue Specificity         Heart,Lung,           Function         function:Accessory subunit of the mitochondrial membrane respiratory chain NADH dehydrogenase (Complex I), that is believed to be not involved in catalysis. Complex I functions in the transfer of electrons from NADH to the respiratory chain. The immediate electron acceptor for the enzyme is believed to be ublquinone, similarity. Belongs to the complex I NDUFB3 subunit family. subunit:Complex I is composed of 45 different subunits.           Background         This gene encodes an accessory subunit of the mitochondrial membrane respiratory chain NADH dehydrogenase (Complex I) which is the first enzyme in the electron transport chain of mitochondrion as a sin	Gene Name	NDUFB3
Specificity         NDUB3 Monoclonal Antibody detects endogenous levels of protein.           Formulation         Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.           Source         Monoclonal, Mouse, IgG           Purification         The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.           Dilution         WB 1:500-2000           Concentration         1 mg/ml           Purity         ≥90%           Storage Stability         -20°C/1 year           Synonyms         Observed Band         10kD           Cell Pathway         Mitochondrion inner membrane; Single-pass membrane protein; Matrix side.           Tissue Specificity         Heart, Lung,           Function         function: Accessory subunit of the mitochondrial membrane respiratory chain NADH dehydrogenase (Complex I), that is believed to be not involved in catalysis. Complex I functions in the transfer of electrons from NADH to the respiratory chain. The immediate electron acceptor for the enzyme is believed to be ubiquinone, similarity: Belongs to the complex I NDUFB3 subunit family., subunit: Complex I is composed of 45 different subunitis.,           Background         This gene encodes an accessory subunit of the mitochondrial membrane respiratory chain NADH dehydrogenase (Complex I) which is the first enzyme in the electron transport chain of mitochondrial or as a single-pass membrane protein. Mutatitons in the membrane protein. Mutatitons in the mitochondrial membrane respiratory chain femilic chain of mito	Protein Name	NADH dehydrogenase [ubiquinone] 1 beta subcomplex subunit 3 (Complex I-B12) (CI-B12) (NADH-ubiquinone oxidoreductase B12 subunit)
Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.  Source Monoclonal, Mouse,IgG  Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-2000  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms  Observed Band 10kD  Cell Pathway Mitochondrion inner membrane ; Single-pass membrane protein ; Matrix side .  Tissue Specificity Heart,Lung,  Function function:Accessory subunit of the mitochondrial membrane respiratory chain NADH dehydrogenase (Complex I), that is believed to be not involved in catalysis. Complex I functions in the transfer of electrons from NADH to the respiratory chain. The immediate electron acceptor for the enzyme is believed to be ubiquinone, similarity:Belongs to the complex I NDUFB3 subunit family, subunit:Complex I is composed of 45 different subunits.,  Background This gene encodes an accessory subunit of the mitochondrial membrane respiratory chain NADH dehydrogenase (Complex I) which is the first enzyme in the electron transport chain of mitochondria. This protein localizes to the inner membrane of the mitochondrial membrane protein. Mutations in	Immunogen	Synthesized peptide derived from human protein . at AA range: 40-120
Source         Monoclonal, Mouse,IgG           Purification         The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.           Dilution         WB 1:500-2000           Concentration         1 mg/ml           Purity         ≥90%           Storage Stability         -20°C/1 year           Synonyms         Observed Band         10kD           Cell Pathway         Mitochondrion inner membrane; Single-pass membrane protein; Matrix side.           Tissue Specificity         Heart,Lung,           Function         function:Accessory subunit of the mitochondrial membrane respiratory chain NADH dehydrogenase (Complex I), that is believed to be not involved in catalysis complex I functions in the transfer of electrons from NADH to the respiratory chain. The immediate electron acceptor for the enzyme is believed to be ubiquinone.,similarity:Belongs to the complex I NDUFB3 subunit family.,subunit:Complex I is composed of 45 different subunits.,           Background         This gene encodes an accessory subunit of the mitochondrial membrane respiratory chain NADH dehydrogenase (Complex I) which is the first enzyme in the electron transport chain of mitochondria. This protein localizes to the inner membrane of the mitochondrion as a single-pass membrane protein. Mutations in	Specificity	NDUB3 Monoclonal Antibody detects endogenous levels of protein.
Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year  Synonyms Observed Band 10kD Cell Pathway Mitochondrion inner membrane ; Single-pass membrane protein ; Matrix side .  Tissue Specificity Heart,Lung, Function function:Accessory subunit of the mitochondrial membrane respiratory chain NADH dehydrogenase (Complex I), that is believed to be not involved in catalysis. Complex I functions in the transfer of electrons from NADH to the respiratory chain. The immediate electron acceptor for the enzyme is believed to be ubiquinone, similarity:Belongs to the complex I NDUFB3 subunit family, subunit:Complex I is composed of 45 different subunits.  Background This gene encodes an accessory subunit of the mitochondrial membrane respiratory chain NADH dehydrogenase (Complex I) which is the first enzyme in the electron transport chain of mitochondria. This protein localizes to the inner membrane of the mitochondrion as a single-pass membrane protein. Mutations in	Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-2000  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms  Observed Band 10kD  Cell Pathway Mitochondrion inner membrane ; Single-pass membrane protein ; Matrix side .  Tissue Specificity Heart, Lung,  Function function: Accessory subunit of the mitochondrial membrane respiratory chain NADH dehydrogenase (Complex I), that is believed to be not involved in catalysis. Complex I functions in the transfer of electrons from NADH to the respiratory chain. The immediate electron acceptor for the enzyme is believed to be ubiquinone., similarity: Belongs to the complex I NDUFB3 subunit family, subunit: Complex I is composed of 45 different subunits.,  Background This gene encodes an accessory subunit of the mitochondrial membrane respiratory chain NADH dehydrogenase (Complex I) which is the first enzyme in the electron transport chain of mitochondria. This protein localizes to the inner membrane of the mitochondrion as a single-pass membrane protein. Mutations in	Source	Monoclonal, Mouse,IgG
Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms  Observed Band 10kD  Cell Pathway Mitochondrion inner membrane; Single-pass membrane protein; Matrix side.  Tissue Specificity Heart,Lung,  Function function:Accessory subunit of the mitochondrial membrane respiratory chain NADH dehydrogenase (Complex I), that is believed to be not involved in catalysis. Complex I functions in the transfer of electrons from NADH to the respiratory chain. The immediate electron acceptor for the enzyme is believed to be ubiquinone.,similarity:Belongs to the complex I NDUFB3 subunit family.,subunit:Complex I is composed of 45 different subunits.,  Background This gene encodes an accessory subunit of the mitochondrial membrane respiratory chain NADH dehydrogenase (Complex I) which is the first enzyme in the electron transport chain of mitochondria. This protein localizes to the inner membrane of the mitochondrion as a single-pass membrane protein. Mutations in	Purification	, , , , , , , , , , , , , , , , , , , ,
Purity ≥90%  Storage Stability -20°C/1 year  Synonyms  Observed Band 10kD  Cell Pathway Mitochondrion inner membrane; Single-pass membrane protein; Matrix side.  Tissue Specificity Heart,Lung,  Function function:Accessory subunit of the mitochondrial membrane respiratory chain NADH dehydrogenase (Complex I), that is believed to be not involved in catalysis. Complex I functions in the transfer of electrons from NADH to the respiratory chain. The immediate electron acceptor for the enzyme is believed to be ubiquinone, similarity:Belongs to the complex I NDUFB3 subunit family.,subunit:Complex I is composed of 45 different subunits.,  Background This gene encodes an accessory subunit of the mitochondrial membrane respiratory chain NADH dehydrogenase (Complex I) which is the first enzyme in the electron transport chain of mitochondria. This protein localizes to the inner membrane of the mitochondrion as a single-pass membrane protein. Mutations in	Dilution	WB 1:500-2000
Synonyms  Observed Band 10kD  Cell Pathway Mitochondrion inner membrane; Single-pass membrane protein; Matrix side.  Tissue Specificity Heart, Lung,  Function function: Accessory subunit of the mitochondrial membrane respiratory chain NADH dehydrogenase (Complex I), that is believed to be not involved in catalysis. Complex I functions in the transfer of electrons from NADH to the respiratory chain. The immediate electron acceptor for the enzyme is believed to be ubiquinone., similarity: Belongs to the complex I NDUFB3 subunit family., subunit: Complex I is composed of 45 different subunits.,  Background This gene encodes an accessory subunit of the mitochondrial membrane respiratory chain NADH dehydrogenase (Complex I) which is the first enzyme in the electron transport chain of mitochondria. This protein localizes to the inner membrane of the mitochondrion as a single-pass membrane protein. Mutations in	Concentration	1 mg/ml
Synonyms  Observed Band  10kD  Cell Pathway  Mitochondrion inner membrane; Single-pass membrane protein; Matrix side.  Tissue Specificity  Heart,Lung,  function: Accessory subunit of the mitochondrial membrane respiratory chain NADH dehydrogenase (Complex I), that is believed to be not involved in catalysis. Complex I functions in the transfer of electrons from NADH to the respiratory chain. The immediate electron acceptor for the enzyme is believed to be ubiquinone.,similarity:Belongs to the complex I NDUFB3 subunit family.,subunit:Complex I is composed of 45 different subunits.,  Background  This gene encodes an accessory subunit of the mitochondrial membrane respiratory chain NADH dehydrogenase (Complex I) which is the first enzyme in the electron transport chain of mitochondria. This protein localizes to the inner membrane of the mitochondrion as a single-pass membrane protein. Mutations in	Purity	≥90%
Observed Band  Cell Pathway  Mitochondrion inner membrane; Single-pass membrane protein; Matrix side.  Tissue Specificity  Heart,Lung,  function:Accessory subunit of the mitochondrial membrane respiratory chain NADH dehydrogenase (Complex I), that is believed to be not involved in catalysis. Complex I functions in the transfer of electrons from NADH to the respiratory chain. The immediate electron acceptor for the enzyme is believed to be ubiquinone.,similarity:Belongs to the complex I NDUFB3 subunit family.,subunit:Complex I is composed of 45 different subunits.,  Background  This gene encodes an accessory subunit of the mitochondrial membrane respiratory chain NADH dehydrogenase (Complex I) which is the first enzyme in the electron transport chain of mitochondria. This protein localizes to the inner membrane of the mitochondrion as a single-pass membrane protein. Mutations in	Storage Stability	-20°C/1 year
Cell Pathway         Mitochondrion inner membrane ; Single-pass membrane protein ; Matrix side .           Tissue Specificity         Heart,Lung,           Function         function:Accessory subunit of the mitochondrial membrane respiratory chain NADH dehydrogenase (Complex I), that is believed to be not involved in catalysis. Complex I functions in the transfer of electrons from NADH to the respiratory chain. The immediate electron acceptor for the enzyme is believed to be ubiquinone.,similarity:Belongs to the complex I NDUFB3 subunit family.,subunit:Complex I is composed of 45 different subunits.,           Background         This gene encodes an accessory subunit of the mitochondrial membrane respiratory chain NADH dehydrogenase (Complex I) which is the first enzyme in the electron transport chain of mitochondria. This protein localizes to the inner membrane of the mitochondrion as a single-pass membrane protein. Mutations in	Synonyms	
Tissue Specificity  Heart,Lung,  function:Accessory subunit of the mitochondrial membrane respiratory chain NADH dehydrogenase (Complex I), that is believed to be not involved in catalysis. Complex I functions in the transfer of electrons from NADH to the respiratory chain. The immediate electron acceptor for the enzyme is believed to be ubiquinone., similarity:Belongs to the complex I NDUFB3 subunit family., subunit:Complex I is composed of 45 different subunits.,  This gene encodes an accessory subunit of the mitochondrial membrane respiratory chain NADH dehydrogenase (Complex I) which is the first enzyme in the electron transport chain of mitochondria. This protein localizes to the inner membrane of the mitochondrion as a single-pass membrane protein. Mutations in	Observed Band	401-0
Function  function:Accessory subunit of the mitochondrial membrane respiratory chain NADH dehydrogenase (Complex I), that is believed to be not involved in catalysis. Complex I functions in the transfer of electrons from NADH to the respiratory chain. The immediate electron acceptor for the enzyme is believed to be ubiquinone.,similarity:Belongs to the complex I NDUFB3 subunit family.,subunit:Complex I is composed of 45 different subunits.,  This gene encodes an accessory subunit of the mitochondrial membrane respiratory chain NADH dehydrogenase (Complex I) which is the first enzyme in the electron transport chain of mitochondria. This protein localizes to the inner membrane of the mitochondrion as a single-pass membrane protein. Mutations in		TUKD
NADH dehydrogenase (Complex I), that is believed to be not involved in catalysis. Complex I functions in the transfer of electrons from NADH to the respiratory chain. The immediate electron acceptor for the enzyme is believed to be ubiquinone., similarity:Belongs to the complex I NDUFB3 subunit family., subunit:Complex I is composed of 45 different subunits.,  This gene encodes an accessory subunit of the mitochondrial membrane respiratory chain NADH dehydrogenase (Complex I) which is the first enzyme in the electron transport chain of mitochondria. This protein localizes to the inner membrane of the mitochondrion as a single-pass membrane protein. Mutations in	Cell Pathway	
respiratory chain NADH dehydrogenase (Complex I) which is the first enzyme in the electron transport chain of mitochondria. This protein localizes to the inner membrane of the mitochondrion as a single-pass membrane protein. Mutations in	-	Mitochondrion inner membrane ; Single-pass membrane protein ; Matrix side .
	Tissue Specificity	Mitochondrion inner membrane; Single-pass membrane protein; Matrix side.  Heart,Lung,  function:Accessory subunit of the mitochondrial membrane respiratory chain NADH dehydrogenase (Complex I), that is believed to be not involved in catalysis. Complex I functions in the transfer of electrons from NADH to the respiratory chain. The immediate electron acceptor for the enzyme is believed to be ubiquinone.,similarity:Belongs to the complex I NDUFB3 subunit

Nanjing BYabscience technology Co.,Ltd


网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658







	results in multiple transcript variants encoding the same protein. Humans have multiple pseudogenes of this gene. [provided by RefSeq, Mar 2012],
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.



网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658