



## NASP Monoclonal Antibody

Catalog No	BYmab-05857
Isotype	lgG
Reactivity	Human;Rat;Mouse
Applications	WB
Gene Name	NASP
Protein Name	Nuclear autoantigenic sperm protein (NASP)
Immunogen	Synthesized peptide derived from human protein . at AA range: 680-760
Specificity	NASP Monoclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	86kD
Cell Pathway	Cytoplasm . Nucleus .
Tissue Specificity	Isoform 1 is testis- and sperm-specific.
Function	function:Required for DNA replication, normal cell cycle progression and cell proliferation. Forms a cytoplasmic complex with HSP90 and H1 linker histones and stimulates HSP90 ATPase activity. NASP and H1 histone are subsequently released from the complex and translocate to the nucleus where the histone is released for binding to DNA.,PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Belongs to the NASP family.,similarity:Contains 3 TPR repeats.,subunit:Binds to linker H1 histones but not to core histones. Also binds to HSP90 in the cytoplasm. This interaction stimulates binding of NASP to HIST1H1T/H1T.,tissue specificity:Isoform 1 is testis- and sperm-specific.,
Background	This gene encodes a H1 histone binding protein that is involved in transporting histones into the nucleus of dividing cells. Multiple isoforms are encoded by
	Naniing Byahscience technology Co. Ltd

## Nanjing BYabscience technology Co.,Ltd



国内优质抗体供应商

精准的 WB 检测服务 431 -----

24H 在线服务,欢迎咨询



	transcript variants of this gene. The somatic form is expressed in all mitotic cells, is localized to the nucleus, and is coupled to the cell cycle. The testicular form is expressed in embryonic tissues, tumor cells, and the testis. In male germ cells, this protein is localized to the cytoplasm of primary spermatocytes, the nucleus of spermatids, and the periacrosomal region of mature spermatozoa. [provided by RefSeq, Jul 2008],	
matters needing attention	Avoid repeated freezing and thawing!	
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.	
Products Images		

Nanjing BYabscience technology Co.,Ltd