




NUFP1 Monoclonal Antibody

Catalog No BYmab-05861 Isotype IgG Reactivity Human;Rat;Mouse Applications WB Gene Name NUFIP1 Protein Name Nuclear fragile X mental retardation-interacting protein 1 (Nuclear FMRP-interacting protein 1) Immunogen Synthesized peptide derived from human protein . at AA range: 180-260 Specificity NUFP1 Monoclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 54kD Cell Pathway Nucleus . Distributed in the nucleus in a dot-like pattern. Tissue Specificity Expressed in spleen, thymus, prostate, testis, ovary, small intestine, colon, peripheral blood leukocyte, heart, brain, placenta, lung, liver, skeletal muscle, kidney, and pancreas. Function Incolona Binds RNA, similarity:Contains 1		
Reactivity Human;Rat;Mouse Applications WB Gene Name NUFIP1 Protein Name Nuclear fragile X mental retardation-interacting protein 1 (Nuclear FMRP-interacting protein 1) Immunogen Synthesized peptide derived from human protein . at AA range: 180-260 Specificity NUFP1 Monoclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 54kD Cell Pathway Nucleus . Distributed in the nucleus in a dot-like pattern. Tissue Specificity Expressed in spleen, thymus, prostate, testis, ovary, small intestine, colon, peripheral blood leukocyte, heart, brain, placenta, lung, liver, skeletal muscle, kidney, and pancreas. Function function:Binds RNA, similarity:Contains 1 C2H2-type zinc finger, subcollar location:Distributed in the nucleus in a dot-like pattern, subunit-Interacts with FMR1, tissue specificity:	Catalog No	BYmab-05861
Applications WB Gene Name NUFIP1 Protein Name FMRP-interacting protein 1 (Nuclear FMRP-interacting protein 1) Immunogen Synthesized peptide derived from human protein . at AA range: 180-260 Specificity NUFP1 Monoclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 54kD Cell Pathway Nucleus . Distributed in the nucleus in a dot-like pattern. Tissue Specificity Expressed in spleen, thymus, prostate, testis, ovary, small intestine, colon, peripheral blood leukocyte, heart, brain, placenta, lung, liver, skeletal muscle, kidney, and pancreas. Function function:Binds RNA, similarity:Contains 1 C2H2-type zinc finger, subcellular location:Distributed in the nucleus in a dot-like pattern. subunit Interacts with FMR1, tissue specificity:Expressed in spleen, thymus, prostate, testis, ovary, small intestine, colon, peripheral blood leukocyte, heart, brain, placenta, lung, liver, skeletal muscle, kidney, and pancreas. Function function:Binds RNA, similarity:Contains 1 C2H2-type zinc finger, subcellular location:Distributed in the nucleus in a dot-like pattern, subunit Interacts with FMR1, tissue specificity:Expressed in spleen, thymus, prostate, testis, ovary, small intestine, colon, peripheral blood leukocyte, heart, brain, placenta, lung, liver, skeletal muscle, kidney, and pancreas. Background This gene encodes a nuclear RNA binding protein that contains a C2H2 zinc finger motif and a nuclear localization signal. This protein is associated with the nuclear matrix in perichromatin fibrils and, in neurons, localizes to the cytoplasm in association with endoplasmic retriculum ribosomes. This protein interacts with	Isotype	IgG
Gene Name NUFIP1 Protein Name Nuclear fragile X mental retardation-interacting protein 1 (Nuclear FMRP-interacting protein 1) Immunogen Synthesized peptide derived from human protein . at AA range: 180-260 Specificity NUFP1 Monoclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 54kD Cell Pathway Nucleus . Distributed in the nucleus in a dot-like pattern. Tissue Specificity Expressed in spleen, thymus, prostate, testis, ovary, small intestine, colon, peripheral blood leukocyte, heart, brain, placenta, lung, liver, skeletal muscle, kidney, and pancreas. Function function: Binds RNA., similarity. Contains 1 C2H2-type zinc finger, subcellular location: Distributed in the nucleus in a dot-like pattern, subunit: Interacts with FMR1. tissue specificity: Expressed in spleen, thymus, prostate, testis, ovary, small intestine, colon, peripheral blood leukocyte, heart, brain, placenta, lung, live	Reactivity	Human;Rat;Mouse
Protein Name Nuclear fragile X mental retardation-interacting protein 1 (Nuclear FMRP-interacting protein 1) Immunogen Synthesized peptide derived from human protein . at AA range: 180-260 Specificity NUFP1 Monoclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 54kD Cell Pathway Nucleus . Distributed in the nucleus in a dot-like pattern. Tissue Specificity Expressed in spleen, thymus, prostate, testis, ovary, small intestine, colon, peripheral blood leukocyte, heart, brain, placenta, lung, liver, skeletal muscle, kidney, and pancreas. Function function:Binds RNA, similarity:Contains 1 C2H2-type zinc finger, subcellular location:Distributed in the nucleus in a dot-like pattern, subunit:Interacts with FMR1, tissue specificity:Expressed in spleen, thymus, prostate, testis, ovary, small intestine, colon, peripheral blood leukocyte, heart, brain, placenta, lung, liver, skeletal muscle, kidney, and pancreas. <t< th=""><th>Applications</th><th>WB</th></t<>	Applications	WB
Immunogen Synthesized peptide derived from human protein . at AA range: 180-260	Gene Name	NUFIP1
Specificity NUFP1 Monoclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 54kD Cell Pathway Nucleus . Distributed in the nucleus in a dot-like pattern. Tissue Specificity Expressed in spleen, thymus, prostate, testis, ovary, small intestine, colon, peripheral blood leukocyte, heart, brain, placenta, lung, liver, skeletal muscle, kidney, and pancreas. Function function:Binds RNA., similarity:Contains 1 C2H2-type zinc finger., subcellular location:Distributed in the nucleus in a dot-like pattern, subunit:Interacts with FMR1, tissue specificity:Expressed in spleen, thymus, prostate, testis, ovary, small intestine, colon, peripheral blood leukocyte, heart, brain, placenta, lung, liver, skeletal muscle, kidney, and pancreas. Background This gene encodes a nuclear RNA binding protein that contains a C2H2 zinc finger motif and a nuclear localization signal. This protein is associated with the nucleur matrix in perichromatin fibrils and, in neurons, localizes to the cytoplasm in	Protein Name	Nuclear fragile X mental retardation-interacting protein 1 (Nuclear FMRP-interacting protein 1)
Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 54kD Cell Pathway Nucleus . Distributed in the nucleus in a dot-like pattern. Tissue Specificity Expressed in spleen, thymus, prostate, testis, ovary, small intestine, colon, peripheral blood leukocyte, heart, brain, placenta, lung, liver, skeletal muscle, kidney, and pancreas. Function function:Binds RNA, similarity:Contains 1 C2H2-type zinc finger., subcellular location:Distributed in the nucleus in a dot-like pattern, subunit: Interacts with FMR1, tissue specificity: Expressed in spleen, thymus, prostate, testis, ovary, small intestine, colon, peripheral blood leukocyte, heart, brain, placenta, lung, liver, skeletal muscle, kidney, and pancreas. Background This gene encodes a nuclear RNA binding protein that contains a C2H2 zinc finger motif and a nuclear localization signal. This protein is associated with the nuclear matrix in perichromatin fibrils and, in neurons, localizes to the cytoplasmin association with endoplasmic reticulum ribosomes. This protein interacts with	Immunogen	Synthesized peptide derived from human protein . at AA range: 180-260
Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 54kD Cell Pathway Nucleus . Distributed in the nucleus in a dot-like pattern. Tissue Specificity Expressed in spleen, thymus, prostate, testis, ovary, small intestine, colon, peripheral blood leukocyte, heart, brain, placenta, lung, liver, skeletal muscle, kidney, and pancreas. Function function:Binds RNA, similarity:Contains 1 C2H2-type zinc finger, subcellular location:Distributed in the nucleus in a dot-like pattern, subunit:Interacts with FMR1, tissue specificity:Expressed in spleen, thymus, prostate, testis, ovary, small intestine, colon, peripheral blood leukocyte, heart, brain, placenta, lung, liver, skeletal muscle, kidney, and pancreas. Background This gene encodes a nuclear RNA binding protein that contains a C2H2 zinc finger motif and a nuclear localization signal. This protein is associated with the nuclear matrix in perichromatin fibrils and, in neurons, localizes to the cytoplasm in association with endoplasmic reticulum ribosomes. This protein interacts with	Specificity	NUFP1 Monoclonal Antibody detects endogenous levels of protein.
Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 54kD Cell Pathway Nucleus . Distributed in the nucleus in a dot-like pattern. Tissue Specificity Expressed in spleen, thymus, prostate, testis, ovary, small intestine, colon, peripheral blood leukocyte, heart, brain, placenta, lung, liver, skeletal muscle, kidney, and pancreas. Function function:Binds RNA., similarity:Contains 1 C2H2-type zinc finger., subcellular location:Distributed in the nucleus in a dot-like pattern., subunit:Interacts with FMR1., tissue specificity:Expressed in spleen, thymus, prostate, testis, ovary, small intestine, colon, peripheral blood leukocyte, heart, brain, placenta, lung, liver, skeletal muscle, kidney, and pancreas. Background This gene encodes a nuclear RNA binding protein that contains a C2H2 zinc finger motif and a nuclear localization signal. This protein is associated with the nuclear matrix in perichromatin fibrils and, in neurons, localizes to the cytoplasm in association with endoplasmic reticulum ribosomes. This protein interacts with	Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 54kD Cell Pathway Nucleus . Distributed in the nucleus in a dot-like pattern. Tissue Specificity Expressed in spleen, thymus, prostate, testis, ovary, small intestine, colon, peripheral blood leukocyte, heart, brain, placenta, lung, liver, skeletal muscle, kidney, and pancreas. Function function:Binds RNA., similarity:Contains 1 C2H2-type zinc finger., subcellular location:Distributed in the nucleus in a dot-like pattern., subunit:Interacts with FMR1., tissue specificity:Expressed in spleen, thymus, prostate, testis, ovary, small intestine, colon, peripheral blood leukocyte, heart, brain, placenta, lung, liver, skeletal muscle, kidney, and pancreas. Background This gene encodes a nuclear RNA binding protein that contains a C2H2 zinc finger motif and a nuclear localization signal. This protein is associated with the nuclear matrix in perichromatin fibrils and, in neurons, localizes to the cytoplasm in association with endoplasmic reticulum ribosomes. This protein interacts with	Source	Monoclonal, Mouse,IgG
Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 54kD Cell Pathway Nucleus . Distributed in the nucleus in a dot-like pattern. Tissue Specificity Expressed in spleen, thymus, prostate, testis, ovary, small intestine, colon, peripheral blood leukocyte, heart, brain, placenta, lung, liver, skeletal muscle, kidney, and pancreas. Function function:Binds RNA.,similarity:Contains 1 C2H2-type zinc finger.,subcellular location:Distributed in the nucleus in a dot-like pattern.,subunit:Interacts with FMR1, tissue specificity:Expressed in spleen, thymus, prostate, testis, ovary, small intestine, colon, peripheral blood leukocyte, heart, brain, placenta, lung, liver, skeletal muscle, kidney, and pancreas. Background This gene encodes a nuclear RNA binding protein that contains a C2H2 zinc finger motif and a nuclear localization signal. This protein is associated with the nuclear matrix in perichromatin fibrils and, in neurons, localizes to the cytoplasm in association with endoplasmic reticulum ribosomes. This protein interacts with	Purification	·
Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 54kD Cell Pathway Nucleus . Distributed in the nucleus in a dot-like pattern. Tissue Specificity Expressed in spleen, thymus, prostate, testis, ovary, small intestine, colon, peripheral blood leukocyte, heart, brain, placenta, lung, liver, skeletal muscle, kidney, and pancreas. Function function:Binds RNA.,similarity:Contains 1 C2H2-type zinc finger.,subcellular location:Distributed in the nucleus in a dot-like pattern.,subunit:Interacts with FMR1.,tissue specificity:Expressed in spleen, thymus, prostate, testis, ovary, small intestine, colon, peripheral blood leukocyte, heart, brain, placenta, lung, liver, skeletal muscle, kidney, and pancreas., Background This gene encodes a nuclear RNA binding protein that contains a C2H2 zinc finger motif and a nuclear localization signal. This protein is associated with the nuclear matrix in perichromatin fibrils and, in neurons, localizes to the cytoplasm in association with endoplasmic reticulum ribosomes. This protein interacts with	Dilution	WB 1:500-2000
Synonyms Observed Band Cell Pathway Nucleus . Distributed in the nucleus in a dot-like pattern. Expressed in spleen, thymus, prostate, testis, ovary, small intestine, colon, peripheral blood leukocyte, heart, brain, placenta, lung, liver, skeletal muscle, kidney, and pancreas. Function function:Binds RNA.,similarity:Contains 1 C2H2-type zinc finger.,subcellular location:Distributed in the nucleus in a dot-like pattern.,subunit:Interacts with FMR1.,tissue specificity:Expressed in spleen, thymus, prostate, testis, ovary, small intestine, colon, peripheral blood leukocyte, heart, brain, placenta, lung, liver, skeletal muscle, kidney, and pancreas. Background This gene encodes a nuclear RNA binding protein that contains a C2H2 zinc finger motif and a nuclear localization signal. This protein is associated with the nuclear matrix in perichromatin fibrils and, in neurons, localizes to the cytoplasm in association with endoplasmic reticulum ribosomes. This protein interacts with	Concentration	1 mg/ml
Synonyms Observed Band 54kD Cell Pathway Nucleus . Distributed in the nucleus in a dot-like pattern. Tissue Specificity Expressed in spleen, thymus, prostate, testis, ovary, small intestine, colon, peripheral blood leukocyte, heart, brain, placenta, lung, liver, skeletal muscle, kidney, and pancreas. Function function:Binds RNA.,similarity:Contains 1 C2H2-type zinc finger.,subcellular location:Distributed in the nucleus in a dot-like pattern.,subunit:Interacts with FMR1.,tissue specificity:Expressed in spleen, thymus, prostate, testis, ovary, small intestine, colon, peripheral blood leukocyte, heart, brain, placenta, lung, liver, skeletal muscle, kidney, and pancreas., Background This gene encodes a nuclear RNA binding protein that contains a C2H2 zinc finger motif and a nuclear localization signal. This protein is associated with the nuclear matrix in perichromatin fibrils and, in neurons, localizes to the cytoplasm in association with endoplasmic reticulum ribosomes. This protein interacts with	Purity	≥90%
Observed Band 54kD Cell Pathway Nucleus . Distributed in the nucleus in a dot-like pattern. Tissue Specificity Expressed in spleen, thymus, prostate, testis, ovary, small intestine, colon, peripheral blood leukocyte, heart, brain, placenta, lung, liver, skeletal muscle, kidney, and pancreas. Function function:Binds RNA.,similarity:Contains 1 C2H2-type zinc finger.,subcellular location:Distributed in the nucleus in a dot-like pattern.,subunit:Interacts with FMR1.,tissue specificity:Expressed in spleen, thymus, prostate, testis, ovary, small intestine, colon, peripheral blood leukocyte, heart, brain, placenta, lung, liver, skeletal muscle, kidney, and pancreas. Background This gene encodes a nuclear RNA binding protein that contains a C2H2 zinc finger motif and a nuclear localization signal. This protein is associated with the nuclear matrix in perichromatin fibrils and, in neurons, localizes to the cytoplasm in association with endoplasmic reticulum ribosomes. This protein interacts with	Storage Stability	-20°C/1 year
Cell Pathway Nucleus . Distributed in the nucleus in a dot-like pattern. Tissue Specificity Expressed in spleen, thymus, prostate, testis, ovary, small intestine, colon, peripheral blood leukocyte, heart, brain, placenta, lung, liver, skeletal muscle, kidney, and pancreas. Function function:Binds RNA.,similarity:Contains 1 C2H2-type zinc finger.,subcellular location:Distributed in the nucleus in a dot-like pattern.,subunit:Interacts with FMR1.,tissue specificity:Expressed in spleen, thymus, prostate, testis, ovary, small intestine, colon, peripheral blood leukocyte, heart, brain, placenta, lung, liver, skeletal muscle, kidney, and pancreas., Background This gene encodes a nuclear RNA binding protein that contains a C2H2 zinc finger motif and a nuclear localization signal. This protein is associated with the nuclear matrix in perichromatin fibrils and, in neurons, localizes to the cytoplasm in association with endoplasmic reticulum ribosomes. This protein interacts with	Synonyms	
Tissue Specificity Expressed in spleen, thymus, prostate, testis, ovary, small intestine, colon, peripheral blood leukocyte, heart, brain, placenta, lung, liver, skeletal muscle, kidney, and pancreas. Function function:Binds RNA.,similarity:Contains 1 C2H2-type zinc finger.,subcellular location:Distributed in the nucleus in a dot-like pattern.,subunit:Interacts with FMR1.,tissue specificity:Expressed in spleen, thymus, prostate, testis, ovary, small intestine, colon, peripheral blood leukocyte, heart, brain, placenta, lung, liver, skeletal muscle, kidney, and pancreas., This gene encodes a nuclear RNA binding protein that contains a C2H2 zinc finger motif and a nuclear localization signal. This protein is associated with the nuclear matrix in perichromatin fibrils and, in neurons, localizes to the cytoplasm in association with endoplasmic reticulum ribosomes. This protein interacts with	Observed Band	54kD
peripheral blood leukocyte, heart, brain, placenta, lung, liver, skeletal muscle, kidney, and pancreas. Function function:Binds RNA.,similarity:Contains 1 C2H2-type zinc finger.,subcellular location:Distributed in the nucleus in a dot-like pattern.,subunit:Interacts with FMR1.,tissue specificity:Expressed in spleen, thymus, prostate, testis, ovary, small intestine, colon, peripheral blood leukocyte, heart, brain, placenta, lung, liver, skeletal muscle, kidney, and pancreas., This gene encodes a nuclear RNA binding protein that contains a C2H2 zinc finger motif and a nuclear localization signal. This protein is associated with the nuclear matrix in perichromatin fibrils and, in neurons, localizes to the cytoplasm in association with endoplasmic reticulum ribosomes. This protein interacts with	Cell Pathway	N. J. British I. H. J. J. J. H. H.
location:Distributed in the nucleus in a dot-like pattern.,subunit:Interacts with FMR1.,tissue specificity:Expressed in spleen, thymus, prostate, testis, ovary, small intestine, colon, peripheral blood leukocyte, heart, brain, placenta, lung, liver, skeletal muscle, kidney, and pancreas., This gene encodes a nuclear RNA binding protein that contains a C2H2 zinc finger motif and a nuclear localization signal. This protein is associated with the nuclear matrix in perichromatin fibrils and, in neurons, localizes to the cytoplasm in association with endoplasmic reticulum ribosomes. This protein interacts with		Nucleus . Distributed in the nucleus in a dot-like pattern.
finger motif and a nuclear localization signal. This protein is associated with the nuclear matrix in perichromatin fibrils and, in neurons, localizes to the cytoplasm in association with endoplasmic reticulum ribosomes. This protein interacts with		Expressed in spleen, thymus, prostate, testis, ovary, small intestine, colon, peripheral blood leukocyte, heart, brain, placenta, lung, liver, skeletal muscle,
	Tissue Specificity	Expressed in spleen, thymus, prostate, testis, ovary, small intestine, colon, peripheral blood leukocyte, heart, brain, placenta, lung, liver, skeletal muscle, kidney, and pancreas. function:Binds RNA.,similarity:Contains 1 C2H2-type zinc finger.,subcellular location:Distributed in the nucleus in a dot-like pattern.,subunit:Interacts with FMR1.,tissue specificity:Expressed in spleen, thymus, prostate, testis, ovary, small intestine, colon, peripheral blood leukocyte, heart, brain, placenta, lung,

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658







	BRCA1, upregulates RNA polymerase II transcription, and is involved in box C/D snoRNP biogenesis. A pseudogene of this gene resides on chromosome 6q12. [provided by RefSeq, Feb 2012],
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658