



HCN1 Monoclonal Antibody

Catalog No	BYmab-05961
Isotype	IgG
Reactivity	Human;Rat;Mouse
Applications	WB
Gene Name	HCN1 BCNG1
Protein Name	Potassium/sodium hyperpolarization-activated cyclic nucleotide-gated channel 1 (Brain cyclic nucleotide-gated channel 1) (BCNG-1)
Immunogen	Synthesized peptide derived from human protein . at AA range: 380-460
Specificity	HCN1 Monoclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	97kD
Cell Pathway	Cell membrane ; Multi-pass membrane protein .
Tissue Specificity	Detected in brain, in particular in amygdala and hippocampus, while expression in caudate nucleus, corpus callosum, substantia nigra, subthalamic nucleus and thalamus is very low or not detectable. Detected at very low levels in muscle and pancreas.
Function	domain:The segment S4 is probably the voltage-sensor and is characterized by a series of positively charged amino acids at every third position.,function:Hyperpolarization-activated ion channel exhibiting weak selectivity for potassium over sodium ions. Contributes to the native pacemaker currents in heart (If) and in neurons (Ih). Activated by cAMP, and at 10-100 times higher concentrations, also by cGMP. May mediate responses to sour stimuli.,sequence caution:Contaminating sequence. Potential poly-A sequence.,similarity:Belongs to the potassium channel HCN family.,similarity:Contains 1 cyclic nucleotide-binding domain.,subunit:The

Nanjing BYabscience technology Co.,Ltd



potassium channel is probably composed of a homo- or heterotetrameric complex of pore-forming subunits. Heteromultimer with HCN2 (By similarity). Interacts with KCNE2.,tissue specificity:Detected in brain, in particular in amygdala and hippocampus, while expr

Background

The membrane protein encoded by this gene is a hyperpolarization-activated cation channel that contributes to the native pacemaker currents in heart and neurons. The encoded protein can homodimerize or heterodimerize with other pore-forming subunits to form a potassium channel. This channel may act as a receptor for sour tastes. [provided by RefSeq, Oct 2011],

matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images