



ATP4A Monoclonal Antibody

Catalog No	BYmab-05964
Isotype	IgG
Reactivity	Human;Rat;Mouse
Applications	WB
Gene Name	ATP4A
Protein Name	Potassium-transporting ATPase alpha chain 1 (EC 3.6.3.10) (Gastric H(+)/K(+) ATPase subunit alpha) (Proton pump)
Immunogen	Synthesized peptide derived from human protein . at AA range: 800-880
Specificity	ATP4A Monoclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Storage Stability Synonyms	-20°C/1 year
	-20°C/1 year 113kD
Synonyms	
Synonyms Observed Band	113kD Apical cell membrane; Multi-pass membrane protein. Localized in the apical canalicular membrane of parietal cells (PubMed:24188822). Localized in the
Synonyms Observed Band Cell Pathway	113kD Apical cell membrane; Multi-pass membrane protein. Localized in the apical canalicular membrane of parietal cells (PubMed:24188822). Localized in the apical canalicular membrane of parietal cells (PubMed:24188822).
Synonyms Observed Band Cell Pathway Tissue Specificity	Apical cell membrane; Multi-pass membrane protein. Localized in the apical canalicular membrane of parietal cells (PubMed:24188822). Localized in the apical canalicular membrane of parietal cells (PubMed:24188822). Expressed in gastric parietal cells (at protein level). catalytic activity:ATP + H(2)O + H(+)(In) + K(+)(Out) = ADP + phosphate + H(+)(Out) + K(+)(In).,function:Catalyzes the hydrolysis of ATP coupled with the exchange of H(+) and K(+) ions across the plasma membrane. Responsible for acid production in the stomach.,similarity:Belongs to the cation transport ATPase (P-type) family.,similarity:Belongs to the cation transport ATPase (P-type) family. Type IIC subfamily.,subunit:Composed of two subunits: alpha (catalytic) and

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658



国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务,欢迎咨询



	heavily glycosylated beta subunit. This enzyme is a proton pump that catalyzes the hydrolysis of ATP coupled with the exchange of H(+) and K(+) ions across the plasma membrane. It is also responsible for gastric acid secretion. This gene encodes a catalytic alpha subunit of the gastric H+, K+-ATPase. [provided by RefSeq, Jul 2008],
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images

Nanjing BYabscience technology Co.,Ltd