



PPR1A Monoclonal Antibody

Catalog No	BYmab-05981
Isotype	lgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	PPP1R1A IPP1
Protein Name	Protein phosphatase 1 regulatory subunit 1A (Protein phosphatase inhibitor 1) (I-1) (IPP-1)
Immunogen	Synthesized peptide derived from human protein . at AA range: 40-120
Specificity	PPR1A Monoclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	18kD
Cell Pathway	extracellular space,cytoplasm,
Tissue Specificity	Brain,Heart,
Function	function:Inhibitor of protein-phosphatase 1. This protein may be important in hormonal control of glycogen metabolism. Hormones that elevate intracellular cAMP increase I-1 activity in many tissues. I-1 activation may impose cAMP control over proteins that are not directly phosphorylated by PKA. Following a rise in intracellular calcium, I-1 is inactivated by calcineurin (or PP2B). Does not inhibit type-2 phosphatases.,PTM:Phosphorylation of Thr-35 is required for activity.,similarity:Belongs to the protein phosphatase inhibitor 1 family.,subunit:Interacts with PPP1R15A.,
Background	function:Inhibitor of protein-phosphatase 1. This protein may be important in hormonal control of glycogen metabolism. Hormones that elevate intracellular cAMP increase I-1 activity in many tissues. I-1 activation may impose cAMP
	Nanjing BYabscience technology Co.,Ltd



国内优质抗体供应商

国内加质加冲供应商 — 2 3 4 精准的 WB 检测服务 2 2 ----

24H 在线服务, 欢迎咨询



Products Images		
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.	
matters needing attention	Avoid repeated freezing and thawing!	
	control over proteins that are not directly phosphorylated by PKA. Following a rise in intracellular calcium, I-1 is inactivated by calcineurin (or PP2B). Does not inhibit type-2 phosphatases.,PTM:Phosphorylation of Thr-35 is required for activity.,similarity:Belongs to the protein phosphatase inhibitor 1 family.,subunit:Interacts with PPP1R15A.,	

Nanjing BYabscience technology Co.,Ltd