



# RIN2 Monoclonal Antibody

<b>Catalog No</b>	BYmab-06022
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human;Mouse
<b>Applications</b>	WB
<b>Gene Name</b>	RIN2 RASSF4
<b>Protein Name</b>	Ras and Rab interactor 2 (Ras association domain family 4) (Ras inhibitor JC265) (Ras interaction/interference protein 2)
<b>Immunogen</b>	Synthesized peptide derived from human protein . at AA range: 70-150
<b>Specificity</b>	RIN2 Monoclonal Antibody detects endogenous levels of protein.
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
<b>Source</b>	Monoclonal, Mouse,IgG
<b>Purification</b>	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Dilution</b>	WB 1:500-2000
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	
<b>Observed Band</b>	98kD
<b>Cell Pathway</b>	Cytoplasm .
<b>Tissue Specificity</b>	Widely expressed. Expressed in heart, kidney, lung placenta. Expressed at low level in skeletal muscle, spleen and peripheral blood.
<b>Function</b>	function:Ras effector protein. May function as an upstream activator and/or downstream effector for RAB5B in endocytic pathway. May function as a guanine nucleotide exchange (GEF) of RAB5B, required for activating the RAB5 proteins by exchanging bound GDP for free GTP.,similarity:Belongs to the RIN (Ras interaction/interference) family.,similarity:Contains 1 Ras-associating domain.,similarity:Contains 1 SH2 domain.,similarity:Contains 1 VPS9 domain.,subunit:Homotetramer; probably composed of anti-parallel linkage of two parallel dimers. Interacts with Ras. Interacts with RAB5B, with a much higher affinity for GTP-bound activated RAB5B. Does not interact with other members of the Rab family.,tissue specificity:Widely expressed. Expressed in heart, kidney, lung placenta. Expressed at low level in skeletal muscle, spleen and peripheral

**Nanjing BYabscience technology Co.,Ltd**



blood.,

**Background**

The RAB5 protein is a small GTPase involved in membrane trafficking in the early endocytic pathway. The protein encoded by this gene binds the GTP-bound form of the RAB5 protein preferentially over the GDP-bound form, and functions as a guanine nucleotide exchange factor for RAB5. The encoded protein is found primarily as a tetramer in the cytoplasm and does not bind other members of the RAB family. Mutations in this gene cause macrocephaly alopecia cutis laxa and scoliosis (MACS) syndrome, an elastic tissue disorder, as well as the related connective tissue disorder, RIN2 syndrome. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Jun 2011],

**matters needing attention**

Avoid repeated freezing and thawing!

**Usage suggestions**

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## Products Images

Nanjing BYabscience technology Co.,Ltd

网址: [www.njbybio.com](http://www.njbybio.com)

官方热线: 025-5229-8998

监督电话: 15950492658