



RAIN Monoclonal Antibody

| Catalog No | BYmab-06032 |
|---|---|
| Isotype | IgG |
| Reactivity | Human;Rat;Mouse; |
| Applications | WB |
| Gene Name | RASIP1 |
| Protein Name | Ras-interacting protein 1 (Rain) |
| Immunogen | Synthesized peptide derived from human protein . at AA range: 620-700 |
| Specificity | RAIN Monoclonal Antibody detects endogenous levels of protein. |
| Formulation | Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. |
| Source | Monoclonal, Mouse,IgG |
| Purification | The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. |
| Dilution | WB 1:500-2000 |
| Concentration | 1 mg/ml |
| Concentiation | i ing/iii |
| Purity | ≥90% |
| | • |
| Purity | ≥90% |
| Purity Storage Stability | ≥90% |
| Purity Storage Stability Synonyms | ≥90% -20°C/1 year |
| Purity Storage Stability Synonyms Observed Band | ≥90% -20°C/1 year 105kD Cytoplasm, perinuclear region . Golgi apparatus, Golgi stack . Associated with |
| Purity Storage Stability Synonyms Observed Band Cell Pathway | ≥90% -20°C/1 year 105kD Cytoplasm, perinuclear region . Golgi apparatus, Golgi stack . Associated with perinuclear vesicles. Is recruited to Golgi stacks by activated HRAS. Highly expressed in heart. Detected at lower levels in placenta and pancreas. function:May act as effector for Golgi-bound HRAS and other Ras-like proteins. May promote HRAS-mediated transformation.,similarity:Contains 1 dilute domain.,similarity:Contains 1 Ras-associating domain.,subcellular location:Associated with perinuclear vesicles. Is recruited to Golgi stacks by activated HRAS.,subunit:Interacts with Ras family members that have been |
| Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity | ≥90% -20°C/1 year 105kD Cytoplasm, perinuclear region . Golgi apparatus, Golgi stack . Associated with perinuclear vesicles. Is recruited to Golgi stacks by activated HRAS. Highly expressed in heart. Detected at lower levels in placenta and pancreas. function:May act as effector for Golgi-bound HRAS and other Ras-like proteins. May promote HRAS-mediated transformation.,similarity:Contains 1 dilute domain.,similarity:Contains 1 Ras-associating domain.,subcellular location:Associated with perinuclear vesicles. Is recruited to Golgi stacks by activated HRAS.,subunit:Interacts with Ras family members that have been activated by GTP binding. Interacts with HRAS, RAP1A, RAP2, RRAS, RAF1 and RRAS2.,tissue specificity:Highly expressed in heart. Detected at lower levels in |

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658





| location:Associated with perinuclear vesicles. Is recruited to Golgi stacks by activated HRAS.,subunit:Interacts with Ras family members that have been activated by GTP binding. Interacts with HRAS, RAP1A, RAP2, RRAS, RAF1 and RRAS2.,tissue specificity:Highly expressed in heart. Detected at lower levels in placenta and pancreas., |
|---|
| Avaid non-actable for-aming and the avairant |

matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using RAIN Monoclonal Antibody

Nanjing BYabscience technology Co.,Ltd