



PTPRT Monoclonal Antibody

Catalog No	BYmab-06069
Isotype	IgG
Reactivity	Human;Mouse
Applications	WB
Gene Name	PTPRT KIAA0283
Protein Name	Receptor-type tyrosine-protein phosphatase T (R-PTP-T) (EC 3.1.3.48) (Receptor-type tyrosine-protein phosphatase rho) (RPTP-rho)
Immunogen	Synthesized peptide derived from human protein . at AA range: 200-280
Specificity	PTPRT Monoclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	158kD
Cell Pathway	Membrane; Single-pass type I membrane protein.
Tissue Specificity	Expressed in colon, lung, heart and testis, as well as in fetal and adult brain. Not detected in muscle and peripheral blood leukocytes.
Function	catalytic activity:Protein tyrosine phosphate + H(2)O = protein tyrosine + phosphate.,function:May be involved in both signal transduction and cellular adhesion in the CNS.,similarity:Belongs to the protein-tyrosine phosphatase family. Receptor class 2B subfamily.,similarity:Contains 1 Ig-like C2-type (immunoglobulin-like) domain.,similarity:Contains 1 MAM domain.,similarity:Contains 2 tyrosine-protein phosphatase domains.,similarity:Contains 4 fibronectin type-III domains.,tissue specificity:Expressed in colon, lung, heart and testis, as well as in fetal and adult brain. Not detected in muscle and peripheral blood leukocytes.,
Background	The protein encoded by this gene is a member of the protein tyrosine phosphatase (PTP) family. PTPs are known to be signaling molecules that

Nanjing BYabscience technology Co.,Ltd



regulate a variety of cellular processes including cell growth, differentiation, mitotic cycle, and oncogenic transformation. This PTP possesses an extracellular region, a single transmembrane region, and two tandem intracellular catalytic domains, and thus represents a receptor-type PTP. The extracellular region contains a meprin-A5 antigen-PTP (MAM) domain, Ig-like and fibronectin type III-like repeats. The protein domain structure and the expression pattern of the mouse counterpart of this PTP suggest its roles in both signal transduction and cellular adhesion in the central nervous system. Two alternatively spliced transcript variants of this gene, which encode distinct proteins, have been reported. [provided by RefSeq, Jul 2008],

matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images