



## **RGS22** Monoclonal Antibody

Isotype   IgG     Reactivity   Human;Rat;Mouse;     Applications   WB     Gene Name   RGS22     Protein Name   Regulator of G-protein signaling 22 (RGS22)     Immunogen   Synthesized peptide derived from human protein . at AA range: 320-400     Specificity   RGS22 Monoclonal Antibody detects endogenous levels of protein.     Formulation   Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.     Source   Monoclonal, Mouse, IgG     Purification   The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.     Dilution   WB 1:500-2000     Concentration   1 mg/ml     Purity   ≥90%     Storage Stability   -20°C/1 year     Synonyms   Cytoplasm . Nucleus . Expressed in the cytoplasm of spermatogonia and spermatogonia to spermatids, also expressed in the nucleus.     Tissue Specificity   Testis-specific. Expressed in Leydig cells and spermatogenic cells from the spermatogenia to spermatid stages (at protein level).     Function   function. Inhibits signal transduction by increasing the GTPase activity of G protein apple suburity thereby driving them into their inactive GDP-bound form. similarity:Contains 2 RGS domains.,		
Reactivity Human;Rat;Mouse;   Applications WB   Gene Name RGS22   Protein Name Regulator of G-protein signaling 22 (RGS22)   Immunogen Synthesized peptide derived from human protein , at AA range; 320-400   Specificity RGS22 Monoclonal Antibody detects endogenous levels of protein.   Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.   Source Monoclonal, Mouse, IgG   Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.   Dilution WB 1:500-2000   Concentration 1 mg/ml   Purity ≥90%   Storage Stability -20°C/1 year   Synonyms -20°C/1 year   Observed Band 139kD   Cell Pathway Cytoplasm . Nucleus . Expressed in the cytoplasm of spermatogonia and spermatogenia cells from the spermatogonia to spermatids, also expressed in the nucleus.   Tissue Specificity Testis-specific. Expressed in Leydig cells and spermatogenic cells from the spermatogonia to spermatids also expressed and the nucleus.   Function function-Inhibits signal transduction by increasing the GTPase activity of G protein alpha subunits thereby driving them into their inactive GDP-bound formsimilarity:Contains 2 RGS domains	Catalog No	BYmab-06075
Applications WB   Gene Name RGS22   Protein Name Regulator of G-protein signaling 22 (RGS22)   Immunogen Synthesized peptide derived from human protein . at AA range: 320-400   Specificity RGS22 Monoclonal Antibody detects endogenous levels of protein.   Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.   Source Monoclonal, Mouse, IgG   Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.   Dilution WB 1:500-2000   Concentration 1 mg/ml   Purity ≥90%   Storage Stability -20°C/1 year   Synonyms Cytoplasm . Nucleus . Expressed in the cytoplasm of spermatogonia and spermatogenia to spermatide, also expressed in the nucleus.   Tissue Specificity Testis-specific. Expressed in Leydig cells and spermatogenic cells from the spermatogenia to spermatide, also expressed in the nucleus.   Function function.Inhibits signal transduction by increasing the GTPase activity of G protein alpha subunits thereby driving them into their inactive GDP-bound form., similarity:Contains 2 RGS domains., matters needing	lsotype	IgG
Gene Name   RGS22     Protein Name   Regulator of G-protein signaling 22 (RGS22)     Immunogen   Synthesized peptide derived from human protein . at AA range: 320-400     Specificity   RGS22 Monoclonal Antibody detects endogenous levels of protein.     Formulation   Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.     Source   Monoclonal, Mouse, IgG     Purtification   The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.     Dilution   WB 1:500-2000     Concentration   1 mg/ml     Purity   ≥90%     Storage Stability   -20°C/1 year     Synonyms   Cytoplasm . Nucleus . Expressed in the cytoplasm of spermatogonia and spermatocytes. In spermatids, also expressed in the nucleus.     Tissue Specificity   Testis-specific. Expressed in Leydig cells and spermatogonia and spermatocytes. In spermatids, also expressing the GTPase activity of G protein alpha subunits thereby driving them into their inactive GDP-bound form., similarity:Contains 2 RGS domains.,     Background   function.Inhibits signal transduction by increasing the GTPase activity of G protein alpha subunits thereby driving them into their inactive GDP-bound form., similarity:Contains 2 RGS domains.,	Reactivity	Human;Rat;Mouse;
Protein Name Regulator of G-protein signaling 22 (RGS22)   Immunogen Synthesized peptide derived from human protein . at AA range: 320-400   Specificity RGS22 Monoclonal Antibody detects endogenous levels of protein.   Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.   Source Monoclonal, Mouse, IgG   Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.   Dilution WB 1:500-2000   Concentration 1 mg/ml   Purity ≥90%   Storage Stability -20°C/1 year   Synonyms Cytoplasm. Nucleus . Expressed in the cytoplasm of spermatogonia and spermatocytes. In spermatids, also expressed in the nucleus.   Tissue Specificity Testis-specific. Expressed in Leydig cells and spermatogenic cells from the spermatogonia to spermatid stages (at protein level).   Function function:Inhibits signal transduction by increasing the GTPase activity of G protein alpha subunits thereby driving them into their inactive GDP-bound form., similarity:Contains 2 RGS domains.,   Background function:Inhibits signal transduction by increasing the GTPase activity of G protein alpha subunits thereby driving them into their inactive GDP-bound form., similarity:Contains 2 RGS domains.,	Applications	WB
Immunogen   Synthesized peptide derived from human protein . at AA range: 320-400     Specificity   RGS22 Monoclonal Antibody detects endogenous levels of protein.     Formulation   Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.     Source   Monoclonal, Mouse,IgG     Purification   The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.     Dilution   WB 1:500-2000     Concentration   1 mg/ml     Purity   ≥90%     Storage Stability   -20°C/1 year     Synonyms   Cytoplasm . Nucleus . Expressed in the cytoplasm of spermatogonia and spermatogonia to spermatids, also expressed in the nucleus.     Tissue Specificity   Testis-specific. Expressed in Leydig cells and spermatogonic cells from the spermatogonia to spermatid stages (at protein level).     Function   function: Inhibits signal transduction by increasing the GTPase activity of G protein alpha subunits thereby driving them into their inactive GDP-bound form., similarity: Contains 2 RGS domains.,     Background   function:Inhibits signal transduction by increasing the GTPase activity of G protein alpha subunits thereby driving them into their inactive GDP-bound form., similarity:Contains 2 RGS domains.,	Gene Name	RGS22
Specificity RGS22 Monoclonal Antibody detects endogenous levels of protein.   Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.   Source Monoclonal, Mouse,IgG   Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.   Dilution WB 1:500-2000   Concentration 1 mg/ml   Purity ≥90%   Storage Stability -20°C/1 year   Synonyms Observed Band   Observed Band 139kD   Cell Pathway Cytoplasm . Nucleus . Expressed in the cytoplasm of spermatogonia and spermatogenia to spermatids, also expressed in the nucleus.   Tissue Specificity Testis-specific. Expressed in Leydig cells and spermatogenic cells from the spermatogonia to spermatid stages (at protein level).   Function function:Inhibits signal transduction by increasing the GTPase activity of G protein alpha subunits thereby driving them into their inactive GDP-bound form, similarity.Contains 2 RGS domains.   Background function:Inhibits signal transduction by increasing the GTPase activity of G protein alpha subunits thereby driving them into their inactive GDP-bound form, similarity.contains 2 RGS domains.	Protein Name	Regulator of G-protein signaling 22 (RGS22)
Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.   Source Monoclonal, Mouse, IgG   Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.   Dilution WB 1:500-2000   Concentration 1 mg/ml   Purity ≥90%   Storage Stability -20°C/1 year   Synonyms -20°C/1 year   Observed Band 139kD   Cell Pathway Cytoplasm . Nucleus . Expressed in the cytoplasm of spermatogonia and spermatocytes. In spermatids, also expressed in the nucleus.   Tissue Specificity Testis-specific. Expressed in Leydig cells and spermatogenic cells from the spermatogonia to spermatid stages (at protein level).   Function function:Inhibits signal transduction by increasing the GTPase activity of G protein alpha subunits thereby driving them into their inactive GDP-bound form., similarity:Contains 2 RGS domains.,   Background function:Inhibits signal transduction by increasing the GTPase activity of G protein alpha subunits thereby driving them into their inactive GDP-bound form., similarity:Contains 2 RGS domains.,   matters needing Avoid repeated freezing and thawing!	Immunogen	Synthesized peptide derived from human protein . at AA range: 320-400
Source Monoclonal, Mouse,IgG   Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.   Dilution WB 1:500-2000   Concentration 1 mg/ml   Purity ≥90%   Storage Stability -20°C/1 year   Synonyms Cytoplasm . Nucleus . Expressed in the cytoplasm of spermatogonia and spermatocytes. In spermatids, also expressed in the nucleus.   Tissue Specificity Testis-specific. Expressed in Leydig cells and spermatogenic cells from the spermatogonia to spermatid stages (at protein level).   Function function:Inhibits signal transduction by increasing the GTPase activity of G protein alpha subunits thereby driving them into their inactive GDP-bound form.,similarity:Contains 2 RGS domains.,   Background function:Inhibits signal transduction by increasing the GTPase activity of G protein alpha subunits thereby driving them into their inactive GDP-bound form.,similarity:Contains 2 RGS domains.,	Specificity	RGS22 Monoclonal Antibody detects endogenous levels of protein.
Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.   Dilution WB 1:500-2000   Concentration 1 mg/ml   Purity ≥90%   Storage Stability -20°C/1 year   Synonyms -20°C/1 year   Observed Band 139kD   Cell Pathway Cytoplasm . Nucleus . Expressed in the cytoplasm of spermatogonia and spermatocytes. In spermatids, also expressed in the nucleus.   Tissue Specificity Testis-specific. Expressed in Leydig cells and spermatogenic cells from the spermatogonia to spermatid stages (at protein level).   Function function:Inhibits signal transduction by increasing the GTPase activity of G protein alpha subunits thereby driving them into their inactive GDP-bound form., similarity:Contains 2 RGS domains.,   Background function:Inhibits signal transduction by increasing the GTPase activity of G protein alpha subunits thereby driving them into their inactive GDP-bound form., similarity:Contains 2 RGS domains.,   matters needing Avoid repeated freezing and thawing!	Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
affinity-chromatography using epitope-specific immunogen.   Dilution WB 1:500-2000   Concentration 1 mg/ml   Purity ≥90%   Storage Stability -20°C/1 year   Synonyms -20°C/1 year   Observed Band 139kD   Cell Pathway Cytoplasm . Nucleus . Expressed in the cytoplasm of spermatogonia and spermatocytes. In spermatids, also expressed in the nucleus.   Tissue Specificity Testis-specific. Expressed in Leydig cells and spermatogenic cells from the spermatogonia to spermatid stages (at protein level).   Function function:Inhibits signal transduction by increasing the GTPase activity of G protein alpha subunits thereby driving them into their inactive GDP-bound form., similarity:Contains 2 RGS domains.,   Background function:Inhibits signal transduction by increasing the GTPase activity of G protein alpha subunits thereby driving them into their inactive GDP-bound form., similarity:Contains 2 RGS domains.,   matters needing Avoid repeated freezing and thawing!	Source	Monoclonal, Mouse,IgG
Concentration 1 mg/ml   Purity ≥90%   Storage Stability -20°C/1 year   Synonyms -20°C/1 year   Observed Band 139kD   Cell Pathway Cytoplasm . Nucleus . Expressed in the cytoplasm of spermatogonia and spermatocytes. In spermatids, also expressed in the nucleus.   Tissue Specificity Testis-specific. Expressed in Leydig cells and spermatogenic cells from the spermatogonia to spermatid stages (at protein level).   Function function:Inhibits signal transduction by increasing the GTPase activity of G protein alpha subunits thereby driving them into their inactive GDP-bound formsimilarity:Contains 2 RGS domains.,   Background function:Inhibits signal transduction by increasing the GTPase activity of G protein alpha subunits thereby driving them into their inactive GDP-bound formsimilarity:Contains 2 RGS domains.,   matters needing Avoid repeated freezing and thawing!	Purification	
Purity ≥90%   Storage Stability -20°C/1 year   Synonyms -20°C/1 year   Observed Band 139kD   Cell Pathway Cytoplasm . Nucleus . Expressed in the cytoplasm of spermatogonia and spermatocytes. In spermatids, also expressed in the nucleus.   Tissue Specificity Testis-specific. Expressed in Leydig cells and spermatogenic cells from the spermatogonia to spermatid stages (at protein level).   Function function:Inhibits signal transduction by increasing the GTPase activity of G protein alpha subunits thereby driving them into their inactive GDP-bound form., similarity:Contains 2 RGS domains.,   Background function:Inhibits signal transduction by increasing the GTPase activity of G protein alpha subunits thereby driving them into their inactive GDP-bound form., similarity:Contains 2 RGS domains.,   matters needing Avoid repeated freezing and thawing!	Dilution	WB 1:500-2000
Storage Stability -20°C/1 year   Synonyms -20°C/1 year   Observed Band 139kD   Cell Pathway Cytoplasm . Nucleus . Expressed in the cytoplasm of spermatogonia and spermatocytes. In spermatids, also expressed in the nucleus.   Tissue Specificity Testis-specific. Expressed in Leydig cells and spermatogenic cells from the spermatogonia to spermatid stages (at protein level).   Function function:Inhibits signal transduction by increasing the GTPase activity of G protein alpha subunits thereby driving them into their inactive GDP-bound form.,similarity:Contains 2 RGS domains.,   Background function:Inhibits signal transduction by increasing the GTPase activity of G protein alpha subunits thereby driving them into their inactive GDP-bound form.,similarity:Contains 2 RGS domains.,   matters needing Avoid repeated freezing and thawing!	Concentration	1 mg/ml
Synonyms   Observed Band 139kD   Cell Pathway Cytoplasm . Nucleus . Expressed in the cytoplasm of spermatogonia and spermatocytes. In spermatids, also expressed in the nucleus.   Tissue Specificity Testis-specific. Expressed in Leydig cells and spermatogenic cells from the spermatogonia to spermatid stages (at protein level).   Function function:Inhibits signal transduction by increasing the GTPase activity of G protein alpha subunits thereby driving them into their inactive GDP-bound form.,similarity:Contains 2 RGS domains.,   Background function:Inhibits signal transduction by increasing the GTPase activity of G protein alpha subunits thereby driving them into their inactive GDP-bound form.,similarity:Contains 2 RGS domains.,   matters needing Avoid repeated freezing and thawing!	Purity	≥90%
Observed Band139kDCell PathwayCytoplasm . Nucleus . Expressed in the cytoplasm of spermatogonia and spermatocytes. In spermatids, also expressed in the nucleus.Tissue SpecificityTestis-specific. Expressed in Leydig cells and spermatogenic cells from the spermatogonia to spermatid stages (at protein level).Functionfunction:Inhibits signal transduction by increasing the GTPase activity of G protein alpha subunits thereby driving them into their inactive GDP-bound form., similarity:Contains 2 RGS domains.,Backgroundfunction:Inhibits signal transduction by increasing the GTPase activity of G protein alpha subunits thereby driving them into their inactive GDP-bound form., similarity:Contains 2 RGS domains.,matters needingAvoid repeated freezing and thawing!	Storage Stability	-20°C/1 year
Cell PathwayCytoplasm . Nucleus . Expressed in the cytoplasm of spermatogonia and spermatocytes. In spermatids, also expressed in the nucleus.Tissue SpecificityTestis-specific. Expressed in Leydig cells and spermatogenic cells from the spermatogonia to spermatid stages (at protein level).Functionfunction:Inhibits signal transduction by increasing the GTPase activity of G protein alpha subunits thereby driving them into their inactive GDP-bound form.,similarity:Contains 2 RGS domains.,Backgroundfunction:Inhibits signal transduction by increasing the GTPase activity of G protein alpha subunits thereby driving them into their inactive GDP-bound form.,similarity:Contains 2 RGS domains.,matters needingAvoid repeated freezing and thawing!	Synonyms	
spermatocytes. In spermatids, also expressed in the nucleus.Tissue SpecificityTestis-specific. Expressed in Leydig cells and spermatogenic cells from the spermatogonia to spermatid stages (at protein level).Functionfunction:Inhibits signal transduction by increasing the GTPase activity of G protein alpha subunits thereby driving them into their inactive GDP-bound form.,similarity:Contains 2 RGS domains.,Backgroundfunction:Inhibits signal transduction by increasing the GTPase activity of G protein alpha subunits thereby driving them into their inactive GDP-bound form.,similarity:Contains 2 RGS domains.,matters needingAvoid repeated freezing and thawing!	Observed Band	139kD
Spermatogonia to spermatid stages (at protein level).Functionfunction:Inhibits signal transduction by increasing the GTPase activity of G protein alpha subunits thereby driving them into their inactive GDP-bound form.,similarity:Contains 2 RGS domains.,Backgroundfunction:Inhibits signal transduction by increasing the GTPase activity of G protein alpha subunits thereby driving them into their inactive GDP-bound form.,similarity:Contains 2 RGS domains.,matters needingAvoid repeated freezing and thawing!	Cell Pathway	Cytoplasm . Nucleus . Expressed in the cytoplasm of spermatogonia and spermatocytes. In spermatids, also expressed in the nucleus.
alpha subunits thereby driving them into their inactive GDP-bound form.,similarity:Contains 2 RGS domains.,Backgroundfunction:Inhibits signal transduction by increasing the GTPase activity of G protein alpha subunits thereby driving them into their inactive GDP-bound form.,similarity:Contains 2 RGS domains.,matters needingAvoid repeated freezing and thawing!	Tissue Specificity	Testis-specific. Expressed in Leydig cells and spermatogenic cells from the spermatogonia to spermatid stages (at protein level).
alpha subunits thereby driving them into their inactive GDP-bound form.,similarity:Contains 2 RGS domains.,   matters needing Avoid repeated freezing and thawing!	Function	function:Inhibits signal transduction by increasing the GTPase activity of G protein alpha subunits thereby driving them into their inactive GDP-bound form.,similarity:Contains 2 RGS domains.,
	Background	function:Inhibits signal transduction by increasing the GTPase activity of G protein alpha subunits thereby driving them into their inactive GDP-bound form.,similarity:Contains 2 RGS domains.,
	matters needing attention	Avoid repeated freezing and thawing!

## Nanjing BYabscience technology Co.,Ltd



国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务,欢迎咨询



Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## **Products Images**

Nanjing BYabscience technology Co.,Ltd