



STAC Monoclonal Antibody

Catalog No	BYmab-06168
Isotype	IgG
Reactivity	Human;Mouse
Applications	WB
Gene Name	STAC
Protein Name	SH3 and cysteine-rich domain-containing protein (Src homology 3 and cysteine-rich domain-containing protein)
Immunogen	Synthesized peptide derived from part region of human protein
Specificity	STAC Monoclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	44kD
Cell Pathway	Cytoplasm, cytosol . Cell membrane ; Peripheral membrane protein ; Cytoplasmic side . Cell membrane, sarcolemma ; Peripheral membrane protein ; Cytoplasmic side .
Tissue Specificity	Brain,Kidney,
Function	function:Probably involved in a neuron-specific signal transduction.,similarity:Contains 1 phorbol-ester/DAG-type zinc finger.,similarity:Contains 1 SH3 domain.,
Background	function:Probably involved in a neuron-specific signal transduction.,similarity:Contains 1 phorbol-ester/DAG-type zinc finger.,similarity:Contains 1 SH3 domain.,
matters needing attention	Avoid repeated freezing and thawing!


Nanjing BYabscience technology Co.,Ltd



Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using STAC Monoclonal Antibody

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com

官方热线: 025-5229-8998

监督电话: 15950492658