




## **UBS3B Monoclonal Antibody**

Catalog No         BYmab-06249           Isotype         IgG           Reactivity         Human;Mouse           Applications         WB           Gene Name         UBASH3B KIAA1959 STS1           Protein Name         Ubiquitin-associated and SH3 domain-containing protein B (EC 3.1.3.48) (Cbi-interacting protein p70) (Suppressor of T-cell receptor signaling 1) (STS-1) (T-cell ubiquitin ligand 2) (TrULA-2) (Tyrosine-           Immunogen         Synthesized peptide derived from part region of human protein           Specificity         UBS3B Monoclonal Antibody detects endogenous levels of protein.           Formulation         Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.           Source         Monoclonal, Mouse, IgG           Purification         The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.           Dilution         WB 1:500-2000           Concentration         1 mg/ml           Purity         ≥90%           Storage Stability         -20°C/1 year           Synonyms         Observed Band         71kD           Cell Pathway         Cytoplasm . Nucleus .           Tissue Specificity         Brain, Kidney, Liver, Placenta, Platelet, function: Interferes with CBL. mediated down-regulation and degradation of receptor-type tyrosine kinases. Promotes accumulation of activated target receptors, such as T-c		
Applications WB Gene Name UBASH3B KIAA1959 STS1  Protein Name Ubiquitin-associated and SH3 domain-containing protein B (EC 3.1.3.48) (Cbl-interacting protein p70) (Suppressor of T-cell receptor signaling 1) (STS-1) (T-cell ubiquitin ligand 2) (TULA-2) (Tyrosine- Immunogen Synthesized peptide derived from part region of human protein Specificity UBS3B Monoclonal Antibody detects endogenous levels of protein.  Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.  Source Monoclonal, Mouse,IgG  Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-2000  Concentration 1 mg/ml  Purity ≥90% Storage Stability -20°C/1 year  Synonyms  Observed Band 71kD  Cell Pathway Cytoplasm . Nucleus .  Tissue Specificity Brain,Kidney,Liver,Placenta,Platelet, Function function:Interferes with CBL-mediated down-regulation and degradation of receptor-type tryosine kinases. Promotes accumulation of activated target receptors, such as T-cell receptors and EGFR, on the cell surface, similarity Contains 1 UBA domain, subunit/Interacts with CBL-part of a complex containing CBL and activated EGFR. Interacts with ubiquitin and with mono-ubiquitinated proteins., This gene encodes a protein that contains a ubiquitin associated domain at the N-terminus, an SH3 domain, and a C-terminal domain with similarities to the catalytic motif of phosphoglycerate mutase. The encoded proteins was found to	Catalog No	BYmab-06249
Applications  Gene Name  UBASH3B KIAA1959 STS1  Protein Name  Ubiquitin-associated and SH3 domain-containing protein B (EC 3.1.3.48) (CbI-interacting protein p70) (Suppressor of T-cell receptor signaling 1) (STS-1) (T-cell ubiquitin ligand 2) (TULA-2) (Tyrosine-Immunogen  Synthesized peptide derived from part region of human protein  Specificity  UBS3B Monoclonal Antibody detects endogenous levels of protein.  Formulation  Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.  Source  Monoclonal, Mouse, IgG  Purification  The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution  WB 1:500-2000  Concentration  1 mg/ml  Purity  ≥90%  Storage Stability  -20°C/1 year  Synonyms  Observed Band  71kD  Cell Pathway  Cytoplasm , Nucleus .  Tissue Specificity  Brain, Kidney, Liver, Placenta, Platelet,  Function  function: Interferes with CBL—mediated down-regulation and degradation of receptor-type tyrosine kinases. Promotes accumulation of activated target receptors, such as T-cell receptors and EGFR, on the cell surface, similarity Contains 1 SH3 domain, similarity. Contains 1 UBA domain, subunit. Interacts with CBL. Part of a complex containing CBL and activated EGFR. Interacts with ubiquitin and with mono-ubiquitinated proteins.  This gene encodes a protein that contains a ubiquitin associated domain at the N-terminus, an SH3 domain, and a C-terminal domain with similarities to the catalytic motif of phosphoglycerate mutase. The encoded proteins.	Isotype	IgG
Gene Name         UBASH3B KIAA1959 STS1           Protein Name         Ubiquitin-associated and SH3 domain-containing protein B (EC 3.1.3.48) (Cbl-interacting protein p70) (Suppressor of T-cell receptor signaling 1) (STS-1) (T-cell ubiquitin ligand 2) (TULA-2) (Tyrosine-limmunogen           Immunogen         Synthesized peptide derived from part region of human protein           Specificity         UBS3B Monoclonal Antibody detects endogenous levels of protein.           Formulation         Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.           Source         Monoclonal, Mouse, lgG           Purification         The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.           Dilution         WB 1:500-2000           Concentration         1 mg/ml           Purity         ≥90%           Storage Stability         -20°C/1 year           Synonyms         Observed Band         71kD           Cell Pathway         Cytoplasm . Nucleus .           Tissue Specificity         Brain, Kidney, Liver, Placenta, Platelet,           Function         function: Interferes with CBL-mediated down-regulation and degradation of receptor-type tyrosine kinases. Promotes accumulation of activated target receptors, such as T-cell receptors and EGFR, on the cell surface, similarity. Contains 1 UBA domain, similarity. Contains 1 UBA domain, subunit: Interacts with ubiquitin and with mono-ubiquitinated proteins.	Reactivity	Human;Mouse
Protein Name  Ubiquitin-associated and SH3 domain-containing protein B (EC 3.1.3.48) (Cbl-interacting protein p70) (Suppressor of T-cell receptor signaling 1) (STS-1) (T-cell ubiquitin ligand 2) (TULA-2) (Tyrosine-  Immunogen  Synthesized peptide derived from part region of human protein  Specificity  UBS3B Monoclonal Antibody detects endogenous levels of protein.  Formulation  Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.  Source  Monoclonal, Mouse, IgG  Purification  The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution  WB 1:500-2000  Concentration  1 mg/ml  Purity  ≥90%  Storage Stability  -20°C/1 year  Synonyms  Observed Band  71kD  Cell Pathway  Cytoplasm . Nucleus .  Tissue Specificity  Brain,Kidney,Liver,Placenta,Platelet, function:Interferes with CBL-mediated down-regulation and degradation of receptor-type tyrosine kinases. Promotes accumulation of activated target receptors, such as T-cell receptors and EGFR, on the cell surface, similarity:Contains 1 UBA domain, submit the tracts with CBL. Part of a complex containing CBL and activated EGFR. Interacts with CBL. Part of a complex containing CBL and activated EGFR. Interacts with Usiquitin and with mono-ubiquitinated proteins.,  This gene encodes a protein that contains a ubiquitin associated domain at the N-terminus, an SH3 domain, and a C-terminal domain with similarities to the callytic motif of phosphoglycerate mutase. The encoded protein was found to	Applications	WB
Immunogen         Synthesized peptide derived from part region of human protein           Specificity         UBS3B Monoclonal Antibody detects endogenous levels of protein.           Formulation         Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.           Source         Monoclonal, Mouse, IgG           Purification         The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.           Dilution         WB 1:500-2000           Concentration         1 mg/ml           Purity         ≥90%           Storage Stability         -20°C/1 year           Synonyms         Observed Band         71kD           Cell Pathway         Cytoplasm . Nucleus .           Tissue Specificity         Brain,Kidney,Liver,Placenta,Platelet,           Function         function:Interferes with CBL-mediated down-regulation and degradation of receptor-type tyrosine kinases. Promotes accumulation of activated target receptors, such as T-cell receptors and EGFR, on the cell surface, similarity:Contains 1 SH3 domain, similarity:Contains 1 UBA domain, similarity:Contains 1 SH3 domain, similarity:Containing CBL and activated EGFR. Interacts with CBL. Part of a complex containing CBL and activated EGFR. Interacts with bidjuitin and with mono-ubiquitinated proteins.,           Background         This gene encodes a protein that contains a ubiquitin associated domain at the N-terminus, an SH3 domain, and a C-terminal domain with similarities to the catalytic motif of phosphoglydycera	Gene Name	UBASH3B KIAA1959 STS1
Specificity         UBS3B Monoclonal Antibody detects endogenous levels of protein.           Formulation         Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.           Source         Monoclonal, Mouse,IgG           Purification         The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.           Dilution         WB 1:500-2000           Concentration         1 mg/ml           Purity         ≥90%           Storage Stability         -20°C/1 year           Synonyms         Observed Band           Observed Band         71kD           Cell Pathway         Cytoplasm . Nucleus .           Tissue Specificity         Brain,Kidney,Liver,Placenta,Platelet,           Function         function:Interferes with CBL-mediated down-regulation and degradation of receptor-type tyrosine kinases. Promotes accumulation of activated target receptors, such as T-cell receptors and EGFR, on the cell surface, similarity:Contains 1 UBA, domain, similarity:Contains 1 UBA, domain, similarity:Contains 1 UBA, domain, similarity:Contain activated EGFR. Interacts with Ubiquitin and with mono-ubiquitinated proteins.,           Background         This gene encodes a protein that contains a ubiquitin associated domain at the N-terminus, an SH3 domain, and a C-terminal domain with similarities to the catalytic motif of phosphoglycerate mutase. The encoded protein was found to	Protein Name	Ubiquitin-associated and SH3 domain-containing protein B (EC 3.1.3.48) (Cbl-interacting protein p70) (Suppressor of T-cell receptor signaling 1) (STS-1) (T-cell ubiquitin ligand 2) (TULA-2) (Tyrosine-
Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.  Source Monoclonal, Mouse,IgG  Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-2000 Concentration 1 mg/ml  Purity 290% Storage Stability -20°C/1 year  Synonyms Observed Band 71kD Cell Pathway Cytoplasm . Nucleus .  Tissue Specificity Brain,Kidney,Liver,Placenta,Platelet,  Function function:Interferes with CBL-mediated down-regulation and degradation of receptors, such as T-cell receptors and EGFR, on the cell surface, similarity:Contains 1 UBA domain, subunit:Interacts with CBL. Part of a complex containing CBL and activated EGFR. Interacts with ubiquitin and with mono-ubiquitinated proteins.,  Background This gene encodes a protein that contains a ubiquitin associated domain at the N-terminus, an SH3 domain, and a C-terminal domain with similarities to the catalytic motif of phosphoglycerate mutase. The encoded protein was found to	Immunogen	Synthesized peptide derived from part region of human protein
Source       Monoclonal, Mouse,IgG         Purification       The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.         Dilution       WB 1:500-2000         Concentration       1 mg/ml         Purity       ≥90%         Storage Stability       -20°C/1 year         Synonyms       Observed Band       71kD         Cell Pathway       Cytoplasm . Nucleus .         Tissue Specificity       Brain,Kidney,Liver,Placenta,Platelet,         Function       function:Interferes with CBL-mediated down-regulation and degradation of receptor-type tyrosine kinases. Promotes accumulation of activated target receptors, such as T-cell receptors and EGFR, on the cell surface.;similarity:Contains 1 SH3 domain.,similarity:Contains 1 UBA domain.,similarity:Contains 1 UBA domain.,similarity:Contains 1 UBA admain.,similarity:Contains 1 UBA activated EGFR. Interacts with ubiquitin and with mono-ubiquitinated proteins.,         Background       This gene encodes a protein that contains a ubiquitin associated domain at the N-terminus, an SH3 domain, and a C-terminal domain with similarities to the catalytic motif of phosphoglycerate mutase. The encoded protein was found to	Specificity	UBS3B Monoclonal Antibody detects endogenous levels of protein.
Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 71kD Cell Pathway Cytoplasm . Nucleus .  Tissue Specificity Brain,Kidney,Liver,Placenta,Platelet, function function:Interferes with CBL-mediated down-regulation and degradation of receptor-type tyrosine kinases. Promotes accumulation of activated target receptors, such as T-cell receptors and EGFR, on the cell surface.,similarity:Contains 1 SH3 domain.,similarity:Contains 1 UBA domain.,subunit:Interacts with CBL. Part of a complex containing CBL and activated EGFR. Interacts with ubiquitin and with mono-ubiquitinated proteins.,  Background This gene encodes a protein that contains a ubiquitin associated domain at the N-terminus, an SH3 domain, and a C-terminal domain with similarities to the catalytic motif of phosphoglycerate mutase. The encoded protein was found to	Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-2000  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms  Observed Band 71kD  Cell Pathway Cytoplasm . Nucleus .  Tissue Specificity Brain,Kidney,Liver,Placenta,Platelet,  Function function:Interferes with CBL-mediated down-regulation and degradation of receptor-type tyrosine kinases. Promotes accumulation of activated target receptors, such as T-cell receptors and EGFR, on the cell surface.,similarity:Contains 1 JBA domain.,subunit:Interacts with CBL. Part of a complex containing CBL and activated EGFR. Interacts with ubiquitin and with mono-ubiquitinated proteins.,  Background This gene encodes a protein that contains a ubiquitin associated domain at the N-terminus, an SH3 domain, and a C-terminal domain with similarities to the catalytic motif of phosphoglycerate mutase. The encoded protein was found to	Source	Monoclonal, Mouse,IgG
Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms  Observed Band 71kD  Cell Pathway Cytoplasm . Nucleus .  Tissue Specificity Brain,Kidney,Liver,Placenta,Platelet,  Function function:Interferes with CBL-mediated down-regulation and degradation of receptor-type tyrosine kinases. Promotes accumulation of activated target receptors, such as T-cell receptors and EGFR, on the cell surface.,similarity:Contains 1 SH3 domain.,similarity:Contains 1 UBA domain.,subunit:Interacts with CBL. Part of a complex containing CBL and activated EGFR. Interacts with ubiquitin and with mono-ubiquitinated proteins.,  Background This gene encodes a protein that contains a ubiquitin associated domain at the N-terminus, an SH3 domain, and a C-terminal domain with similarities to the catalytic motif of phosphoglycerate mutase. The encoded protein was found to	Purification	· · · · · · · · · · · · · · · · · · ·
Purity ≥90%  Storage Stability -20°C/1 year  Synonyms  Observed Band 71kD  Cell Pathway Cytoplasm . Nucleus .  Tissue Specificity Brain,Kidney,Liver,Placenta,Platelet,  Function function:Interferes with CBL-mediated down-regulation and degradation of receptor-type tyrosine kinases. Promotes accumulation of activated target receptors, such as T-cell receptors and EGFR, on the cell surface.,similarity:Contains 1 SH3 domain.,similarity:Contains 1 UBA domain.,subunit:Interacts with CBL. Part of a complex containing CBL and activated EGFR. Interacts with ubiquitin and with mono-ubiquitinated proteins.,  Background This gene encodes a protein that contains a ubiquitin associated domain at the N-terminus, an SH3 domain, and a C-terminal domain with similarities to the catalytic motif of phosphoglycerate mutase. The encoded protein was found to	Dilution	WB 1:500-2000
Storage Stability  -20°C/1 year  Synonyms  Observed Band  71kD  Cell Pathway  Cytoplasm . Nucleus .  Tissue Specificity  Brain,Kidney,Liver,Placenta,Platelet,  Function  function:Interferes with CBL-mediated down-regulation and degradation of receptor-type tyrosine kinases. Promotes accumulation of activated target receptors, such as T-cell receptors and EGFR, on the cell surface.,similarity:Contains 1 SH3 domain.,similarity:Contains 1 UBA domain.,subunit:Interacts with CBL. Part of a complex containing CBL and activated EGFR. Interacts with ubiquitin and with mono-ubiquitinated proteins.,  Background  This gene encodes a protein that contains a ubiquitin associated domain at the N-terminus, an SH3 domain, and a C-terminal domain with similarities to the catalytic motif of phosphoglycerate mutase. The encoded protein was found to	Concentration	1 mg/ml
Synonyms  Observed Band  71kD  Cell Pathway  Cytoplasm . Nucleus .  Tissue Specificity  Brain,Kidney,Liver,Placenta,Platelet,  Function  function:Interferes with CBL-mediated down-regulation and degradation of receptor-type tyrosine kinases. Promotes accumulation of activated target receptors, such as T-cell receptors and EGFR, on the cell surface.,similarity:Contains 1 SH3 domain.,similarity:Contains 1 UBA domain.,subunit:Interacts with CBL. Part of a complex containing CBL and activated EGFR. Interacts with ubiquitin and with mono-ubiquitinated proteins.,  Background  This gene encodes a protein that contains a ubiquitin associated domain at the N-terminus, an SH3 domain, and a C-terminal domain with similarities to the catalytic motif of phosphoglycerate mutase. The encoded protein was found to	Purity	≥90%
Observed Band  Cell Pathway  Cytoplasm . Nucleus .  Tissue Specificity  Brain,Kidney,Liver,Placenta,Platelet,  function  function:Interferes with CBL-mediated down-regulation and degradation of receptor-type tyrosine kinases. Promotes accumulation of activated target receptors, such as T-cell receptors and EGFR, on the cell surface.,similarity:Contains 1 SH3 domain.,similarity:Contains 1 UBA domain.,subunit:Interacts with CBL. Part of a complex containing CBL and activated EGFR. Interacts with ubiquitin and with mono-ubiquitinated proteins.,  This gene encodes a protein that contains a ubiquitin associated domain at the N-terminus, an SH3 domain, and a C-terminal domain with similarities to the catalytic motif of phosphoglycerate mutase. The encoded protein was found to	Storage Stability	-20°C/1 year
Cell Pathway         Cytoplasm . Nucleus .           Tissue Specificity         Brain,Kidney,Liver,Placenta,Platelet,           Function         function:Interferes with CBL-mediated down-regulation and degradation of receptor-type tyrosine kinases. Promotes accumulation of activated target receptors, such as T-cell receptors and EGFR, on the cell surface.,similarity:Contains 1 SH3 domain.,similarity:Contains 1 UBA domain.,subunit:Interacts with CBL. Part of a complex containing CBL and activated EGFR. Interacts with ubiquitin and with mono-ubiquitinated proteins.,           Background         This gene encodes a protein that contains a ubiquitin associated domain at the N-terminus, an SH3 domain, and a C-terminal domain with similarities to the catalytic motif of phosphoglycerate mutase. The encoded protein was found to	Synonyms	
Tissue Specificity  Brain,Kidney,Liver,Placenta,Platelet,  function  function:Interferes with CBL-mediated down-regulation and degradation of receptor-type tyrosine kinases. Promotes accumulation of activated target receptors, such as T-cell receptors and EGFR, on the cell surface.,similarity:Contains 1 SH3 domain.,similarity:Contains 1 UBA domain.,subunit:Interacts with CBL. Part of a complex containing CBL and activated EGFR. Interacts with ubiquitin and with mono-ubiquitinated proteins.,  This gene encodes a protein that contains a ubiquitin associated domain at the N-terminus, an SH3 domain, and a C-terminal domain with similarities to the catalytic motif of phosphoglycerate mutase. The encoded protein was found to	Observed Band	71kD
Function  function:Interferes with CBL-mediated down-regulation and degradation of receptor-type tyrosine kinases. Promotes accumulation of activated target receptors, such as T-cell receptors and EGFR, on the cell surface.,similarity:Contains 1 SH3 domain.,similarity:Contains 1 UBA domain.,subunit:Interacts with CBL. Part of a complex containing CBL and activated EGFR. Interacts with ubiquitin and with mono-ubiquitinated proteins.,  This gene encodes a protein that contains a ubiquitin associated domain at the N-terminus, an SH3 domain, and a C-terminal domain with similarities to the catalytic motif of phosphoglycerate mutase. The encoded protein was found to	Cell Pathway	Cytoplasm . Nucleus .
receptor-type tyrosine kinases. Promotes accumulation of activated target receptors, such as T-cell receptors and EGFR, on the cell surface., similarity:Contains 1 SH3 domain., similarity:Contains 1 UBA domain., subunit:Interacts with CBL. Part of a complex containing CBL and activated EGFR. Interacts with ubiquitin and with mono-ubiquitinated proteins.,  This gene encodes a protein that contains a ubiquitin associated domain at the N-terminus, an SH3 domain, and a C-terminal domain with similarities to the catalytic motif of phosphoglycerate mutase. The encoded protein was found to	Tissue Specificity	Brain,Kidney,Liver,Placenta,Platelet,
N-terminus, an SH3 domain, and a C-terminal domain with similarities to the catalytic motif of phosphoglycerate mutase. The encoded protein was found to	Function	receptor-type tyrosine kinases. Promotes accumulation of activated target receptors, such as T-cell receptors and EGFR, on the cell surface.,similarity:Contains 1 SH3 domain.,similarity:Contains 1 UBA domain.,subunit:Interacts with CBL. Part of a complex containing CBL and
, , , ,	Background	N-terminus, an SH3 domain, and a C-terminal domain with similarities to the catalytic motif of phosphoglycerate mutase. The encoded protein was found to

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658







	platelet-derived growth factor receptor. [provided by RefSeq, Jul 2008],
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## **Products Images**

Nanjing BYabscience technology Co.,Ltd