



SYT5 Monoclonal Antibody

Catalog No	BYmab-06260
lsotype	lgG
Reactivity	Human;Rat;Mouse
Applications	WB
Gene Name	SYT5
Protein Name	Synaptotagmin-5 (Synaptotagmin V) (SytV)
Immunogen	Synthesized peptide derived from part region of human protein
Specificity	SYT5 Monoclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	42kD
Cell Pathway	Cytoplasmic vesicle, secretory vesicle, synaptic vesicle membrane ; Single-pass membrane protein . Recycling endosome membrane ; Single-pass membrane protein . In mast cells, localizes to the endocytic recycling compartment
Tissue Specificity	Brain,
Function	cofactor:Binds 3 calcium ions per subunit. The ions are bound to the C2 domains.,function:May be involved in Ca(2+)-dependent exocytosis of secretory vesicles through Ca(2+) and phospholipid binding to the C2 domain or may serve as Ca(2+) sensors in the process of vesicular trafficking and exocytosis. Regulates the Ca(2+)-dependent secretion of norepinephrine in PC12 cells. Required for export from the endocytic recycling compartment to the cell surface.,similarity:Belongs to the synaptotagmin family.,similarity:Contains 2 C2
Synonyms Observed Band Cell Pathway Tissue Specificity	42kD Cytoplasmic vesicle, secretory vesicle, synaptic vesicle membrane ; Sin membrane protein . Recycling endosome membrane ; Single-pass mem protein . In mast cells, localizes to the endocytic recycling compartment. Brain, cofactor:Binds 3 calcium ions per subunit. The ions are bound to the C2


Nanjing BYabscience technology Co.,Ltd



国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务,欢迎咨询

Background	Synaptotagmins, such as SYT5, are a family of type III membrane proteins characterized by cytoplasmic repeats related to protein kinase C (see MIM 176960) regulatory (C2) domains, which are thought to bind calcium. Synaptotagmins may act both as negative regulators of vesicle fusion, allowing fusion in the presence of calcium, and as calcium receptors or sensor molecules (summary by Hudson and Birnbaum, 1995 [PubMed 7597049]).[supplied by OMIM, Feb 2011],
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using SYT5 Monoclonal Antibody

Nanjing BYabscience technology Co., Ltd