

TPC6B Monoclonal Antibody

BYmab-06288
D1111aD-00200
IgG
Human;Mouse
WB
TRAPPC6B
Trafficking protein particle complex subunit 6B
Synthesized peptide derived from part region of human protein
TPC6B Monoclonal Antibody detects endogenous levels of protein.
Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Monoclonal, Mouse,IgG
The antibody was affinity-purified from mouse antiserum by
affinity-chromatography using epitope-specific immunogen.
affinity-chromatography using epitope-specific immunogen. WB 1:500-2000
WB 1:500-2000
WB 1:500-2000 1 mg/ml
WB 1:500-2000 1 mg/ml ≥90%
WB 1:500-2000 1 mg/ml ≥90%
WB 1:500-2000 1 mg/ml ≥90% -20°C/1 year
WB 1:500-2000 1 mg/ml ≥90% -20°C/1 year 17kD
WB 1:500-2000 1 mg/ml ≥90% -20°C/1 year 17kD Golgi apparatus, cis-Golgi network . Endoplasmic reticulum . Both isoforms are expressed ubiquitously (at transcript level), isoform 1 being the most predominant (PubMed:28626029). Expressed in the fetal brain and different
WB 1:500-2000 1 mg/ml ≥90% -20°C/1 year 17kD Golgi apparatus, cis-Golgi network . Endoplasmic reticulum . Both isoforms are expressed ubiquitously (at transcript level), isoform 1 being the most predominant (PubMed:28626029). Expressed in the fetal brain and different regions of the adult brain and spinal cord (PubMed:28626029). function:May play a role in vesicular transport from endoplasmic reticulum to Golgi.,similarity:Belongs to the TRAPP small subunits family. BET3 subfamilysubunit:Part of the multisubunit TRAPP (transport protein particle)

Nanjing BYabscience technology Co.,Ltd




国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务, 欢迎咨询



Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.





Western Blot analysis of various cells using TPC6B Monoclonal Antibody

Nanjing BYabscience technology Co.,Ltd