

Catalog No



EI2BB Monoclonal Antibody

BYmab-06316


Catalog No	D1111ab-00510
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	EIF2B2 EIF2BB
Protein Name	Translation initiation factor eIF-2B subunit beta (S20I15) (S20III15) (eIF-2B GDP-GTP exchange factor subunit beta)
Immunogen	Synthesized peptide derived from human protein . at AA range: 50-130
Specificity	El2BB Monoclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	38kD
Cell Pathway	cytoplasm,cytosol,eukaryotic translation initiation factor 2B complex,
Tissue Specificity	Brain,Lung,Placenta,
Function	disease:Defects in EIF2B2 are a cause of leukodystrophy with vanishing white matter (VWM) [MIM:603896]. VWM is a leukodystrophy that occurs mainly in children. Neurological signs include progressive cerebellar ataxia, spasticity, inconstant optic atrophy and relatively preserved mental abilities. The disease is chronic-progressive with, in most individuals, additional episodes of rapid deterioration following febrile infections or minor head trauma. While childhood onset is the most common form of the disorder, some severe forms are apparent at birth. A severe, early-onset form seen among the Cree and Chippewayan populations of Quebec and Manitoba is called Cree leukoencephalopathy. Milder forms may not become evident until adolescence or adulthood. Some females with milder forms of the disease who survive to adolescence exhibit ovarian

Nanjing BYabscience technology Co.,Ltd

with milder forms of the disease who survive to adolescence exhibit ovarian dysfunction. This variant of the disorder is called

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658







Background	This gene encodes the beta subunit of eukaryotic initiation factor-2B (EIF2B). EIF2B is involved in protein synthesis and exchanges GDP and GTP for its activation and deactivation. [provided by RefSeq, Aug 2011],
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images Western Blot analysis of various cells using EI2BB 1 Monoclonal Antibody kDa 180 ---140 --100 --75 ---60 --45 --EI2BB 35 ---25 ---15 --10 --

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658