



IF2M Monoclonal Antibody

Catalog No	BYmab-06317
Isotype	lgG
Reactivity	Human;Mouse
Applications	WB
Gene Name	MTIF2
Protein Name	Translation initiation factor IF-2, mitochondrial (IF-2(Mt)) (IF-2Mt) (IF2(mt))
Immunogen	Synthesized peptide derived from human protein . at AA range: 450-530
Specificity	IF2M Monoclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	79kD
Cell Pathway	Mitochondrion.
Tissue Specificity	Expressed in all tissues examined. Highest level in skeletal muscle.
Function	function:One of the essential components for the initiation of protein synthesis. Protects formylmethionyl-tRNA from spontaneous hydrolysis and promotes its binding to the 30S ribosomal subunits. Also involved in the hydrolysis of GTP during the formation of the 70S ribosomal complex.,similarity:Belongs to the IF-2 family.,subunit:Monomer.,tissue specificity:Expressed in all tissues examined. Highest level in skeletal muscle.,
Background	During the initiation of protein biosynthesis, initiation factor-2 (IF-2) promotes the binding of the initiator tRNA to the small subunit of the ribosome in a GTP-dependent manner. Prokaryotic IF-2 is a single polypeptide, while eukaryotic cytoplasmic IF-2 (eIF-2) is a trimeric protein. Bovine liver mitochondria contain IF-2(mt), an 85-kD monomeric protein that is equivalent to prokaryotic IF-2. The
	Naniing Byabscience technology Co. Ltd

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658



国内优质抗体供应商

ШR 精准的 WB 检测服务 24H 在线服务,欢迎咨询

	predicted 727-amino acid human protein contains a 29-amino acid presequence. Human IF-2(mt) shares 32 to 38% amino acid sequence identity with yeast IF-2(mt) and several prokaryotic IF-2s, with the greatest degree of conservation in the G domains of the proteins. [provided by RefSeq, Mar 2016],
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.
Products Images	

Nanjing BYabscience technology Co.,Ltd