



UBAP1 Monoclonal Antibody


BYmab-06354
IgG
Human;Rat;Mouse
WB
UBAP1 NAG20
Ubiquitin-associated protein 1 (UBAP-1) (Nasopharyngeal carcinoma-associated gene 20 protein)
Synthesized peptide derived from part region of human protein
UBAP1 Monoclonal Antibody detects endogenous levels of protein.
Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Monoclonal, Mouse,IgG
The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
WB 1:500-2000
1 mg/ml
≥90%
-20°C/1 year
55kD
Cytoplasm, cytosol . Endosome . Predominantly cytosolic (PubMed:21757351). Recruited to endosomes as part of the ESCRT-I complex (PubMed:21757351)
Ubiquitous. Highly expressed in heart, brain, placenta, lung, liver, skeletal muscle and pancreas.
similarity:Contains 2 UBA domains.,tissue specificity:Ubiquitous. Highly expressed in heart, brain, placenta, lung, liver, skeletal muscle and pancreas.,
This gene is a member of the UBA domain family, whose members include proteins having connections to ubiquitin and the ubiquitination pathway. The ubiquitin associated domain is thought to be a non-covalent ubiquitin binding domain consisting of a compact three helix bundle. This particular protein originates from a gene locus in a refined region on chromosome 9 undergoing loss of heterozygosity in nasopharyngeal carcinoma (NPC). Taking into account its cytogenetic location, this UBA domain family member is being studies as a putative target for mutation in nasopharyngeal carcinomas. Multiple alternatively spliced transcript variants encoding different isoforms have been found for this

Nanjing BYabscience technology Co.,Ltd



matters needing attention

Usage suggestions





gene. [provided by RefSeq, Jan 2010],

Avoid repeated freezing and thawing!

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images

Nanjing BYabscience technology Co.,Ltd