



UB2E3 Monoclonal Antibody

| Catalog No | BYmab-06360 |
|--------------------|---|
| Isotype | lgG |
| Reactivity | Human;Mouse |
| Applications | WB |
| Gene Name | UBE2E3 UBCE4 UBCH9 |
| Protein Name | Ubiquitin-conjugating enzyme E2 E3 (EC 6.3.2.19) (UbcH9) (Ubiquitin carrier protein E3) (Ubiquitin-conjugating enzyme E2-23 kDa) (Ubiquitin-protein ligase E3) |
| Immunogen | Synthesized peptide derived from part region of human protein |
| Specificity | UB2E3 Monoclonal Antibody detects endogenous levels of protein. |
| Formulation | Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. |
| Source | Monoclonal, Mouse,IgG |
| Purification | The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. |
| Dilution | WB 1:500-2000 |
| Concentration | 1 mg/ml |
| Purity | ≥90% |
| Storage Stability | -20°C/1 year |
| Synonyms | |
| Observed Band | 22kD |
| Cell Pathway | Nucleus . Cytoplasm . Shuttles between the nucleus and cytoplasm in a IPO11-dependent manner. |
| Tissue Specificity | Ubiquitously expressed at low levels. Highly expressed in skeletal muscle. |
| Function | catalytic activity:ATP + ubiquitin + protein lysine = AMP + diphosphate + protein N-ubiquityllysine.,function:Catalyzes the covalent attachment of ubiquitin to other proteins. Participates in the regulation of transepithelial sodium transport in renal cells. May be involved in cell growth arrest.,pathway:Protein modification; protein ubiquitination.,similarity:Belongs to the ubiquitin-conjugating enzyme family.,subcellular location:Shuttles between the nucleus and cytoplasm in a IPO11-dependent manner.,subunit:The ubiquitin-loaded form interacts specifically with importin-11 (IPO11), leading to its import into the nucleus (By similarity). Interacts with NEDD4L.,tissue specificity:Ubiquitously expressed at low levels. Highly expressed in skeletal muscle., |

Nanjing BYabscience technology Co.,Ltd


| 博研生物 BYabscience |
|----------------------------|
|----------------------------|

国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务,欢迎咨询



| Background | The modification of proteins with ubiquitin is an important cellular mechanism for targeting abnormal or short-lived proteins for degradation. Ubiquitination involves at least three classes of enzymes: ubiquitin-activating enzymes, or E1s, ubiquitin-conjugating enzymes, or E2s, and ubiquitin-protein ligases, or E3s. This gene encodes a member of the E2 ubiquitin-conjugating enzyme family. The encoded protein shares 100% sequence identity with the mouse and rat counterparts, which indicates that this enzyme is highly conserved in eukaryotes. Multiple alternatively spliced transcript variants encoding the same protein have been found for this gene. [provided by RefSeq, Jun 2013], |
|---------------------------|--|
| matters needing attention | Avoid repeated freezing and thawing! |
| Usage suggestions | This product can be used in immunological reaction related experiments. For more information, please consult technical personnel. |

Products Images



Western Blot analysis of various cells using UB2E3 Monoclonal Antibody

Nanjing BYabscience technology Co.,Ltd