



SOGA2 Monoclonal Antibody

Catalog No	BYmab-06375
Isotype	IgG
Reactivity	Human;Mouse
Applications	WB
Gene Name	SOGA2 CCDC165 KIAA0802
Protein Name	Protein SOGA2 (Coiled-coil domain-containing protein 165)
Immunogen	Synthesized peptide derived from human protein . at AA range: 1770-1850
Specificity	SOGA2 Monoclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	209kD
Cell Pathway	Lateral cell membrane . Apical cell membrane . Cytoplasm, cytoskeleton, spindle pole . Midbody . Cytoplasm, cytoskeleton . Colocalized with microtubules at the base of cilia. Gradually accumulates on the apicobasal microtubule bundles during epithelial cell polarization (By similarity). Colocalized with the apicobasal microtubule bundles running beneath the lateral membrane. Colocalized with microtubule bundles in the spindle pole in mitotic cells and in the midbodies at the end of cytokinesis.
Tissue Specificity	Brain,Cerebellum,Epithelium,Testis,
Function	PTM:Isoform 4 is phosphorylated upon DNA damage, probably by ATM or ATR.,
Background	PTM:Isoform 4 is phosphorylated upon DNA damage, probably by ATM or ATR.,

Nanjing BYabscience technology Co.,Ltd



国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务, 欢迎咨询



matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images

Nanjing BYabscience technology Co.,Ltd