




## ZN281 Monoclonal Antibody

Catalog No         BYmab-06412           Isotype         IgG           Reactivity         Human;Mouse           Applications         WB           Gene Name         ZNF281 GZP1 ZBP99           Protein Name         Zinc finger protein 281 (GC-box-binding zinc finger protein 1) (Transcription factor ZBP-99) (Zinc finger DNA-binding protein 99)           Immunogen         Synthesized peptide derived from part region of human protein           Specificity         ZN281 Monoclonal Antibody detects endogenous levels of protein.           Formulation         Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.           Source         Monoclonal, Mouse, IgG           Purification         The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.           Dilution         WB 1:500-2000           Concentration         1 mg/ml           Purity         ≥90%           Storage Stability         -20*C/1 year           Synonyms         Observed Band           Observed Band         98kD           Cell Pathway         Nucleus .           Tissue Specificity         Liver,Placenta,Skin,           Function         function:Involved in transcriptional regulation. Represses the transcription of a number of genes including gastrin and ornithine decarboxylase. Binds to the G-rich box		
Applications  WB  Gene Name  ZNF281 GZP1 ZBP99  Protein Name  Zinc finger protein 281 (GC-box-binding zinc finger protein 1) (Transcription factor ZBP-99) (Zinc finger DNA-binding protein 99)  Immunogen  Synthesized peptide derived from part region of human protein  Specificity  ZN281 Monoclonal Antibody detects endogenous levels of protein.  Formulation  Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.  Source  Monoclonal, Mouse, IgG  Purification  The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution  WB 1:500-2000  Concentration  1 mg/ml  Purity  ≥90%  Storage Stability  -20°C/1 year  Synonyms  Observed Band  98kD  Cell Pathway  Nucleus .  Tissue Specificity  Liver, Placenta, Skin,  function:Involved in transcriptional regulation. Represses the transcription of a number of genes including gastrin and ornithine decarboxylase, Binds to the G-rich box in the enhancer region of these genes, PTM:Phosphorylated upon DNA damage, probably by ATM or ATR., similarity; Belongs to the krueppel including gastrin and ornithine decarboxylase. Binds to the G-rich box in the enhancer region of these genes. PTM:Phosphorylated upon DNA damage, probably by ATM or ATR., similarity; Belongs to the krueppel cingers.  Background	Catalog No	BYmab-06412
Applications  Gene Name  ZNF281 GZP1 ZBP99  Protein Name  Zinc finger protein 281 (GC-box-binding zinc finger protein 1) (Transcription factor ZBP-99) (Zinc finger DNA-binding protein 99)  Immunogen  Synthesized peptide derived from part region of human protein  Specificity  ZN281 Monoclonal Antibody detects endogenous levels of protein.  Formulation  Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.  Source  Monoclonal, Mouse, IgG  Purification  The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution  WB 1:500-2000  Concentration  1 mg/ml  Purity  ≥90%  Storage Stability  -20°C/1 year  Synonyms  Observed Band  98kD  Cell Pathway  Nucleus .  Tissue Specificity  Liver, Placenta, Skin,  function: Involved in transcriptional regulation. Represses the transcription of a number of genes including gastrin and ornithine decarboxylase. Binds to the G-rich box in the enhancer region of these genes. PTM: Phosphorylated upon DNA damage, probably by ATM or ATR, similarity, Belongs to the krueppel (2H2-type zinc fingers.)  Background  Function: Involved in transcriptional regulation. Represses the transcription of a number of genes including gastrin and ornithine decarboxylase. Binds to the G-rich box in the enhancer region of these genes. PTM: Phosphorylated upon DNA damage, probably by ATM or ATR, similarity. Belongs to the krueppel ornown and the enhancer region of these genes. PTM: Phosphorylated upon DNA damage, probably by ATM or ATR, similarity. Belongs to the krueppel ornown and the enhancer region of these genes. PTM: Phosphorylated upon DNA damage, probably by ATM or ATR, similarity. Belongs to the krueppel ornown and the enhancer region of these genes. PTM: Phosphorylated upon DNA damage, probably by ATM or ATR, similarity. Belongs to the krueppel	Isotype	IgG
Gene Name         ZNF281 GZP1 ZBP99           Protein Name         Zinc finger protein 281 (GC-box-binding zinc finger protein 1) (Transcription factor ZBP-99) (Zinc finger DNA-binding protein 99)           Immunogen         Synthesized peptide derived from part region of human protein           Specificity         ZN281 Monoclonal Antibody detects endogenous levels of protein.           Formulation         Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.           Source         Monoclonal, Mouse,lgG           Purification         The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.           Dilution         WB 1:500-2000           Concentration         1 mg/ml           Purity         ≥90%           Storage Stability         -20°C/1 year           Synonyms         Observed Band         98kD           Cell Pathway         Nucleus .           Tissue Specificity         Liver, Placenta, Skin,           Function         function: Involved in transcriptional regulation. Represses the transcription of a number of genes including gastrin and ornithine decarboxylase. Binds to the G-rich box in the enhancer region of these genes., PTM-Phosphorylated upon DNA damage, probably by ATM or ATR., similarity: Belongs to the krueppel of genes including gastrin and ornithine decarboxylase. Binds to the G-rich box in the enhancer region of these genese., PTM-Phosphorylated upon DNA damage, probably by ATM or ATR., similarity: Be	Reactivity	Human;Mouse
Protein Name  Zinc finger protein 281 (GC-box-binding zinc finger protein 1) (Transcription factor ZBP-99) (Zinc finger DNA-binding protein 99)  Immunogen  Synthesized peptide derived from part region of human protein  Specificity  ZN281 Monoclonal Antibody detects endogenous levels of protein.  Formulation  Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.  Source  Monoclonal, Mouse, IgG  Purification  The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution  WB 1:500-2000  Concentration  1 mg/ml  Purity  ≥90%  Storage Stability  -20°C/1 year  Synonyms  Observed Band  98kD  Cell Pathway  Nucleus .  Tissue Specificity  Liver, Placenta, Skin,  function: Involved in transcriptional regulation. Represses the transcription of a number of genes including gastrin and ornithine decarboxylase. Binds to the G-rich box in the enhancer region of these genes. PTM: Phosphorylated upon DNA damage, probably by ATM or ATR, similarity: Contains 4 C2H2-type zinc fingers.,  Background	Applications	WB
Immunogen         Synthesized peptide derived from part region of human protein           Specificity         ZN281 Monoclonal Antibody detects endogenous levels of protein.           Formulation         Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.           Source         Monoclonal, Mouse, IgG           Purification         The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.           Dilution         WB 1:500-2000           Concentration         1 mg/ml           Purity         ≥90%           Storage Stability         -20°C/1 year           Synonyms         Observed Band           Observed Band         98kD           Cell Pathway         Nucleus.           Tissue Specificity         Liver, Placenta, Skin,           Function         function:Involved in transcriptional regulation. Represses the transcription of a number of genes including gastrin and ornithine decarboxylase. Binds to the G-rich box in the enhancer region of these genes. PTM.Phosphorylated upon DNA damage, probably by ATM or ATR., similarity: Contains 4 C2H2-type zinc-fingers., function:Involved in transcriptional regulation. Represses the transcription of a number of genes including gastrin and ornithine decarboxylase. Binds to the G-rich box in the enhancer region of these genes. PTM.Phosphorylated upon DNA damage, probably by ATM or ATR, similarity:Gontains 4 C2H2-type zinc-fingers in the enhancer region of these genes. PTM.Phosphorylated upon DNA damage, probably by ATM or ATR, s	Gene Name	ZNF281 GZP1 ZBP99
Specificity ZN281 Monoclonal Antibody detects endogenous levels of protein.  Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.  Source Monoclonal, Mouse, IgG  Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-2000  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms  Observed Band 98kD  Cell Pathway Nucleus.  Tissue Specificity Liver, Placenta, Skin,  Function function: Involved in transcriptional regulation. Represses the transcription of a number of genes including gastrin and ornithine decarboxylase. Binds to the G-rich box in the enhancer region of these genes. PTM:Phosphorylated upon DNA damage, probably by ATM or ATR., similarity: Belongs to the krueppel C2H2-type zinc-finger protein family., similarity: Contains 4 C2H2-type zinc fingers.,  Background function: Involved in transcriptional regulation. Represses the transcription of a number of genes including gastrin and ornithine decarboxylase. Binds to the G-rich box in the enhancer region of these genes. PTM:Phosphorylated upon DNA damage, probably by ATM or ATR., similarity:Contains 4 C2H2-type zinc fingers.,  Function: Involved in transcriptional regulation. Represses the transcription of a number of genes including gastrin and ornithine decarboxylase. Binds to the G-rich box in the enhancer region of these genes., PTM:Phosphorylated upon DNA damage, probably by ATM or ATR, similarity Bolongs to the Krueppel C2H2-type zinc-finger protein family Similarity:Bolongs to the Krueppel C2H2-type zinc-finger protein family Similarity:Bolongs to the Krueppel C2H2-type zinc-finger protein family Similarity Bolongs to the Krueppel C2H2-type zinc-finger protein family Similarity Bolongs to the Krueppel C2H2-type zinc-finger protein family Similarity Bolongs to the Krueppel C2H2-type zinc-finger protein family Similarity Bolongs to the Krueppel C2H2-type zinc-finger protein family Similarity Bolo	Protein Name	Zinc finger protein 281 (GC-box-binding zinc finger protein 1) (Transcription factor ZBP-99) (Zinc finger DNA-binding protein 99)
Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.  Source Monoclonal, Mouse, IgG  Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-2000 Concentration 1 mg/ml  Purity ≥90% Storage Stability -20°C/1 year  Synonyms  Observed Band 98kD  Cell Pathway Nucleus.  Tissue Specificity Liver, Placenta, Skin,  function: Involved in transcriptional regulation. Represses the transcription of a number of genes including gastrin and ornithine decarboxylase. Binds to the G-rich box in the enhancer region of these genes. PTM: Phosphorylated upon DNA damage, probably by ATM or ATR. similarity: Belongs to the krueppel C2H2-type zinc-finger protein family., similarity: Contains 4 C2H2-type zinc-fingers, function: Involved in transcriptional regulation. Represses the transcription of a number of genes including gastrin and ornithine decarboxylase. Binds to the G-rich box in the enhancer region of these genes., PTM: Phosphorylated upon DNA damage, probably by ATM or ATR. similarity: Belongs to the krueppel G-rich box in the enhancer region of these genes., PTM: Phosphorylated upon DNA damage, probably by ATM or ATR. similarity: Belongs to the krueppel	Immunogen	Synthesized peptide derived from part region of human protein
Source       Monoclonal, Mouse,IgG         Purification       The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.         Dilution       WB 1:500-2000         Concentration       1 mg/ml         Purity       ≥90%         Storage Stability       -20°C/1 year         Synonyms       Observed Band       98kD         Cell Pathway       Nucleus .         Tissue Specificity       Liver,Placenta,Skin,         Function       function:Involved in transcriptional regulation. Represses the transcription of a number of genes including gastrin and ornithine decarboxylase. Binds to the G-rich box in the enhancer region of these genes. PTM:Phosphorylated upon DNA damage, probably by ATM or ATR., similarity:Contains 4 C2H2-type zinc fingers.,         Background       function:Involved in transcriptional regulation. Represses the transcription of a number of genes including gastrin and ornithine decarboxylase. Binds to the G-rich box in the enhancer region of these genes. PTM:Phosphorylated upon DNA damage, probably by ATM or ATR., similarity:Belongs to the krueppel	Specificity	ZN281 Monoclonal Antibody detects endogenous levels of protein.
Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year  Synonyms Observed Band 98kD Cell Pathway Nucleus .  Tissue Specificity Liver,Placenta,Skin,  Function function:Involved in transcriptional regulation. Represses the transcription of a number of genes including gastrin and ornithine decarboxylase. Binds to the G-rich box in the enhancer region of these genes.,PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Belongs to the Krueppel C2H2-type zinc-finger protein family.,similarity:Contains 4 C2H2-type zinc fingers.,  Background function:Involved in transcriptional regulation. Represses the transcription of a number of genes including gastrin and ornithine decarboxylase. Binds to the G-rich box in the enhancer region of these genes.,PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Belongs to the Krueppel G-rich box in the enhancer region of these genes.,PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Belongs to the Krueppel O-Rich box in the enhancer region of these genes.,PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Belongs to the Krueppel	Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-2000  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms  Observed Band 98kD  Cell Pathway Nucleus .  Tissue Specificity Liver, Placenta, Skin,  Function function:Involved in transcriptional regulation. Represses the transcription of a number of genes including gastrin and ornithine decarboxylase. Binds to the G-rich box in the enhancer region of these genes. PTM:Phosphorylated upon DNA damage, probably by ATM or ATR., similarity: Belongs to the krueppel C2H2-type zinc-fingers.,  Background function:Involved in transcriptional regulation. Represses the transcription of a number of genes including gastrin and ornithine decarboxylase. Binds to the G-rich box in the enhancer region of these genes. PTM:Phosphorylated upon DNA damage, probably by ATM or ATR., similarity: Belongs to the krueppel of the proper of these genes. PTM:Phosphorylated upon DNA damage, probably by ATM or ATR, similarity:Belongs to the krueppel or the enhancer region of these genes. PTM:Phosphorylated upon DNA damage, probably by ATM or ATR, similarity:Belongs to the krueppel or the enhancer region of these genes. PTM:Phosphorylated upon DNA damage, probably by ATM or ATR, similarity:Belongs to the krueppel	Source	Monoclonal, Mouse,IgG
Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms  Observed Band 98kD  Cell Pathway Nucleus .  Tissue Specificity Liver,Placenta,Skin,  Function function:Involved in transcriptional regulation. Represses the transcription of a number of genes including gastrin and ornithine decarboxylase. Binds to the G-rich box in the enhancer region of these genes.,PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Belongs to the krueppel C2H2-type zinc-finger protein family.,similarity:Contains 4 C2H2-type zinc fingers.,  Background function:Involved in transcriptional regulation. Represses the transcription of a number of genes including gastrin and ornithine decarboxylase. Binds to the G-rich box in the enhancer region of these genes.,PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Belongs to the krueppel	Purification	·
Purity ≥90%  Storage Stability -20°C/1 year  Synonyms  Observed Band 98kD  Cell Pathway Nucleus .  Tissue Specificity Liver,Placenta,Skin,  Function function:Involved in transcriptional regulation. Represses the transcription of a number of genes including gastrin and ornithine decarboxylase. Binds to the G-rich box in the enhancer region of these genes.,PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Belongs to the krueppel C2H2-type zinc-finger protein family.,similarity:Contains 4 C2H2-type zinc fingers.,  Background function:Involved in transcriptional regulation. Represses the transcription of a number of genes including gastrin and ornithine decarboxylase. Binds to the G-rich box in the enhancer region of these genes.,PTM:Phosphorylated upon DNA damage, probably by ATM or ATRsimilarity:Belongs to the krueppel	Dilution	WB 1:500-2000
Synonyms  Observed Band 98kD  Cell Pathway Nucleus .  Tissue Specificity Liver, Placenta, Skin,  Function function: Involved in transcriptional regulation. Represses the transcription of a number of genes including gastrin and ornithine decarboxylase. Binds to the G-rich box in the enhancer region of these genes., PTM: Phosphorylated upon DNA damage, probably by ATM or ATR., similarity: Belongs to the krueppel C2H2-type zinc-finger protein family., similarity: Contains 4 C2H2-type zinc fingers.,  Background function: Involved in transcriptional regulation. Represses the transcription of a number of genes including gastrin and ornithine decarboxylase. Binds to the G-rich box in the enhancer region of these genes., PTM: Phosphorylated upon DNA damage, probably by ATM or ATR., similarity: Belongs to the krueppel	Concentration	1 mg/ml
Synonyms  Observed Band 98kD  Cell Pathway Nucleus .  Tissue Specificity Liver, Placenta, Skin,  Function function: Involved in transcriptional regulation. Represses the transcription of a number of genes including gastrin and ornithine decarboxylase. Binds to the G-rich box in the enhancer region of these genes., PTM: Phosphorylated upon DNA damage, probably by ATM or ATR., similarity: Belongs to the krueppel C2H2-type zinc-finger protein family., similarity: Contains 4 C2H2-type zinc fingers.,  Background function: Involved in transcriptional regulation. Represses the transcription of a number of genes including gastrin and ornithine decarboxylase. Binds to the G-rich box in the enhancer region of these genes., PTM: Phosphorylated upon DNA damage, probably by ATM or ATR., similarity: Belongs to the krueppel	Purity	≥90%
Observed Band  Pathway  Nucleus .  Tissue Specificity  Liver,Placenta,Skin,  function:Involved in transcriptional regulation. Represses the transcription of a number of genes including gastrin and ornithine decarboxylase. Binds to the G-rich box in the enhancer region of these genes.,PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Belongs to the krueppel C2H2-type zinc-finger protein family.,similarity:Contains 4 C2H2-type zinc fingers.,  Background  function:Involved in transcriptional regulation. Represses the transcription of a number of genes including gastrin and ornithine decarboxylase. Binds to the G-rich box in the enhancer region of these genes.,PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Belongs to the krueppel	Storage Stability	-20°C/1 year
Tissue Specificity  Liver, Placenta, Skin,  function  function: Involved in transcriptional regulation. Represses the transcription of a number of genes including gastrin and ornithine decarboxylase. Binds to the G-rich box in the enhancer region of these genes. PTM: Phosphorylated upon DNA damage, probably by ATM or ATR., similarity: Belongs to the krueppel C2H2-type zinc-finger protein family., similarity: Contains 4 C2H2-type zinc fingers.,  Background  function: Involved in transcriptional regulation. Represses the transcription of a number of genes including gastrin and ornithine decarboxylase. Binds to the G-rich box in the enhancer region of these genes., PTM: Phosphorylated upon DNA damage, probably by ATM or ATR., similarity: Belongs to the krueppel	Synonyms	
Tissue Specificity  Liver, Placenta, Skin,  function: Involved in transcriptional regulation. Represses the transcription of a number of genes including gastrin and ornithine decarboxylase. Binds to the G-rich box in the enhancer region of these genes., PTM: Phosphorylated upon DNA damage, probably by ATM or ATR., similarity: Belongs to the krueppel C2H2-type zinc-finger protein family., similarity: Contains 4 C2H2-type zinc fingers.,  Background  function: Involved in transcriptional regulation. Represses the transcription of a number of genes including gastrin and ornithine decarboxylase. Binds to the G-rich box in the enhancer region of these genes., PTM: Phosphorylated upon DNA damage, probably by ATM or ATR., similarity: Belongs to the krueppel	Observed Band	98kD
Function  function:Involved in transcriptional regulation. Represses the transcription of a number of genes including gastrin and ornithine decarboxylase. Binds to the G-rich box in the enhancer region of these genes.,PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Belongs to the krueppel C2H2-type zinc-finger protein family.,similarity:Contains 4 C2H2-type zinc fingers.,  Background  function:Involved in transcriptional regulation. Represses the transcription of a number of genes including gastrin and ornithine decarboxylase. Binds to the G-rich box in the enhancer region of these genes.,PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Belongs to the krueppel	Cell Pathway	Nucleus .
number of genes including gastrin and ornithine decarboxylase. Binds to the G-rich box in the enhancer region of these genes.,PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Belongs to the krueppel C2H2-type zinc-finger protein family.,similarity:Contains 4 C2H2-type zinc fingers.,  Background  function:Involved in transcriptional regulation. Represses the transcription of a number of genes including gastrin and ornithine decarboxylase. Binds to the G-rich box in the enhancer region of these genes.,PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Belongs to the krueppel		
number of genes including gastrin and ornithine decarboxylase. Binds to the G-rich box in the enhancer region of these genes.,PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Belongs to the krueppel	Tissue Specificity	Liver,Placenta,Skin,
	•	function:Involved in transcriptional regulation. Represses the transcription of a number of genes including gastrin and ornithine decarboxylase. Binds to the G-rich box in the enhancer region of these genes.,PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Belongs to the krueppel C2H2-type zinc-finger protein family.,similarity:Contains 4 C2H2-type zinc

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658







	fingers.,
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## **Products Images**

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658