



## **ARAID Monoclonal Antibody**

Catalog No         BYmab-06423           Isotype         IgG           Reactivity         Human;Rat;Mouse;           Applications         WB           Gene Name         ATRAID APR3 C2orf28 HSPC013 UNQ214/PRO240           Protein Name         All-trans retinoic acid-induced differentiation factor (Apoptosis-related protein 3) (APR-3) (p18)           Immunogen         Synthesized peptide derived from human protein . at AA range: 1-80           Specificity         ARAID Monoclonal Antibody detects endogenous levels of protein.           Formulation         Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.           Source         Monoclonal, Mouse, IgG           Purification         The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.           Dilution         WB 1:500-2000           Concentration         1 mg/ml           Purity         290%           Storage Stability         -20°C/1 year           Synonyms         StbD           Cell Pathway         Nucleus envelope . Cell membrane ; Single-pass membrane protein . Colocalizes with NetL11 on the Learn envelope and the perinuclear region (PubMed: 21723284).           Tissue Specificity         Weakly expressed in hematopoietic cell lines.           Function         sequence caution: Translation N-terminally shortened, similarity: Contain		
Reactivity Human;Rat;Mouse;  Applications WB  Gene Name ATRAID APR3 C2orf28 HSPC013 UNQ214/PR0240  Protein Name All-trans retinoic acid-induced differentiation factor (Apoptosis-related protein 3) (APR-3) (p18)  Immunogen Synthesized peptide derived from human protein . at AA range: 1-80  Specificity ARAID Monoclonal Antibody detects endogenous levels of protein.  Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.  Source Monoclonal, Mouse,IgG  Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-2000  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms  Observed Band 25kD  Cell Pathway Nucleus envelope . Cell membrane ; Single-pass membrane protein . Colocalizes with NELL1 on the nuclear envelope and the perinuclear region (PubMed:21723284).  Tissue Specificity Weakly expressed in hematopoietic cell lines.  Function sequence caution:Translation N-terminally shortened, similarity:Contains 1 EGF-like domain.,tissue specificity:Weakly expressed in hematopoietic cell lines.  This gene is thought to be involved in apoptosis, and may also be involved in hematopoietic development and differentiation. The use of alternative splice sites and promotors result in multiple transcript variants encoding different isoforms.[provided by RefSeq, Dec 2009],  matters needing Avoid repeated freezing and thawing!	Catalog No	BYmab-06423
Applications WB  Gene Name ATRAID APR3 C2orf28 HSPC013 UNQ214/PRO240  Protein Name (All-trans retinoic acid-induced differentiation factor (Apoptosis-related protein 3) (APR-3) (p18)  Immunogen Synthesized peptide derived from human protein . at AA range: 1-80  Specificity ARAID Monoclonal Antibody detects endogenous levels of protein.  Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.  Source Monoclonal, Mouse, IgG  Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-2000  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms  Observed Band 25kD  Cell Pathway Nucleus envelope . Cell membrane ; Single-pass membrane protein . Colocalizes with NELL1 on the nuclear envelope and the perinuclear region (PubMed:21723284).  Tissue Specificity Weakly expressed in hematopoietic cell lines.  Function sequence caution:Translation N-terminally shortened, similarity:Contains 1 EGF-like domain.,tissue specificity:Weakly expressed in hematopoietic cell lines., This gene is thought to be involved in apoptosis, and may also be linvolved in hematopoietic cell perinutive splice sites and promotors result in multiple transcript variants encoding different isoforms.[provided by RefSeq, Dec 2009],	Isotype	IgG
Gene Name ATRAID APR3 C2orf28 HSPC013 UNQ214/PRO240  Protein Name (All-trans retinoic acid-induced differentiation factor (Apoptosis-related protein 3) (APR-3) (p18)  Immunogen Synthesized peptide derived from human protein . at AA range: 1-80  Specificity ARAID Monoclonal Antibody detects endogenous levels of protein.  Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.  Source Monoclonal, Mouse, IgG  Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-2000  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms  Observed Band 25kD  Cell Pathway Nucleus envelope . Cell membrane ; Single-pass membrane protein . Colocalizes with NELL1 on the nuclear envelope and the perinuclear region (PubMed:21723284).  Tissue Specificity Weakly expressed in hematopoietic cell lines.  Function sequence caution:Translation N-terminally shortened, similarity:Contains 1 EGF-like domain.,tissue specificity:Weakly expressed in hematopoietic cell lines., This gene is thought to be involved in apoptosis, and may also be linvolved in hematopoietic cell offerentiation. The use of alternative splice sites and promotors result in multiple transcript variants encoding different isoforms.[provided by RefSeq, Dec 2009],	Reactivity	Human;Rat;Mouse;
Protein Name All-trans retinoic acid-induced differentiation factor (Apoptosis-related protein 3) (APR-3) (p18)  Immunogen Synthesized peptide derived from human protein . at AA range: 1-80  Specificity ARAID Monoclonal Antibody detects endogenous levels of protein.  Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.  Source Monoclonal, Mouse, IgG  Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-2000  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms  Observed Band 25kD  Cell Pathway Nucleus envelope Cell membrane; Single-pass membrane protein . Colocalizes with NELL1 on the nuclear envelope and the perinuclear region (PubMed:21723284).  Tissue Specificity Weakly expressed in hematopoietic cell lines.  Function sequence caution:Translation N-terminally shortened.,similarity:Contains 1 EGF-like domain, tissue specificity: Weakly expressed in hematopoietic cell lines., This gene is thought to be involved in apoptosis, and may also be involved in hematopoietic development and differentiation. The use of alternative splice sites and promotors result in multiple transcript variants encoding different isoforms. [provided by RefSeq. Dec 2009],  matters needing Avoid repeated freezing and thawing!	Applications	WB
Immunogen   Synthesized peptide derived from human protein . at AA range: 1-80	Gene Name	ATRAID APR3 C2orf28 HSPC013 UNQ214/PRO240
Specificity ARAID Monoclonal Antibody detects endogenous levels of protein.  Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.  Source Monoclonal, Mouse, lgG  Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-2000  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms  Observed Band 25kD  Cell Pathway Nucleus envelope . Cell membrane ; Single-pass membrane protein . Colocalizes with NELL1 on the nuclear envelope and the perinuclear region (PubMed:21723284).  Tissue Specificity Weakly expressed in hematopoietic cell lines.  Function sequence caution:Translation N-terminally shortened., similarity:Contains 1 EGF-like domain., tissue specificity:Weakly expressed in hematopoietic cell lines, and may also be involved in hematopoietic development and differentiation. The use of alternative splice sites and promotors result in multiple transcript variants encoding different isoforms. [provided by RefSeq., Dec 2009], matters needing Avoid repeated freezing and thawing!	Protein Name	All-trans retinoic acid-induced differentiation factor (Apoptosis-related protein 3) (APR-3) (p18)
Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.  Source Monoclonal, Mouse, IgG  Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-2000 Concentration 1 mg/ml  Purity ≥90% Storage Stability -20°C/1 year  Synonyms  Observed Band 25kD  Cell Pathway Nucleus envelope . Cell membrane ; Single-pass membrane protein . Colocalizes with NELL1 on the nuclear envelope and the perinuclear region (PubMed:21723284).  Tissue Specificity Weakly expressed in hematopoietic cell lines.  Function Sequence caution:Translation N-terminally shortened., similarity:Contains 1 EGF-like domain., tissue specificity:Weakly expressed in hematopoietic cell lines.,  Background This gene is thought to be involved in apoptosis, and may also be involved in hematopoietic development and differentiation. The use of alternative splice sites and promotors result in multiple transcript variants encoding different isoforms.[provided by RefSeq., Dec 2009],  matters needing Avoid repeated freezing and thawing!	Immunogen	Synthesized peptide derived from human protein . at AA range: 1-80
Source       Monoclonal, Mouse,IgG         Purification       The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.         Dilution       WB 1:500-2000         Concentration       1 mg/ml         Purity       ≥90%         Storage Stability       -20°C/1 year         Synonyms         Observed Band       25kD         Cell Pathway       Nucleus envelope . Cell membrane ; Single-pass membrane protein . Colocalizes with NELL1 on the nuclear envelope and the perinuclear region (PubMed:21723284).         Tissue Specificity       Weakly expressed in hematopoietic cell lines.         Function       sequence caution:Translation N-terminally shortened.,similarity:Contains 1 EGF-like domain.,tissue specificity:Weakly expressed in hematopoietic cell lines.,         Background       This gene is thought to be involved in apoptosis, and may also be involved in hematopoietic development and differentiation. The use of alternative splice sites and promotors result in multiple transcript variants encoding different isoforms.[provided by RefSeq, Dec 2009],         matters needing       Avoid repeated freezing and thawing!	Specificity	ARAID Monoclonal Antibody detects endogenous levels of protein.
Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year  Synonyms  Observed Band 25kD  Cell Pathway Nucleus envelope . Cell membrane ; Single-pass membrane protein . Colocalizes with NELL1 on the nuclear envelope and the perinuclear region (PubMed:21723284).  Tissue Specificity Weakly expressed in hematopoietic cell lines.  Function Sequence caution:Translation N-terminally shortened.,similarity:Contains 1 EGF-like domain.,tissue specificity:Weakly expressed in hematopoietic cell lines.,  This gene is thought to be involved in apoptosis, and may also be involved in hematopoietic development and differentiation. The use of alternative splice sites and promotors result in multiple transcript variants encoding different isoforms.[provided by RefSeq, Dec 2009],  matters needing Avoid repeated freezing and thawing!	Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-2000  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms  Observed Band 25kD  Cell Pathway Nucleus envelope . Cell membrane ; Single-pass membrane protein . Colocalizes with NELL1 on the nuclear envelope and the perinuclear region (PubMed:21723284).  Tissue Specificity Weakly expressed in hematopoietic cell lines.  Function sequence caution:Translation N-terminally shortened.,similarity:Contains 1 EGF-like domain.,tissue specificity:Weakly expressed in hematopoietic cell lines.,  Background This gene is thought to be involved in apoptosis, and may also be involved in hematopoietic development and differentiation. The use of alternative splice sites and promotors result in multiple transcript variants encoding different isoforms.[provided by RefSeq, Dec 2009],  matters needing Avoid repeated freezing and thawing!	Source	Monoclonal, Mouse,IgG
Concentration       1 mg/ml         Purity       ≥90%         Storage Stability       -20°C/1 year         Synonyms       -25kD         Cell Pathway       Nucleus envelope . Cell membrane ; Single-pass membrane protein . Colocalizes with NELL1 on the nuclear envelope and the perinuclear region (PubMed:21723284).         Tissue Specificity       Weakly expressed in hematopoietic cell lines.         Function       sequence caution:Translation N-terminally shortened.,similarity:Contains 1 EGF-like domain.,tissue specificity:Weakly expressed in hematopoietic cell lines.,         Background       This gene is thought to be involved in apoptosis, and may also be involved in hematopoietic development and differentiation. The use of alternative splice sites and promotors result in multiple transcript variants encoding different isoforms.[provided by RefSeq, Dec 2009],         matters needing       Avoid repeated freezing and thawing!	Purification	
Purity ≥90%  Storage Stability -20°C/1 year  Synonyms  Observed Band 25kD  Cell Pathway Nucleus envelope . Cell membrane ; Single-pass membrane protein . Colocalizes with NELL1 on the nuclear envelope and the perinuclear region (PubMed:21723284).  Tissue Specificity Weakly expressed in hematopoietic cell lines.  Function sequence caution:Translation N-terminally shortened.,similarity:Contains 1 EGF-like domain.,tissue specificity:Weakly expressed in hematopoietic cell lines.,  This gene is thought to be involved in apoptosis, and may also be involved in hematopoietic development and differentiation. The use of alternative splice sites and promotors result in multiple transcript variants encoding different isoforms.[provided by RefSeq, Dec 2009],  matters needing Avoid repeated freezing and thawing!	Dilution	WB 1:500-2000
Storage Stability -20°C/1 year  Synonyms  Observed Band 25kD  Cell Pathway Nucleus envelope . Cell membrane ; Single-pass membrane protein . Colocalizes with NELL1 on the nuclear envelope and the perinuclear region (PubMed:21723284).  Tissue Specificity Weakly expressed in hematopoietic cell lines.  Function sequence caution:Translation N-terminally shortened.,similarity:Contains 1 EGF-like domain.,tissue specificity:Weakly expressed in hematopoietic cell lines.,  Background This gene is thought to be involved in apoptosis, and may also be involved in hematopoietic development and differentiation. The use of alternative splice sites and promotors result in multiple transcript variants encoding different isoforms.[provided by RefSeq, Dec 2009],  matters needing Avoid repeated freezing and thawing!	Concentration	1 mg/ml
Synonyms  Observed Band 25kD  Cell Pathway Nucleus envelope . Cell membrane ; Single-pass membrane protein . Colocalizes with NELL1 on the nuclear envelope and the perinuclear region (PubMed:21723284).  Tissue Specificity Weakly expressed in hematopoietic cell lines.  Function sequence caution:Translation N-terminally shortened.,similarity:Contains 1 EGF-like domain.,tissue specificity:Weakly expressed in hematopoietic cell lines.,  Background This gene is thought to be involved in apoptosis, and may also be involved in hematopoietic development and differentiation. The use of alternative splice sites and promotors result in multiple transcript variants encoding different isoforms.[provided by RefSeq, Dec 2009],  matters needing Avoid repeated freezing and thawing!		
Observed Band  Cell Pathway  Nucleus envelope . Cell membrane ; Single-pass membrane protein . Colocalizes with NELL1 on the nuclear envelope and the perinuclear region (PubMed:21723284).  Tissue Specificity  Weakly expressed in hematopoietic cell lines.  Function  sequence caution:Translation N-terminally shortened.,similarity:Contains 1 EGF-like domain.,tissue specificity:Weakly expressed in hematopoietic cell lines.,  This gene is thought to be involved in apoptosis, and may also be involved in hematopoietic development and differentiation. The use of alternative splice sites and promotors result in multiple transcript variants encoding different isoforms.[provided by RefSeq, Dec 2009],  matters needing  Avoid repeated freezing and thawing!	Purity	≥90%
Cell Pathway  Nucleus envelope . Cell membrane ; Single-pass membrane protein . Colocalizes with NELL1 on the nuclear envelope and the perinuclear region (PubMed:21723284).  Tissue Specificity  Weakly expressed in hematopoietic cell lines.  Function  sequence caution:Translation N-terminally shortened.,similarity:Contains 1 EGF-like domain.,tissue specificity:Weakly expressed in hematopoietic cell lines.,  This gene is thought to be involved in apoptosis, and may also be involved in hematopoietic development and differentiation. The use of alternative splice sites and promotors result in multiple transcript variants encoding different isoforms.[provided by RefSeq, Dec 2009],  matters needing  Avoid repeated freezing and thawing!		
(PubMed:21/23284).  Tissue Specificity  Weakly expressed in hematopoietic cell lines.  Function  sequence caution:Translation N-terminally shortened.,similarity:Contains 1 EGF-like domain.,tissue specificity:Weakly expressed in hematopoietic cell lines.,  This gene is thought to be involved in apoptosis, and may also be involved in hematopoietic development and differentiation. The use of alternative splice sites and promotors result in multiple transcript variants encoding different isoforms.[provided by RefSeq, Dec 2009],  Matters needing  Avoid repeated freezing and thawing!	Storage Stability	
Function  sequence caution:Translation N-terminally shortened., similarity:Contains 1 EGF-like domain., tissue specificity:Weakly expressed in hematopoietic cell lines.,  This gene is thought to be involved in apoptosis, and may also be involved in hematopoietic development and differentiation. The use of alternative splice sites and promotors result in multiple transcript variants encoding different isoforms.[provided by RefSeq, Dec 2009],  matters needing  Avoid repeated freezing and thawing!	Storage Stability Synonyms	-20°C/1 year
Background  This gene is thought to be involved in apoptosis, and may also be involved in hematopoietic development and differentiation. The use of alternative splice sites and promotors result in multiple transcript variants encoding different isoforms.[provided by RefSeq, Dec 2009],  Matters needing  EGF-like domain.,tissue specificity:Weakly expressed in hematopoietic cell lines.,  Avoid repeated freezing and thaving!	Storage Stability Synonyms Observed Band	-20°C/1 year  25kD  Nucleus envelope . Cell membrane ; Single-pass membrane protein . Colocalizes with NELL1 on the nuclear envelope and the perinuclear region
hematopoietic development and differentiation. The use of alternative splice sites and promotors result in multiple transcript variants encoding different isoforms.[provided by RefSeq, Dec 2009],  matters needing Avoid repeated freezing and thawing!	Storage Stability Synonyms Observed Band Cell Pathway	-20°C/1 year  25kD  Nucleus envelope . Cell membrane ; Single-pass membrane protein . Colocalizes with NELL1 on the nuclear envelope and the perinuclear region (PubMed:21723284).
	Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity	-20°C/1 year  25kD  Nucleus envelope . Cell membrane ; Single-pass membrane protein . Colocalizes with NELL1 on the nuclear envelope and the perinuclear region (PubMed:21723284).  Weakly expressed in hematopoietic cell lines.  sequence caution:Translation N-terminally shortened.,similarity:Contains 1
	Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function	-20°C/1 year  25kD  Nucleus envelope . Cell membrane ; Single-pass membrane protein . Colocalizes with NELL1 on the nuclear envelope and the perinuclear region (PubMed:21723284).  Weakly expressed in hematopoietic cell lines.  sequence caution:Translation N-terminally shortened.,similarity:Contains 1 EGF-like domain.,tissue specificity:Weakly expressed in hematopoietic cell lines.,  This gene is thought to be involved in apoptosis, and may also be involved in hematopoietic development and differentiation. The use of alternative splice sites and promotors result in multiple transcript variants encoding different

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658




国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务, 欢迎咨询



**Usage suggestions** 

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## **Products Images**



Western Blot analysis of various cells using ARAID Monoclonal Antibody