



TAB3 Monoclonal Antibody

Catalog No	BYmab-06488
Isotype	IgG
Reactivity	Human;Mouse
Applications	WB
Gene Name	TAB3 MAP3K7IP3
Protein Name	TGF-beta-activated kinase 1 and MAP3K7-binding protein 3 (Mitogen-activated protein kinase kinase kinase 7-interacting protein 3) (NF-kappa-B-activating protein 1) (TAK1-binding protein 3) (TAB-3) (TG
Immunogen	Synthesized peptide derived from part region of human protein
Specificity	TAB3 Monoclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	78kD
Cell Pathway	nucleoplasm,cytoplasm,cytosol,plasma membrane,endosome membrane,extracellular exosome,
Tissue Specificity	Widely expressed. Constitutively overexpressed in certain tumor tissues. ; [Isoform 1]: Major transcript. ; [Isoform 2]: Minor transcript.
Function	function:Adapter linking MAP3K7/TAK1 and TRAF6 or TRAF2. Mediator of MAP3K7 activation, respectively in the IL1 and TNF signaling pathways. Plays a role in activation of NF-kappa-B and AP1 transcription factor. Isoform 2 may be an oncogenic factor.,PTM:Phosphorylated.,PTM:Ubiquitinated; following IL1 stimulation or TRAF6 overexpression.,similarity:Contains 1 CUE domain.,similarity:Contains 1 RanBP2-type zinc finger.,subunit:Interacts with TAB1, TAB2, MAP3K7, TRAF2 and TRAF6. The minimal TAB3-containing complex (TAB1-MAP3K7-TAB3) appears not to contain TAB2. However, it seems sensible to consider that TAB2 may also join this complex and may act in a cooperative manner with TAB3.,tissue specificity:Widely expressed. Constitutively

Nanjing BYabscience technology Co.,Ltd



overexpressed in certain tumor tissues. Isoform 1 is a major transcript while isoform 2 is a minor transcript.,

Background

The product of this gene functions in the NF-kapMAB signal transduction pathway. The encoded protein, and the similar and functionally redundant protein MAP3K7IP2/TAB2, forms a ternary complex with the protein kinase MAP3K7/TAK1 and either TRAF2 or TRAF6 in response to stimulation with the pro-inflammatory cytokines TNF or IL-1. Subsequent MAP3K7/TAK1 kinase activity triggers a signaling cascade leading to activation of the NF-kapMAB transcription factor. The human genome contains a related pseudogene. Alternatively spliced transcript variants have been described, but their biological validity has not been determined. [provided by RefSeq, Jul 2008],

matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com

官方热线: 025-5229-8998

监督电话: 15950492658