



## TAF7L Monoclonal Antibody

BYmab-06499
lgG
Human;Rat;Mouse;
WB
TAF7L TAF2Q
Transcription initiation factor TFIID subunit 7-like (Cancer/testis antigen 40) (CT40) (RNA polymerase II TBP-associated factor subunit Q) (TATA box-binding protein-associated factor 50 kDa) (Transcri
Synthesized peptide derived from part region of human protein
TAF7L Monoclonal Antibody detects endogenous levels of protein.
Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Monoclonal, Mouse,IgG
The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
WB 1:500-2000
1 mg/ml
≥90%
-20°C/1 year
50kD
Nucleus . Cytoplasm . Cytoplasmic in spermatogonia and early spermatocytes (preleptotene, leptotene, and zygotene); translocates into the nuclei of pachytene spermatocytes and round spermatids
Testis-specific.
function:Probably functions as a spermatogensis-specific component of the DNA-binding general transcription factor complex TFIID, a multimeric protein complex that plays a central role in mediating promoter responses to various activators and repressors. May play a role in spermatogenesis.,similarity:Belongs to the TAF7 family.,subcellular location:Cytoplasmic in spermatogonia and early spermatocytes (preleptotene, leptotene, and zygotene); translocates into the nuclei of pachytene spermatocytes and round spermatids.,subunit:TFIID is composed of TATA binding protein (TBP) and a number of TBP-associated

## Nanjing BYabscience technology Co.,Ltd



国内优质抗体供应商

小小贞九件供应问 二

精准的 WB 检测服务 24H 在线服务, 欢迎咨询



	————
	TFIID. Interacts with TBP; the interaction occurs in a sub-population of cells (pachytene and haploid round spermatids) and is developmentally regulated through differential intracellular localization of the two
Background	This gene is similar to a mouse gene that encodes a TATA box binding protein-associated factor, and shows testis-specific expression. The encoded protein could be a spermatogenesis-specific component of the DNA-binding general transcription factor complex TFIID. Alternatively spliced transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Dec 2009],
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.
Products Images	

Nanjing BYabscience technology Co.,Ltd