



## **HES3 Monoclonal Antibody**

| Catalog No         | BYmab-06504  |
|--------------------|--|
| Isotype            | IgG  |
| Reactivity         | Human;Mouse;Rat  |
| Applications       | WB   |
| Gene Name          | HES3 BHLHB43   |
| Protein Name       | Transcription factor HES-3 (Class B basic helix-loop-helix protein 43) (bHLHb43) (Hairy and enhancer of split 3)     |
| Immunogen          | Synthesized peptide derived from human protein . at AA range: 20-100   |
| Specificity        | HES3 Monoclonal Antibody detects endogenous levels of protein.   |
| Formulation        | Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.   |
| Source             | Monoclonal, Mouse,IgG  |
| Purification       | The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. |
| Dilution           | WB 1:500-2000  |
| Concentration      | 1 mg/ml  |
| Purity             | ≥90%   |
| Storage Stability  | -20°C/1 year   |
| Synonyms           |  |
| Observed Band      | 20kD   |
| Cell Pathway       | Nucleus .  |
| Tissue Specificity |  |

**Function** 

domain:Has a particular type of basic domain (presence of a helix-interrupting proline) that binds to the N-box (CACNAG), rather than the canonical E-box (CANNTG).,domain:The C-terminal WRPW motif is a transcriptional repression domain necessary for the interaction with Groucho/TLE family members, transcriptional corepressors recruited to specific target DNA by Hairy-related proteins.,function:Transcriptional repressor of genes that require a bHLH protein for their transcription.,similarity:Contains 1 basic helix-loop-helix (bHLH) domain.,similarity:Contains 1 Orange domain.,subunit:Transcription repression requires formation of a complex with a corepressor protein of the Groucho/TLE family

family.,

Nanjing BYabscience technology Co., Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658



国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务,欢迎咨询



| Background                | domain:Has a particular type of basic domain (presence of a helix-interrupting proline) that binds to the N-box (CACNAG), rather than the canonical E-box (CANNTG).,domain:The C-terminal WRPW motif is a transcriptional repression domain necessary for the interaction with Groucho/TLE family members, transcriptional corepressors recruited to specific target DNA by Hairy-related proteins.,function:Transcriptional repressor of genes that require a bHLH protein for their transcription.,similarity:Contains 1 basic helix-loop-helix (bHLH) domain.,similarity:Contains 1 Orange domain.,subunit:Transcription repression requires formation of a complex with a corepressor protein of the Groucho/TLE family., |
|---------------------------|---|
| matters needing attention | Avoid repeated freezing and thawing!  |
| Usage suggestions         | This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.   |

## **Products Images** Western Blot analysis of various cells using HES3 1 Monoclonal Antibody kDa 180 --140 --100 --75 ---60 --45 ---35 --25 --HES3 15 --10 --

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658