



KDM4D Monoclonal Antibody

| | |
|--------------------|--|
| Catalog No | BYmab-06510 |
| Isotype | IgG |
| Reactivity | Human;Rat;Mouse; |
| Applications | WB |
| Gene Name | KDM4D JHDM3D JMJD2D |
| Protein Name | Lysine-specific demethylase 4D (EC 1.14.11.-) (JmjC domain-containing histone demethylation protein 3D) (Jumonji domain-containing protein 2D) |
| Immunogen | Synthesized peptide derived from human protein . at AA range: 400-480 |
| Specificity | KDM4D Monoclonal Antibody detects endogenous levels of protein. |
| Formulation | Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. |
| Source | Monoclonal, Mouse,IgG |
| Purification | The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. |
| Dilution | WB 1:500-2000 |
| Concentration | 1 mg/ml |
| Purity | ≥90% |
| Storage Stability | -20°C/1 year |
| Synonyms | |
| Observed Band | 57kD |
| Cell Pathway | Nucleus . |
| Tissue Specificity | Brain,Embryo,Teratocarcinoma,Testis, |
| Function | caution:It is uncertain whether Met-1 or Met-4 is the initiator.,cofactor:Binds 1 Fe(2+) ion per subunit.,function:Histone demethylase that specifically demethylates 'Lys-9' of histone H3, thereby playing a central role in histone code. Does not demethylate histone H3 'Lys-4', H3 'Lys-27', H3 'Lys-36' nor H4 'Lys-20'. Demethylates both di- and trimethylated H3 'Lys-9' residue, while it has no activity on monomethylated residues. Demethylation of Lys residue generates formaldehyde and succinate.,similarity:Belongs to the JHDM3 histone demethylase family.,similarity:Contains 1 JmjC domain.,similarity:Contains 1 JmjN domain., |
| Background | caution:It is uncertain whether Met-1 or Met-4 is the initiator.,cofactor:Binds 1 Fe(2+) ion per subunit.,function:Histone demethylase that specifically |

Nanjing BYabscience technology Co.,Ltd



demethylates 'Lys-9' of histone H3, thereby playing a central role in histone code. Does not demethylate histone H3 'Lys-4', H3 'Lys-27', H3 'Lys-36' nor H4 'Lys-20'. Demethylates both di- and trimethylated H3 'Lys-9' residue, while it has no activity on monomethylated residues. Demethylation of Lys residue generates formaldehyde and succinate.,similarity:Belongs to the JHDM3 histone demethylase family.,similarity:Contains 1 JmjC domain.,similarity:Contains 1 JmjN domain.,

matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com

官方热线: 025-5229-8998

监督电话: 15950492658