



# WWTR1 Monoclonal Antibody


Catalog No	BYmab-06569
Isotype	IgG
Reactivity	Human;Mouse
Applications	WB
Gene Name	WWTR1 TAZ
Protein Name	WW domain-containing transcription regulator protein 1 (Transcriptional coactivator with PDZ-binding motif)
Immunogen	Synthesized peptide derived from part region of human protein
Specificity	WWTR1 Monoclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	44kD
Cell Pathway	Nucleus . Cytoplasm . Cell membrane . Concentrates along specific portions of the plasma membrane, and accumulates in punctate nuclear bodies (By similarity). When phosphorylated, is retained in the cytoplasm by YWHAZ (By similarity). Can be retained in the nucleus by MED15 (PubMed:18568018). Localized in the cytoplasm in areas of epithelial cell high density (PubMed:21145499). At blastocyst stage expressed in the nucleus in trophectodermal cells, however expressed in the cytoplasm in the inner cell mass (By similarity). .
Tissue Specificity	Highly expressed in kidney, heart, placenta and lung. Expressed in the thyroid tissue.
Function	domain:Binds to transcription factors via its WW domain.,domain:The PDZ-binding motif is essential for stimulated gene transcription. It localizes the protein into both punctate nuclear foci and plasma membrane-associated complexes.,function:Functions as a transcriptional coactivator.,PTM:Phosphorylated. Phosphorylation results in the inhibition of

Nanjing BYabscience technology Co.,Ltd



	transcriptional coactivation through YWHAZ-mediated nuclear export.,similarity:Contains 1 WW domain.,subcellular location:Concentrates along specific portions of the plasma membrane, and accumulates in punctate nuclear bodies. When phosphorylated, is retained in cytoplasm by YWHAZ.,subunit:Binds to SLC9A3R2 via the PDZ motif at the plasma membrane. Binds to YWHAZ in vivo and in vitro through the phosphoserine-binding motif RSHSSP.,tissue specificity:Highly expressed in kidney, heart, placenta and lung.,
<b>Background</b>	domain:Binds to transcription factors via its WW domain.,domain:The PDZ-binding motif is essential for stimulated gene transcription. It localizes the protein into both punctate nuclear foci and plasma membrane-associated complexes.,function:Functions as a transcriptional coactivator.,PTM:Phosphorylated. Phosphorylation results in the inhibition of transcriptional coactivation through YWHAZ-mediated nuclear export.,similarity:Contains 1 WW domain.,subcellular location:Concentrates along specific portions of the plasma membrane, and accumulates in punctate nuclear bodies. When phosphorylated, is retained in cytoplasm by YWHAZ.,subunit:Binds to SLC9A3R2 via the PDZ motif at the plasma membrane. Binds to YWHAZ in vivo and in vitro through the phosphoserine-binding motif RSHSSP.,tissue specificity:Highly expressed in kidney, heart, placenta and lung.,
<b>matters needing attention</b>	Avoid repeated freezing and thawing!
<b>Usage suggestions</b>	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## Products Images



Western Blot analysis of various cells using WWTR1 Monoclonal Antibody

Nanjing BYabs science technology Co.,Ltd

网址: [www.njbybio.com](http://www.njbybio.com)

官方热线: 025-5229-8998

监督电话: 15950492658