



# TF2L1 Monoclonal Antibody

|                           |   |
|---------------------------|---|
| <b>Catalog No</b>         | BYmab-06581   |
| <b>Isotype</b>            | IgG   |
| <b>Reactivity</b>         | Human;Mouse   |
| <b>Applications</b>       | WB  |
| <b>Gene Name</b>          | TFCP2L1 CRTR1 LBP9  |
| <b>Protein Name</b>       | Transcription factor CP2-like protein 1 (CP2-related transcriptional repressor 1) (CRTR-1) (Transcription factor LBP-9)   |
| <b>Immunogen</b>          | Synthesized peptide derived from part region of human protein   |
| <b>Specificity</b>        | TF2L1 Monoclonal Antibody detects endogenous levels of protein.   |
| <b>Formulation</b>        | Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.  |
| <b>Source</b>             | Monoclonal, Mouse,IgG   |
| <b>Purification</b>       | The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.  |
| <b>Dilution</b>           | WB 1:500-2000   |
| <b>Concentration</b>      | 1 mg/ml   |
| <b>Purity</b>             | ≥90%  |
| <b>Storage Stability</b>  | -20°C/1 year  |
| <b>Synonyms</b>           |   |
| <b>Observed Band</b>      | 52kD  |
| <b>Cell Pathway</b>       | Nucleus .   |
| <b>Tissue Specificity</b> | Highly expressed in placental JEG-3 cells and very low levels of expression in non-steroidogenic cells. No expression was seen in adrenal NCI-H295A cells or in adrenal tissue.   |
| <b>Function</b>           | function:Transcriptional suppressor. May suppress UBP1-mediated transcriptional activation. Modulates the placental expression of CYP11A1.,similarity:Belongs to the grh/CP2 family. CP2 subfamily.,tissue specificity:Highly expressed in placental JEG-3 cells and very low levels of expression in non-steroidogenic cells. No expression was seen in adrenal NCI-H295A cells or in adrenal tissue., |
| <b>Background</b>         | function:Transcriptional suppressor. May suppress UBP1-mediated transcriptional activation. Modulates the placental expression of CYP11A1.,similarity:Belongs to the grh/CP2 family. CP2 subfamily.,tissue specificity:Highly expressed in placental JEG-3 cells and very low levels of expression in non-steroidogenic cells. No   |

**Nanjing BYabscience technology Co.,Ltd**



expression was seen in adrenal NCI-H295A cells or in adrenal tissue.,

**matters needing attention**

Avoid repeated freezing and thawing!

**Usage suggestions**

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## Products Images

Nanjing BYabscience technology Co.,Ltd

网址: [www.njbybio.com](http://www.njbybio.com)

官方热线: 025-5229-8998

监督电话: 15950492658