





## **GRIP2** Monoclonal Antibody

Catalog No	BYmab-06614
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	WB
Gene Name	GRIP2 KIAA1719
Protein Name	Glutamate receptor-interacting protein 2 (GRIP-2)
Immunogen	Synthesized peptide derived from human protein . at AA range: 780-860
Specificity	GRIP2 Monoclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	114kD
Cell Pathway	Cytoplasm . Membrane ; Peripheral membrane protein .
Tissue Specificity	Brain,Spleen,
Function	domain:PDZ 5 mediates the C-terminal binding of GRIA2 and GRIA3. PDZ 6 mediates interaction with the PDZ recognition motif of EFNB1. PDZ 7 mediates interaction with CSPG4.,function:May play a role as a localized scaffold for the assembly of a multiprotein signaling complex and as mediator of the trafficking of its binding partners at specific subcellular location in neurons.,similarity:Belongs to the GRIP2 family.,similarity:Contains 7 PDZ (DHR) domains.,subunit:Interacts with EFNB1, EFNB3, GRIA2, GRIA3, CSPG4 and GRIPAP1. Can form homomultimers and heteromultimers with GRIP1 (By similarity). Interacts with the C-terminal tail of PRLHR.,
Background	domain:PDZ 5 mediates the C-terminal binding of GRIA2 and GRIA3. PDZ 6 mediates interaction with the PDZ recognition motif of EFNB1. PDZ 7 mediates

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658



国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务,欢迎咨询



	interaction with CSPG4.,function:May play a role as a localized scaffold for the assembly of a multiprotein signaling complex and as mediator of the trafficking of its binding partners at specific subcellular location in neurons.,similarity:Belongs to the GRIP2 family.,similarity:Contains 7 PDZ (DHR) domains.,subunit:Interacts with EFNB1, EFNB3, GRIA2, GRIA3, CSPG4 and GRIPAP1. Can form homomultimers and heteromultimers with GRIP1 (By similarity). Interacts with the C-terminal tail of PRLHR.,
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## **Products Images**

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658