



## HS74L Monoclonal Antibody

Catalog No       BYmab-06617         Isotype       IgG         Reactivity       Human;Mouse         Applications       WB         Gene Name       HSPA4L APG1 OSP94         Protein Name       Heat shock 70 kDa protein 4L (Heat shock 70-related protein APG-1) (Osmotic stress protein 94)         Immunogen       Synthesized peptide derived from human protein . at AA range: 450-530         Specificity       HS74L Monoclonal Antibody detects endogenous levels of protein.         Formulation       Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.         Source       Monoclonal, Mouse, IgG         Purification       The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.         Dilution       WB 1:500-2000         Concentration       1 mg/ml         Purity       290%         Storage Stability       -20°C/1 year         Synonyms       -20°C/1 year         Observed Band       92kD         Call Pathway       Cytoplasm . Nucleus . May translocate to the nucleus after heat shock		
Reactivity       Human;Mouse         Applications       WB         Gene Name       HSPA4L APG1 OSP94         Protein Name       Heat shock 70 kDa protein 4L (Heat shock 70-related protein APG-1) (Osmotic stress protein 94)         Immunogen       Synthesized peptide derived from human protein . at AA range: 450-530         Specificity       HS74L Monoclonal Antibody detects endogenous levels of protein.         Formulation       Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.         Source       Monoclonal, Mouse, IgG         Purification       The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.         Dilution       WB 1:500-2000         Concentration       1 mg/ml         Purity       ≥90%         Storage Stability       -20°C/1 year         Synonyms       -20°C/1 year         Observed Band       92kD         Cell Pathway       Cytoplasm . Nucleus . May translocate to the nucleus after heat shock         Tissue Specificity       Heart, Testis,         Function       function: Possesses chaperone activity in vitro where it inhibits aggregation of citrate synthase. Induction: May translocate to the nucleus after heat shock damage, probably by ATM or ATR_similarity Belongs to the heat shock reprint shock. subunit: Homodimer.         Background       The protein encoded by	Catalog No	BYmab-06617
ApplicationsWBGene NameHSPA4L APG1 OSP94Protein NameHeat shock 70 kDa protein 4L (Heat shock 70-related protein APG-1) (Osmotic stress protein 94)ImmunogenSynthesized peptide derived from human protein . at AA range: 450-530SpecificityHS74L Monoclonal Antibody detects endogenous levels of protein.FormulationLiquid in PBS containing 50% glycerol, and 0.02% sodium azide.SourceMonoclonal, Mouse.lgGPurificationThe antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.DilutionWB 1:500-2000Concentration1 mg/mlPurity≥90%Storage Stability-20°C/1 yearSynonymsCytoplasm . Nucleus . May translocate to the nucleus after heat shockTissue SpecificityHeart, Testis,Functionfunction:Possesses chaperone activity in vitro where it inhibits aggregation of citrate synthase, induction:By heat shock, PTM:Phosphorylated upon DNA damage, probably by ATM or ATR, similarity Belongs to the heat shock protein 70 family, subcellular location. May translocate to the nucleus after heat shock protein 70 family, subcellular location. May translocate to the nucleus after heat shock, subunit:Homodimer.,BackgroundThe protein encoded by this gene is heat shock inducible and may act as a chaperone. The encoded protein S. This gene is highly expressed in leukemia cells and may be a good target for therapeutic intervention. Several transcripts encoding different isoforms have been found for this gene is highly expressed in leukemia cells and may be a good target for therapeutic intervention. Several transcripts encoding dif	Isotype	lgG
Gene Name       HSPA4L APG1 OSP94         Protein Name       Heat shock 70 kDa protein 4L (Heat shock 70-related protein APG-1) (Osmotic stress protein 94)         Immunogen       Synthesized peptide derived from human protein . at AA range: 450-530         Specificity       HS74L Monoclonal Antibody detects endogenous levels of protein.         Formulation       Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.         Source       Monoclonal, Mouse, IgG         Purification       The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.         Dilution       WB 1:500-2000         Concentration       1 mg/ml         Purity       ≥90%         Storage Stability       -20°C/1 year         Synonyms       Observed Band         Okserved Band       92kD         Cell Pathway       Cytoplasm . Nucleus . May translocate to the nucleus after heat shock         Tissue Specificity       Heart, Testis,         Function       function:Possesses chaperone activity in vitro where it inhibits aggregation of citrate synthase, induction:By heat shock, PTM:Phosphonylated upon DNA dramage, probably by ATM or ATR, similarity-Ebengs to the heat shock protein 70 family. subcellular location:May translocate to the nucleus after heat shock protein 70 family. subcellular location:May translocate to the nucleus after heat shock are cell against the harmful effects of aggregated proteins. This gene is highly ex	Reactivity	Human;Mouse
Protein Name       Heat shock 70 kDa protein 4L (Heat shock 70-related protein APG-1) (Osmotic stress protein 94)         Immunogen       Synthesized peptide derived from human protein . at AA range: 450-530         Specificity       HS74L Monoclonal Antibody detects endogenous levels of protein.         Formulation       Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.         Source       Monoclonal, Mouse, IgG         Purification       The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.         Dilution       WB 1:500-2000         Concentration       1 mg/ml         Purity       ≥90%         Storage Stability       -20°C/1 year         Synonyms       Cytoplasm . Nucleus . May translocate to the nucleus after heat shock         Tissue Specificity       Heart, Testis,         Function       function:Possesses chaperone activity in vitro where it inhibits aggregation of citrate synthase, induction:By heat shock, PTM:Phosphorylated upon DNA damage, probably by ATM or ATR., similarity:Belongs to the heat shock protein 70 familysubcellular location:May translocate to the nucleus after heat a clagainst the hardful effects of aggregated proteins. This gene is heat shock inducible and may act as a chaperone. The encoded by this gene is heat shock inducible and may act as a chaperone. The encoded by this gene is heat shock inducible and may act as a chaperone. The encoded by this gene is heat shock inducible and may act as a chaperone. The encoded by this gene is heat shock inducibl	Applications	WB
stress protein 94)ImmunogenSynthesized peptide derived from human protein . at AA range: 450-530SpecificityHS74L Monoclonal Antibody detects endogenous levels of protein.FormulationLiquid in PBS containing 50% glycerol, and 0.02% sodium azide.SourceMonoclonal, Mouse,IgGPurificationThe antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.DilutionWB 1:500-2000Concentration1 mg/mlPurity≥90%Storage Stability-20°C/1 yearSynonymsCytoplasm . Nucleus . May translocate to the nucleus after heat shockTissue SpecificityHeart, Testis,Functionfunction:Possesse chaperone activity in vitro where it inhibits aggregation of citrate synthase.induction:May translocate to the nucleus after heat shock protein 70 familysubcellular location.May translocate to the nucleus after heat shock, subunit:Homodimer.,BackgroundThe protein encoded by this gene is heat shock inducible and may act as a chaperone. The encoded protein can protein can protein the served in leukemia cells and may be a good target for therapeutic intervention. Several transcripts encoding different isoforms have been found for this gene. [provided by RefSeq.	Gene Name	HSPA4L APG1 OSP94
Specificity       HS74L Monoclonal Antibody detects endogenous levels of protein.         Formulation       Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.         Source       Monoclonal, Mouse,IgG         Purification       The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.         Dilution       WB 1:500-2000         Concentration       1 mg/ml         Purity       ≥90%         Storage Stability       -20°C/1 year         Synonyms       -20°C/1 year         Observed Band       92kD         Cell Pathway       Cytoplasm. Nucleus . May translocate to the nucleus after heat shock         Function       function:Possesses chaperone activity in vitro where it inhibits aggregation of citrate synthaseInduction:By heat shock., PTM:Phosphorylated upon DNA damage, probably by ATM or ATR., similarity:Belongs to the heat shock protein 70 family. subcellular location.May translocate to the nucleus after heat shock protein 70 family. Subcellular location.May translocate to the nucleus after heat shock protein 70 family. Subcellular location May translocate to the nucleus after heat shock rotein activity in vitro where it inhibits aggregation of citrate synthaseInduction By heat shock. TM:Phosphorylated upon DNA damage, probably by ATM or ATR., similarity:Belongs to the heat shock protein 70 family. Subcellular location.May translocate to the nucleus after heat shock rotein 70 family. Subcellular location is heat shock inducible and may act as a chaperone. The encoded proteins. This gene is highly expressed in leukemi	Protein Name	Heat shock 70 kDa protein 4L (Heat shock 70-related protein APG-1) (Osmotic stress protein 94)
Formulation       Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.         Source       Monoclonal, Mouse,IgG         Purification       The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.         Dilution       WB 1:500-2000         Concentration       1 mg/ml         Purity       ≥90%         Storage Stability       -20°C/1 year         Synonyms       -20°C/1 year         Observed Band       92kD         Cell Pathway       Cytoplasm . Nucleus . May translocate to the nucleus after heat shock         Tissue Specificity       Heart, Testis,         Function       function:Possesses chaperone activity in vitro where it inhibits aggregation of citrate synthaseinduction:By heat shockPTM:Phosphorylated upon DNA damage, probably by ATM or ATRsimilarity:Belongs to the heat shock protein 70 familysubcellular location:May translocate to the nucleus after heat shock protein 70 familysubcellular location:May translocate to the nucleus after heat shock subunit:Homodimer.,         Background       The protein encoded protein can protect the heat shock dagainst the harmful effects of aggregated proteins. This gene is highly expressed in leukemia cells and may be a good target for therapeutic intervention. Several transcripts encoding different isoforms have been found for this gene. [provided by RefSeq, Dec 2015],	Immunogen	Synthesized peptide derived from human protein . at AA range: 450-530
Source       Monoclonal, Mouse,IgG         Purification       The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.         Dilution       WB 1:500-2000         Concentration       1 mg/ml         Purity       ≥90%         Storage Stability       -20°C/1 year         Synonyms       -20°C/1 year         Observed Band       92kD         Cell Pathway       Cytoplasm . Nucleus . May translocate to the nucleus after heat shock         Tissue Specificity       Heart,Testis,         Function       function:Possesses chaperone activity in vitro where it inhibits aggregation of citrate synthase.,induction:By heat shockPTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity.Belongs to the heat shock protein 70 familysubcellular location:May translocate to the nucleus after heat shock protein 70 familysubcellular location:May translocate to the nucleus after heat shock protein 70 familysubcellular location:May translocate to the nucleus after heat shock protein 70 familysubcellular location:May translocate to the nucleus after heat shock protein 70 familysubcellular location:May translocate to the nucleus after heat shock protein for family .subcellular location:May translocate to the nucleus after heat shock protein 70 familysubcellular location:May translocate to the nucleus after heat shock protein 70 familysubcellular location:May translocate to the nucleus after heat shock protein for family effects of aggregated proteins. This gene is highly expressed in leukemia cells and may be a good target for therapeutic intervention. Several trans	Specificity	HS74L Monoclonal Antibody detects endogenous levels of protein.
Purification       The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.         Dilution       WB 1:500-2000         Concentration       1 mg/ml         Purity       ≥90%         Storage Stability       -20°C/1 year         Synonyms       -20°C/1 year         Observed Band       92kD         Cell Pathway       Cytoplasm . Nucleus . May translocate to the nucleus after heat shock         Tissue Specificity       Heart, Testis,         Function       function:Possesses chaperone activity in vitro where it inhibits aggregation of citrate synthase, induction:By heat shock, .PTM:Phosphorylated upon DNA damage, probably by ATM or ATR., similarity:Belongs to the heat shock protein 70 family, subcellular location.May translocate to the nucleus after heat shock protein 70 family, subcellular location:May translocate to the nucleus after heat shock protein 70 family. subcellular location indep roteins. This gene is heat shock cell against the harmful effects of aggregated proteins. This gene is highly expressed in leukemia cells and may be a good target for therapeutic intervention. Several transcripts encoding different isoforms have been found for this gene. [provided by RefSeq. Dec 2015],	Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
affinity-chromatography using epitope-specific immunogen.         Dilution       WB 1:500-2000         Concentration       1 mg/ml         Purity       ≥90%         Storage Stability       -20°C/1 year         Synonyms       -20°C/1 year         Observed Band       92kD         Cell Pathway       Cytoplasm . Nucleus . May translocate to the nucleus after heat shock         Tissue Specificity       Heart, Testis,         Function       function:Possesses chaperone activity in vitro where it inhibits aggregation of citrate synthase., induction:By heat shock., PTM:Phosphorylated upon DNA damage, probably by ATM or ATR., similarity:Belongs to the heat shock protein 70 family., subcellular location:May translocate to the nucleus after heat shock protein 70 family. subcellular location:May translocate to the nucleus after heat shock protein 70 family. subcellular location:May translocate to the nucleus after heat shock protein 70 family. subcellular location:May translocate to the nucleus after heat shock inducible and may act as a chaperone. The encoded protein can protect the heat-shocked cell against the hamful effects of aggregated proteins. This gene is highly expressed in leukemia cells and may be a good target for therapeutic intervention. Several transcripts encoding different isoforms have been found for this gene. [provided by RefSeq, Dec 2015],	Source	Monoclonal, Mouse,IgG
Concentration       1 mg/ml         Purity       ≥90%         Storage Stability       -20°C/1 year         Synonyms       -20°C/1 year         Observed Band       92kD         Cell Pathway       Cytoplasm . Nucleus . May translocate to the nucleus after heat shock         Tissue Specificity       Heart,Testis,         Function       function:Possesses chaperone activity in vitro where it inhibits aggregation of citrate synthase.,induction:By heat shock.,PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Belongs to the heat shock protein 70 family.,subcellular location:May translocate to the nucleus after heat shock code against the harmful effects of aggregated proteins. This gene is heat shock inducible and may act as a chaperone. The encoded by this gene is heat shock inducible and may act as a chaperone. The encoded protein can protect the heat-shocked cell against the harmful effects of aggregated proteins. This gene is highly expressed in leukemia cells and may be a good target for therapeutic intervention. Several transcripts encoding different isoforms have been found for this gene. [provided by RefSeq, Dec 2015],	Purification	
Purity       ≥90%         Storage Stability       -20°C/1 year         Synonyms       -20°C/1 year         Observed Band       92kD         Cell Pathway       Cytoplasm . Nucleus . May translocate to the nucleus after heat shock         Tissue Specificity       Heart,Testis,         Function       function:Possesses chaperone activity in vitro where it inhibits aggregation of citrate synthase, induction:By heat shockPTM:Phosphorylated upon DNA damage, probably by ATM or ATRsimilarity:Belongs to the heat shock protein 70 familysubcellular location:May translocate to the nucleus after heat shock protein 70 familysubcellular location:May translocate to the nucleus after heat shock protein for this gene. In encoded by this gene is heat shock inducible and may act as a chaperone. The encoded protein can protect the heat-shocked cell against the harmful effects of aggregated proteins. This gene is highly expressed in leukemia cells and may be a good target for therapeutic intervention. Several transcripts encoding different isoforms have been found for this gene. [provided by RefSeq, Dec 2015],	Dilution	WB 1:500-2000
Storage Stability       -20°C/1 year         Synonyms       -20°C/1 year         Observed Band       92kD         Cell Pathway       Cytoplasm . Nucleus . May translocate to the nucleus after heat shock         Tissue Specificity       Heart, Testis,         Function       function:Possesses chaperone activity in vitro where it inhibits aggregation of citrate synthase.,induction:By heat shock.,PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Belongs to the heat shock protein 70 family.,subcellular location:May translocate to the nucleus after heat shock.,subunit:Homodimer.,         Background       The protein encoded by this gene is heat shock inducible and may act as a chaperone. The encoded protein can protect the heat-shocked cell against the harmful effects of aggregated proteins. This gene is highly expressed in leukemia cells and may be a good target for therapeutic intervention. Several transcripts encoding different isoforms have been found for this gene. [provided by RefSeq, Dec 2015],	Concentration	1 mg/ml
Synonyms         Observed Band       92kD         Cell Pathway       Cytoplasm . Nucleus . May translocate to the nucleus after heat shock         Tissue Specificity       Heart,Testis,         Function       function:Possesses chaperone activity in vitro where it inhibits aggregation of citrate synthase.,induction:By heat shock.,PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Belongs to the heat shock protein 70 family.,subcellular location:May translocate to the nucleus after heat shock protein 70 family.,subcellular location:May translocate to the nucleus after heat shock inducible and may act as a chaperone. The encoded by this gene is heat shock inducible and may act as a chaperone. The encoded protein can protect the heat-shocked cell against the harmful effects of aggregated proteins. This gene is highly expressed in leukemia cells and may be a good target for therapeutic intervention. Several transcripts encoding different isoforms have been found for this gene. [provided by RefSeq, Dec 2015],	Purity	≥90%
Observed Band92kDCell PathwayCytoplasm . Nucleus . May translocate to the nucleus after heat shockTissue SpecificityHeart,Testis,Functionfunction:Possesses chaperone activity in vitro where it inhibits aggregation of citrate synthase.,induction:By heat shock.,PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Belongs to the heat shock protein 70 family.,subcellular location:May translocate to the nucleus after heat shock.,subunit:Homodimer.,BackgroundThe protein encoded by this gene is heat shock inducible and may act as a chaperone. The encoded protein can protect the heat-shocked cell against the harmful effects of aggregated proteins. This gene is highly expressed in leukemia cells and may be a good target for therapeutic intervention. Several transcripts encoding different isoforms have been found for this gene. [provided by RefSeq, Dec 2015],	Storage Stability	-20°C/1 year
Cell PathwayCytoplasm . Nucleus . May translocate to the nucleus after heat shockTissue SpecificityHeart, Testis,Functionfunction:Possesses chaperone activity in vitro where it inhibits aggregation of citrate synthase.,induction:By heat shock.,PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Belongs to the heat shock protein 70 family.,subcellular location:May translocate to the nucleus after heat shock.,subunit:Homodimer.,BackgroundThe protein encoded by this gene is heat shock inducible and may act as a chaperone. The encoded protein can protect the heat-shocked cell against the harmful effects of aggregated proteins. This gene is highly expressed in leukemia cells and may be a good target for therapeutic intervention. Several transcripts encoding different isoforms have been found for this gene. [provided by RefSeq, Dec 2015],	Synonyms	
Tissue SpecificityHeart,Testis,Functionfunction:Possesses chaperone activity in vitro where it inhibits aggregation of citrate synthase.,induction:By heat shock.,PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Belongs to the heat shock protein 70 family.,subcellular location:May translocate to the nucleus after heat shock.,subunit:Homodimer.,BackgroundThe protein encoded by this gene is heat shock inducible and may act as a chaperone. The encoded protein can protect the heat-shocked cell against the harmful effects of aggregated proteins. This gene is highly expressed in leukemia cells and may be a good target for therapeutic intervention. Several transcripts encoding different isoforms have been found for this gene. [provided by RefSeq, Dec 2015],	Observed Band	92kD
Functionfunction:Possesses chaperone activity in vitro where it inhibits aggregation of citrate synthase.,induction:By heat shock.,PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Belongs to the heat shock protein 70 family.,subcellular location:May translocate to the nucleus after heat shock.,subunit:Homodimer.,BackgroundThe protein encoded by this gene is heat shock inducible and may act as a chaperone. The encoded protein can protect the heat-shocked cell against the harmful effects of aggregated proteins. This gene is highly expressed in leukemia cells and may be a good target for therapeutic intervention. Several transcripts encoding different isoforms have been found for this gene. [provided by RefSeq, Dec 2015],	Cell Pathway	Cytoplasm . Nucleus . May translocate to the nucleus after heat shock
<ul> <li>citrate synthase.,induction:By heat shock.,PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Belongs to the heat shock protein 70 family.,subcellular location:May translocate to the nucleus after heat shock.,subunit:Homodimer.,</li> <li>Background The protein encoded by this gene is heat shock inducible and may act as a chaperone. The encoded protein can protect the heat-shocked cell against the harmful effects of aggregated proteins. This gene is highly expressed in leukemia cells and may be a good target for therapeutic intervention. Several transcripts encoding different isoforms have been found for this gene. [provided by RefSeq, Dec 2015],</li> </ul>	Tissue Specificity	Heart, Testis,
chaperone. The encoded protein can protect the heat-shocked cell against the harmful effects of aggregated proteins. This gene is highly expressed in leukemia cells and may be a good target for therapeutic intervention. Several transcripts encoding different isoforms have been found for this gene. [provided by RefSeq, Dec 2015],	Function	citrate synthase.,induction:By heat shock.,PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Belongs to the heat shock protein 70 family.,subcellular location:May translocate to the nucleus after heat
Nanjing BYabscience technology Co.,Ltd	Background	chaperone. The encoded protein can protect the heat-shocked cell against the harmful effects of aggregated proteins. This gene is highly expressed in leukemia cells and may be a good target for therapeutic intervention. Several transcripts encoding different isoforms have been found for this gene. [provided by RefSeq,
		Nanjing BYabscience technology Co.,Ltd



国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务,欢迎咨询



## matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

**Products Images** 

Nanjing BYabscience technology Co.,Ltd