



ZDH17 Monoclonal Antibody

Catalog No	BYmab-06694
Isotype	IgG
Reactivity	Human;Mouse
Applications	WB
Gene Name	ZDHC17 HIP14 HIP3 HYPH KIAA0946 HSPC294
Protein Name	Palmitoyltransferase ZDHC17 (EC 2.3.1.-) (Huntingtin yeast partner H) (Huntingtin-interacting protein 14) (HIP-14) (Huntingtin-interacting protein 3) (HIP-3) (Huntingtin-interacting protein H) (Putat
Immunogen	Synthesized peptide derived from part region of human protein
Specificity	ZDH17 Monoclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	69kD
Cell Pathway	Golgi apparatus membrane ; Multi-pass membrane protein . Cytoplasmic vesicle membrane ; Multi-pass membrane protein . Cell junction, synapse, presynaptic cell membrane ; Multi-pass membrane protein . Low extracellular Mg(2+) induces increase in Golgi and in post-Golgi membrane vesicles. .
Tissue Specificity	Expressed in all brain regions. Expression is highest in the cortex, cerebellum, occipital lobe and caudate and lowest in the spinal cord. Expression is also seen in testis, pancreas, heart and kidney.
Function	catalytic activity:Palmitoyl-CoA + protein-cysteine = S-palmitoyl protein + CoA.,domain:The DHHC domain is required for palmitoyltransferase activity.,function:Palmitoyltransferase specific for a subset of neuronal proteins, including SNAP25, DLG4/PSD95, GAD2, SYT1 and HD. May be involved in the sorting or targeting of critical proteins involved in the initiating events of endocytosis at the plasma membrane. May be involved in the NF-kappa-B signaling pathway. Has transforming activity.,miscellaneous:The early and

Nanjing BYabscience technology Co.,Ltd



	<p>prominent pathology of HD is observed in the medium spiny neurons that project into the globus.,similarity:Belongs to the DHHC palmitoyltransferase family. AKR/ZDHHC17 subfamily.,similarity:Contains 1 DHHC-type zinc finger.,similarity:Contains 5 ANK repeats.,subunit:Binds HD. This interaction is inversely correlated to the length of the polyglutamine tract added to the huntingt</p>
Background	<p>catalytic activity:Palmitoyl-CoA + protein-cysteine = S-palmitoyl protein + CoA.,domain:The DHHC domain is required for palmitoyltransferase activity.,function:Palmitoyltransferase specific for a subset of neuronal proteins, including SNAP25, DLG4/PSD95, GAD2, SYT1 and HD. May be involved in the sorting or targeting of critical proteins involved in the initiating events of endocytosis at the plasma membrane. May be involved in the NF-kappa-B signaling pathway. Has transforming activity.,miscellaneous:The early and prominent pathology of HD is observed in the medium spiny neurons that project into the globus.,similarity:Belongs to the DHHC palmitoyltransferase family. AKR/ZDHHC17 subfamily.,similarity:Contains 1 DHHC-type zinc finger.,similarity:Contains 5 ANK repeats.,subunit:Binds HD. This interaction is inversely correlated to the length of the polyglutamine tract added to the huntingtin protein in Huntington disease.,tissue specificity:Expressed in all brain regions. Expression is highest in the cortex, cerebellum, occipital lobe and caudate and lowest in the spinal cord. Expression is also seen in testis, pancreas, heart and kidney.,</p>
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com

官方热线: 025-5229-8998

监督电话: 15950492658