



# HIPK2 Monoclonal Antibody

<b>Catalog No</b>	BYmab-06734
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human;Rat;Mouse;
<b>Applications</b>	WB
<b>Gene Name</b>	HIPK2
<b>Protein Name</b>	Homeodomain-interacting protein kinase 2 (hHIPk2) (EC 2.7.11.1)
<b>Immunogen</b>	Synthesized peptide derived from part region of human protein
<b>Specificity</b>	HIPK2 Monoclonal Antibody detects endogenous levels of protein.
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
<b>Source</b>	Monoclonal, Mouse,IgG
<b>Purification</b>	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Dilution</b>	WB 1:500-2000
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	
<b>Observed Band</b>	131kD
<b>Cell Pathway</b>	Nucleus, PML body . Cytoplasm. Concentrated in PML/POD/ND10 nuclear bodies. Small amounts are cytoplasmic.
<b>Tissue Specificity</b>	Highly expressed in heart, muscle and kidney. Weakly expressed in a ubiquitous way. Down-regulated in several thyroid and breast tumors.
<b>Function</b>	alternative products:Experimental confirmation may be lacking for some isoforms,catalytic activity:ATP + a protein = ADP + a phosphoprotein.,function:Protein kinase acting as a corepressor of several transcription factors, including SMAD1 and POU4F1/Brn3a and probably NK homeodomain transcription factors. Inhibits cell growth and promotes apoptosis. Involved in transcriptional activation of TP53 and TP73. Phosphorylation of TP53 may be mediated by a TP53-HIPK2-AXIN1 complex. In response to TGFB, cooperates with DAXX to activate JNK. Phosphorylates the antiapoptotic factor CTBP1 and promotes its proteasomal degradation. In the Wnt/beta-catenin signaling pathway acts as an intermediate kinase between TAK1 and NLK to promote the proteasomal degradation of MYB (By similarity). Phosphorylates

**Nanjing BYabscience technology Co.,Ltd**



CBX4 upon DNA damage and promotes its E3 SUMO-protein ligase activity.,induction:By UV.,PTM:Phosphory

**Background**

homeodomain interacting protein kinase 2(HIPK2) Homo sapiens This gene encodes a conserved serine/threonine kinase that is a member of the homeodomain-interacting protein kinase family. The encoded protein interacts with homeodomain transcription factors and many other transcription factors such as p53, and can function as both a corepressor and a coactivator depending on the transcription factor and its subcellular localization. Multiple transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Nov 2011],

**matters needing attention**

Avoid repeated freezing and thawing!

**Usage suggestions**

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## Products Images

Nanjing BYabscience technology Co.,Ltd

网址: [www.njbybio.com](http://www.njbybio.com)

官方热线: 025-5229-8998

监督电话: 15950492658