



# PSD4 Monoclonal Antibody

<b>Catalog No</b>	BYmab-06796
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human;Rat;Mouse;
<b>Applications</b>	WB
<b>Gene Name</b>	PSD4 EFA6B TIC
<b>Protein Name</b>	PH and SEC7 domain-containing protein 4 (Exchange factor for ADP-ribosylation factor guanine nucleotide factor 6 B) (Exchange factor for ARF6 B) (Pleckstrin homology and SEC7 domain-containing protein
<b>Immunogen</b>	Synthesized peptide derived from part region of human protein
<b>Specificity</b>	PSD4 Monoclonal Antibody detects endogenous levels of protein.
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
<b>Source</b>	Monoclonal, Mouse,IgG
<b>Purification</b>	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Dilution</b>	WB 1:500-2000
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	
<b>Observed Band</b>	116kD
<b>Cell Pathway</b>	Cell membrane . Cell projection, ruffle membrane . In interphase associated with the plasma membrane, in particular with membrane ruffling regions (PubMed:23603394). Accumulates in dynamic actin-rich membrane ruffles and microvilli-like structures (PubMed:12082148). Recruited to membranes via phosphatidylinositol phosphate-binding (Probable). .
<b>Tissue Specificity</b>	Widely expressed. Highest levels of expression are found in placenta, pancreas, spleen, thymus and peripheral blood.
<b>Function</b>	function:Guanine nucleotide exchange factor for ARF6-like protein. Involved in membrane recycling.,similarity:Contains 1 PH domain.,similarity:Contains 1 SEC7 domain.,subcellular location:Accumulates in dynamic actin-rich membrane ruffles and microvilli-like structures.,tissue specificity:Widely expressed. Highest levels of expression are found in placenta, pancreas, spleen, thymus and peripheral blood.,

**Nanjing BYabscience technology Co.,Ltd**



Background	function:Guanine nucleotide exchange factor for ARF6-like protein. Involved in membrane recycling.,similarity:Contains 1 PH domain.,similarity:Contains 1 SEC7 domain.,subcellular location:Accumulates in dynamic actin-rich membrane ruffles and microvilli-like structures.,tissue specificity:Widely expressed. Highest levels of expression are found in placenta, pancreas, spleen, thymus and peripheral blood.,
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## Products Images