



## **HES4 Monoclonal Antibody**

Catalog No	BYmab-06852
Catalog No	5 1 max 00002
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	WB
Gene Name	HES4 BHLHB42
Protein Name	Transcription factor HES-4 (hHES4) (Class B basic helix-loop-helix protein 42) (bHLHb42) (Hairy and enhancer of split 4) (bHLH factor Hes4)
Immunogen	Synthesized peptide derived from part region of human protein
Specificity	HES4 Monoclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	24kD
Cell Pathway	Nucleus .
Tissue Specificity	Kidney,
Function	domain:Has a particular type of basic domain (presence of a helix-interrupting proline) that binds to the N-box (CACNAG), rather than the canonical E-box (CANNTG).,domain:The C-terminal WRPW motif is a transcriptional repression domain necessary for the interaction with Groucho/TLE family members, transcriptional corepressors recruited to specific target DNA by Hairy-related proteins.,function:Transcriptional repressor. Binds DNA on N-box motifs: 5'-CACNAG-3'.,similarity:Contains 1 basic helix-loop-helix (bHLH) domain.,similarity:Contains 1 Orange domain.,subunit:Transcription repression requires formation of a complex with a corepressor protein of the Groucho/TLE family.,


Nanjing BYabscience technology Co.,Ltd



国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务,欢迎咨询



Background	domain: Has a particular type of basic domain (presence of a helix-interrupting proline) that binds to the N-box (CACNAG), rather than the canonical E-box (CANNTG)., domain: The C-terminal WRPW motif is a transcriptional repression domain necessary for the interaction with Groucho/TLE family members, transcriptional corepressors recruited to specific target DNA by Hairy-related proteins., function: Transcriptional repressor. Binds DNA on N-box motifs: 5'-CACNAG-3'., similarity: Contains 1 basic helix-loop-helix (bHLH) domain., similarity: Contains 1 Orange domain., subunit: Transcription repression requires formation of a complex with a corepressor protein of the Groucho/TLE family.,
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.



网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658