




## **THOP1 Monoclonal Antibody**

Catalog No         BYmab-06921           Isotype         IgG           Reactivity         Human;Rat;Mouse           Applications         WB           Gene Name         THOP1           Protein Name         Thimet oligopeptidase (EC 3.4.24.15) (Endopeptidase 24.15) (MP78)           Immunogen         Synthesized peptide derived from part region of human protein           Specificity         THOP1 Monoclonal Antibody detects endogenous levels of protein.           Formulation         Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.           Source         Monoclonal, Mouse, IgG           Purification         The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.           Dilution         WB 1:500-2000           Concentration         1 mg/ml           Purity         290%           Storage Stability         -20°C/1 year           Synonyms         Observed Band         75kD           Cell Pathway         Cytoplasm.           Tissue Specificity         Brain, Lung, Placenta, Uterus,           Function         catalytic activity. Perferential cleavage of bonds with hydrophobic residues at P1, residues, orfactor. Sinds 1 zinc ion, function involved in the metabolism of neuropeptides under 20 amino acid residues long. Involved in the metabolism of neuropeptides under 20 amino acid residu		
Reactivity         Human;Rat;Mouse           Applications         WB           Gene Name         THOP1           Protein Name         Thimet oligopeptidase (EC 3.4.24.15) (Endopeptidase 24.15) (MP78)           Immunogen         Synthesized peptide derived from part region of human protein           Specificity         THOP1 Monoclonal Antibody detects endogenous levels of protein.           Formulation         Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.           Source         Monoclonal, Mouse, IgG           Purification         The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.           Dilution         WB 1:500-2000           Concentration         1 mg/ml           Purity         ≥90%           Storage Stability         -20°C/1 year           Synonyms         Observed Band         75kD           Cell Pathway         Cytoplasm.           Tissue Specificity         Brain, Lung, Placenta, Uterus,           Function         catalytic activity-Preferential cleavage of bonds with hydrophobic residues at P1, P2 and P3' and a small residue at P1' in substrates of 5 to 15 residues, cofactor-Binds 1 zinc ion, function: Involved in the metabolism of neuropeptides under 20 anino acid residues long, Involved in explopiamic peptide degradation. Able to degrade the beta-amyloid precursor protein and generate amyloidogenic fragments, similarity, Be	Catalog No	BYmab-06921
Applications WB Gene Name THOP1 Protein Name Thimet oligopeptidase (EC 3.4.24.15) (Endopeptidase 24.15) (MP78) Immunogen Synthesized peptide derived from part region of human protein Specificity THOP1 Monoclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity 290% Storage Stability -20°C/1 year Synonyms Observed Band 75kD Cell Pathway Cytoplasm. Tissue Specificity Brain, Lung, Placenta, Uterus, Function catalytic activity: Preferential cleavage of bonds with hydrophobic residues at P1, P2 and P3' and a small residue at P1' in substrates of 5 to 15 residues, corfactor Binds 1 zinc ion, function: Involved in the metabolism of neuropeptides under 20 amino acid residues long. Involved in explosionic peptide degradation. Able to degrade the beta-amyloid precursor protein and generate amyloidogenic fragments, similarity. Belongs to the peptidase M3 family, subunit: Monomer., Background	Isotype	IgG
Gene Name THOP1 Protein Name Thimet oligopeptidase (EC 3.4.24.15) (Endopeptidase 24.15) (MP78)  Immunogen Synthesized peptide derived from part region of human protein  Specificity THOP1 Monoclonal Antibody detects endogenous levels of protein.  Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.  Source Monoclonal, Mouse, IgG  Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-2000  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms  Observed Band 75kD  Cell Pathway Cytoplasm.  Tissue Specificity Brain, Lung, Placenta, Uterus,  Each of the many of the metabolism of neuropeptides under 20 amino acid residues long. Involved in the metabolism of neuropeptides under 20 amino acid residues long. Involved in the metabolism of neuropeptides under 20 amino acid regardents, similarity: Belongs to the peptidase M3 family, subuniti Monomer.,  The protein encoded by this gene is a kininase that uses zinc as a cofactor. The encoded oligopeptidase cleaves cytosolic peptides, making them unavailable for display on antigen-presenting cells. This protein also cleaves neuropeptides	Reactivity	Human;Rat;Mouse
Protein Name         Thimet oligopeptidase (EC 3.4.24.15) (Endopeptidase 24.15) (MP78)           Immunogen         Synthesized peptide derived from part region of human protein           Specificity         THOP1 Monoclonal Antibody detects endogenous levels of protein.           Formulation         Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.           Source         Monoclonal, Mouse, IgG           Purification         The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.           Dilution         WB 1:500-2000           Concentration         1 mg/ml           Purity         ≥90%           Storage Stability         -20°C/1 year           Synonyms         Observed Band         75kD           Cell Pathway         Cytoplasm.           Tissue Specificity         Brain, Lung, Placenta, Uterus,           Function         catalytic activity: Preferential cleavage of bonds with hydrophobic residues at P1, P2 and P3' and a small residue at P1' in substrates of 5 to 15 residues, cofactor: Binds 1 zinc ion, function: Involved in cytoplasmic peptide degradation. Able to degrade the beta-amyloid precursor protein and generate amyloidogenic fragments, similarity. Belongs to the peptidase M3 family, subunit: Monomer,           Background         The protein encoded by this gene is a kininase that uses zinc as a cofactor. The encoded oligopeptidase cleaves cytosolic peptides, making them unavailable for display on antigen-	Applications	WB
Immunogen         Synthesized peptide derived from part region of human protein           Specificity         THOP1 Monoclonal Antibody detects endogenous levels of protein.           Formulation         Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.           Source         Monoclonal, Mouse, IgG           Purification         The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.           Dilution         WB 1:500-2000           Concentration         1 mg/ml           Purity         ≥90%           Storage Stability         -20°C/1 year           Synonyms         Observed Band         75kD           Cell Pathway         Cytoplasm.           Tissue Specificity         Brain,Lung,Placenta,Uterus,           Function         catalytic activity:Preferential cleavage of bonds with hydrophobic residues at P1, P2 and P3' and a small residue at P1' in substrates of 5 to 15 residues, cofactor:Binds 1 zinc ion per subunit, cofactor:Binds 1 zinc ion, function:Involved in the metabolism of neuropeptides under 20 amino acid residues long, Involved in the metabolism of neuropeptides under 20 amino acid residues long, Involved in the metabolism of neuropeptides under 20 amino acid residues long, Involved in the metabolism of neuropeptides under 20 amino acid residues long, Involved in the metabolism of neuropeptides under 20 amino acid residues long, Involved in the metabolism of neuropeptides under 20 amino acid residues long, Involved in the metabolism of neuropeptides under 20 amino acid resi	Gene Name	THOP1
Specificity       THOP1 Monoclonal Antibody detects endogenous levels of protein.         Formulation       Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.         Source       Monoclonal, Mouse, IgG         Purification       The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.         Dilution       WB 1:500-2000         Concentration       1 mg/ml         Purity       ≥90%         Storage Stability       -20°C/1 year         Synonyms       Observed Band         Cell Pathway       Cytoplasm.         Tissue Specificity       Brain, Lung, Placenta, Uterus,         Function       catalytic activity: Preferential cleavage of bonds with hydrophobic residues at P1, P2 and P3' and a small residue at P1' in substrates of 5 to 15 residues, cofactor: Binds 1 zinc ion per subunit, cofactor: Binds 1 zinc ion, function: Involved in the metabolism of neuropeptides under 20 amino acid residues long. Involved in cytoplasmic peptide degradation. Able to degrade the beta-amyloid precursor protein and generate amyloidogenic fragments, similarity: Belongs to the peptidase M3 family, subunit: Monomer.,         Background       The protein encoded by this gene is a kininase that uses zinc as a cofactor. The encoded oligopeptidase cleaves cytosolic peptides, making them unavailable for display on antigept-presenting cells. This protein also cleaves neuropeptides	Protein Name	Thimet oligopeptidase (EC 3.4.24.15) (Endopeptidase 24.15) (MP78)
Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.  Source Monoclonal, Mouse, IgG  Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-2000  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms  Observed Band 75kD  Cell Pathway Cytoplasm.  Tissue Specificity Brain, Lung, Placenta, Uterus,  Function catalytic activity: Preferential cleavage of bonds with hydrophobic residues at P1, P2 and P3' and a small residue at P1' in substrates of 5 to 15 residues, cofactor:Binds 1 zinc ion, function:Involved in the metabolism of neuropetides under 20 amino acid residues long. Involved in cytoplasmic peptide degradation. Able to degrade the beta-amyloid precursor protein and generate amyloidogenic fragments, similarity:Belongs to the peptides, making them unavailable for display on antigen-presenting cells. This protein also cleaves neuropeptides	Immunogen	Synthesized peptide derived from part region of human protein
Source       Monoclonal, Mouse,IgG         Purification       The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.         Dilution       WB 1:500-2000         Concentration       1 mg/ml         Purity       ≥90%         Storage Stability       -20°C/1 year         Synonyms       Cell Pathway         Cell Pathway       Cytoplasm.         Tissue Specificity       Brain,Lung,Placenta,Uterus,         Function       catalytic activity:Preferential cleavage of bonds with hydrophobic residues at P1, P2 and P3' and a small residue at P1' in substrates of 5 to 15 residues, cofactor:Binds 1 zinc ion per subunit, cofactor:Binds 1 zinc ion, function:Involved in the metabolism of neuropetides under 20 amino acid residues long. Involved in cytoplasmic peptide degradation. Able to degrade the beta-amyloid precursor protein and generate amyloidogenic fragments, similarity:Belongs to the peptidase M3 family., subunit:Monomer.,         Background       The protein encoded by this gene is a kininase that uses zinc as a cofactor. The encoded oligopeptidase cleaves cytosolic peptides, making them unavailable for display on antigen-presenting cells. This protein also cleaves neuropeptides	Specificity	THOP1 Monoclonal Antibody detects endogenous levels of protein.
Purification  The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution  WB 1:500-2000  Concentration  1 mg/ml  Purity  ≥90%  Storage Stability  -20°C/1 year  Synonyms  Observed Band  75kD  Cell Pathway  Cytoplasm.  Tissue Specificity  Brain,Lung,Placenta,Uterus,  Function  catalytic activity:Preferential cleavage of bonds with hydrophobic residues at P1, P2 and P3' and a small residue at P1' in substrates of 5 to 15 residues, cofactor:Binds 1 zinc ion, function:Involved in the metabolism of neuropeptides under 20 amino acid residues long, Involved in cytoplasmic peptide degradation. Able to degrade the beta-amyloid precursor protein and generate amyloidogenic fragments.,similarity:Belongs to the peptidase M3 family.,subunit:Monomer.,  Background  The protein encoded by this gene is a kininase that uses zinc as a cofactor. The encoded oligopeptidase cleaves cytosolic peptides, making them unavailable for display on antigen-presenting cells. This protein also cleaves neuropeptides	Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-2000  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms  Observed Band 75kD  Cell Pathway Cytoplasm.  Tissue Specificity Brain,Lung,Placenta,Uterus,  Function catalytic activity:Preferential cleavage of bonds with hydrophobic residues at P1, P2 and P3' and a small residue at P1' in substrates of 5 to 15 residues.,cofactor:Binds 1 zinc ion, function:Involved in the metabolism of neuropeptides under 20 amino acid residues long. Involved in cytoplasmic peptide degradation. Able to degrade the beta-amyloid precursor protein and generate amyloidogenic fragments.,similarity:Belongs to the peptidase M3 family.,subunit:Monomer.,  Background The protein encoded by this gene is a kininase that uses zinc as a cofactor. The encoded oligopeptidase cleaves cytosolic peptides, making them unavailable for display on antigen-presenting cells. This protein also cleaves neuropeptides	Source	Monoclonal, Mouse,IgG
Concentration       1 mg/ml         Purity       ≥90%         Storage Stability       -20°C/1 year         Synonyms       -20°C/1 year         Observed Band       75kD         Cell Pathway       Cytoplasm.         Tissue Specificity       Brain,Lung,Placenta,Uterus,         Function       catalytic activity:Preferential cleavage of bonds with hydrophobic residues at P1, P2 and P3' and a small residue at P1' in substrates of 5 to 15 residues., cofactor:Binds 1 zinc ion per subunit., cofactor:Binds 1 zinc ion, function:Involved in the metabolism of neuropeptides under 20 amino acid residues long. Involved in cytoplasmic peptide degradation. Able to degrade the beta-amyloid precursor protein and generate amyloidogenic fragments., similarity:Belongs to the peptidase M3 family., subunit:Monomer.,         Background       The protein encoded by this gene is a kininase that uses zinc as a cofactor. The encoded oligopeptidase cleaves cytosolic peptides, making them unavailable for display on antigen-presenting cells. This protein also cleaves neuropeptides	Purification	
Purity ≥90%  Storage Stability -20°C/1 year  Synonyms  Observed Band 75kD  Cell Pathway Cytoplasm.  Tissue Specificity Brain,Lung,Placenta,Uterus,  Function catalytic activity:Preferential cleavage of bonds with hydrophobic residues at P1, P2 and P3' and a small residue at P1' in substrates of 5 to 15 residues.,cofactor:Binds 1 zinc ion per subunit.,cofactor:Binds 1 zinc ion.,function:Involved in the metabolism of neuropeptides under 20 amino acid residues long. Involved in cytoplasmic peptide degradation. Able to degrade the beta-amyloid precursor protein and generate amyloidogenic fragments.,similarity:Belongs to the peptidase M3 family.,subunit:Monomer.,  Background The protein encoded by this gene is a kininase that uses zinc as a cofactor. The encoded oligopeptidase cleaves cytosolic peptides, making them unavailable for display on antigen-presenting cells. This protein also cleaves neuropeptides	Dilution	WB 1:500-2000
Synonyms  Observed Band  75kD  Cell Pathway  Cytoplasm.  Tissue Specificity  Brain,Lung,Placenta,Uterus,  catalytic activity:Preferential cleavage of bonds with hydrophobic residues at P1, P2 and P3' and a small residue at P1' in substrates of 5 to 15 residues.,cofactor:Binds 1 zinc ion.,function:Involved in the metabolism of neuropeptides under 20 amino acid residues long. Involved in cytoplasmic peptide degradation. Able to degrade the beta-amyloid precursor protein and generate amyloidogenic fragments.,similarity:Belongs to the peptidase M3 family.,subunit:Monomer.,  The protein encoded by this gene is a kininase that uses zinc as a cofactor. The encoded oligopeptidase cleaves cytosolic peptides, making them unavailable for display on antigen-presenting cells. This protein also cleaves neuropeptides	Concentration	1 mg/ml
Synonyms  Observed Band  75kD  Cell Pathway  Cytoplasm.  Tissue Specificity  Brain,Lung,Placenta,Uterus,  Function  catalytic activity:Preferential cleavage of bonds with hydrophobic residues at P1, P2 and P3' and a small residue at P1' in substrates of 5 to 15 residues, cofactor:Binds 1 zinc ion, function:Involved in the metabolism of neuropeptides under 20 amino acid residues long. Involved in cytoplasmic peptide degradation. Able to degrade the beta-amyloid precursor protein and generate amyloidogenic fragments.,similarity:Belongs to the peptidase M3 family.,subunit:Monomer.,  Background  The protein encoded by this gene is a kininase that uses zinc as a cofactor. The encoded oligopeptidase cleaves cytosolic peptides, making them unavailable for display on antigen-presenting cells. This protein also cleaves neuropeptides	Purity	≥90%
Observed Band  Cell Pathway  Cytoplasm.  Tissue Specificity  Brain,Lung,Placenta,Uterus,  Catalytic activity:Preferential cleavage of bonds with hydrophobic residues at P1, P2 and P3' and a small residue at P1' in substrates of 5 to 15 residues.,cofactor:Binds 1 zinc ion per subunit.,cofactor:Binds 1 zinc ion.,function:Involved in the metabolism of neuropeptides under 20 amino acid residues long. Involved in cytoplasmic peptide degradation. Able to degrade the beta-amyloid precursor protein and generate amyloidogenic fragments.,similarity:Belongs to the peptidase M3 family.,subunit:Monomer.,  Background  The protein encoded by this gene is a kininase that uses zinc as a cofactor. The encoded oligopeptidase cleaves cytosolic peptides, making them unavailable for display on antigen-presenting cells. This protein also cleaves neuropeptides	Storage Stability	-20°C/1 year
Cell Pathway  Cytoplasm.  Brain,Lung,Placenta,Uterus,  Catalytic activity:Preferential cleavage of bonds with hydrophobic residues at P1, P2 and P3' and a small residue at P1' in substrates of 5 to 15 residues.,cofactor:Binds 1 zinc ion per subunit.,cofactor:Binds 1 zinc ion.,function:Involved in the metabolism of neuropeptides under 20 amino acid residues long. Involved in cytoplasmic peptide degradation. Able to degrade the beta-amyloid precursor protein and generate amyloidogenic fragments.,similarity:Belongs to the peptidase M3 family.,subunit:Monomer.,  Background  The protein encoded by this gene is a kininase that uses zinc as a cofactor. The encoded oligopeptidase cleaves cytosolic peptides, making them unavailable for display on antigen-presenting cells. This protein also cleaves neuropeptides	Synonyms	
Tissue Specificity  Brain,Lung,Placenta,Uterus,  catalytic activity:Preferential cleavage of bonds with hydrophobic residues at P1, P2 and P3' and a small residue at P1' in substrates of 5 to 15 residues.,cofactor:Binds 1 zinc ion per subunit.,cofactor:Binds 1 zinc ion.,function:Involved in the metabolism of neuropeptides under 20 amino acid residues long. Involved in cytoplasmic peptide degradation. Able to degrade the beta-amyloid precursor protein and generate amyloidogenic fragments.,similarity:Belongs to the peptidase M3 family.,subunit:Monomer.,  Background  The protein encoded by this gene is a kininase that uses zinc as a cofactor. The encoded oligopeptidase cleaves cytosolic peptides, making them unavailable for display on antigen-presenting cells. This protein also cleaves neuropeptides		
Function  catalytic activity:Preferential cleavage of bonds with hydrophobic residues at P1, P2 and P3' and a small residue at P1' in substrates of 5 to 15 residues.,cofactor:Binds 1 zinc ion per subunit.,cofactor:Binds 1 zinc ion.,function:Involved in the metabolism of neuropeptides under 20 amino acid residues long. Involved in cytoplasmic peptide degradation. Able to degrade the beta-amyloid precursor protein and generate amyloidogenic fragments.,similarity:Belongs to the peptidase M3 family.,subunit:Monomer.,  Background  The protein encoded by this gene is a kininase that uses zinc as a cofactor. The encoded oligopeptidase cleaves cytosolic peptides, making them unavailable for display on antigen-presenting cells. This protein also cleaves neuropeptides	Observed Band	75kD
P2 and P3' and a small residue at P1' in substrates of 5 to 15 residues.,cofactor:Binds 1 zinc ion per subunit.,cofactor:Binds 1 zinc ion.,function:Involved in the metabolism of neuropeptides under 20 amino acid residues long. Involved in cytoplasmic peptide degradation. Able to degrade the beta-amyloid precursor protein and generate amyloidogenic fragments.,similarity:Belongs to the peptidase M3 family.,subunit:Monomer.,  Background  The protein encoded by this gene is a kininase that uses zinc as a cofactor. The encoded oligopeptidase cleaves cytosolic peptides, making them unavailable for display on antigen-presenting cells. This protein also cleaves neuropeptides		
encoded oligopeptidase cleaves cytosolic peptides, making them unavailable for display on antigen-presenting cells. This protein also cleaves neuropeptides	Cell Pathway	Cytoplasm.
	Cell Pathway Tissue Specificity	Cytoplasm.  Brain,Lung,Placenta,Uterus,  catalytic activity:Preferential cleavage of bonds with hydrophobic residues at P1, P2 and P3' and a small residue at P1' in substrates of 5 to 15 residues.,cofactor:Binds 1 zinc ion per subunit.,cofactor:Binds 1 zinc ion.,function:Involved in the metabolism of neuropeptides under 20 amino acid residues long. Involved in cytoplasmic peptide degradation. Able to degrade the

Nanjing BYabscience technology Co.,Ltd







	amyloidogenic peptides. [provided by RefSeq, Nov 2015],
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## **Products Images**

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658