



## **KLOTB Monoclonal Antibody**

Catalog No	BYmab-06937
Isotype	lgG
Reactivity	Human;Mouse
Applications	WB
Gene Name	KLB
Protein Name	Beta-klotho (BKL) (BetaKlotho) (Klotho beta-like protein)
Immunogen	Synthesized peptide derived from part region of human protein
Specificity	KLOTB Monoclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	114kD
Cell Pathway	Cell membrane ; Single-pass type III membrane protein .
Tissue Specificity	Liver, Testis,
Function	domain:Contains 2 glycosyl hydrolase 1 regions. However, the first region lacks the essential Glu active site residue at position 241, and the second one lacks the essential Glu active site residue at position 889. These domains are therefore predicted to be inactive.,function:Contributes to the transcriptional repression of cholesterol 7-alpha-hydroxylase (CYP7A1), the rate-limiting enzyme in bile acid synthesis. Probably inactive as a glycosidase. Increases the ability of FGFR1 and FGFR4 to bind FGF21.,similarity:Belongs to the glycosyl hydrolase 1 family.,similarity:Belongs to the glycosyl hydrolase 1 family. Klotho subfamily.,subunit:Interacts with FGFR1 and FGFR4.,
Background	domain:Contains 2 glycosyl hydrolase 1 regions. However, the first region lacks the essential Glu active site residue at position 241, and the second one lacks the
Nanjing BYabscience technology Co.,Ltd	

## Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658



国内优质抗体供应商

精准的 WB 检测服务

24H 在线服务,欢迎咨询



	essential Glu active site residue at position 889. These domains are therefore predicted to be inactive.,function:Contributes to the transcriptional repression of cholesterol 7-alpha-hydroxylase (CYP7A1), the rate-limiting enzyme in bile acid synthesis. Probably inactive as a glycosidase. Increases the ability of FGFR1 and FGFR4 to bind FGF21.,similarity:Belongs to the glycosyl hydrolase 1 family.,similarity:Belongs to the glycosyl hydrolase 1 family.,subunit:Interacts with FGFR1 and FGFR4.,
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.
	Products Images

Nanjing BYabscience technology Co.,Ltd