



# CCD80 Monoclonal Antibody

|                           |   |
|---------------------------|---|
| <b>Catalog No</b>         | BYmab-06966   |
| <b>Isotype</b>            | IgG   |
| <b>Reactivity</b>         | Human;Rat;Mouse   |
| <b>Applications</b>       | WB  |
| <b>Gene Name</b>          | CCDC80 DRO1 URB HBE245  |
| <b>Protein Name</b>       | Coiled-coil domain-containing protein 80 (Down-regulated by oncogenes protein 1) (Up-regulated in BRS-3 deficient mouse homolog)  |
| <b>Immunogen</b>          | Synthesized peptide derived from part region of human protein   |
| <b>Specificity</b>        | CCD80 Monoclonal Antibody detects endogenous levels of protein.   |
| <b>Formulation</b>        | Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.  |
| <b>Source</b>             | Monoclonal, Mouse,IgG   |
| <b>Purification</b>       | The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.  |
| <b>Dilution</b>           | WB 1:500-2000   |
| <b>Concentration</b>      | 1 mg/ml   |
| <b>Purity</b>             | ≥90%  |
| <b>Storage Stability</b>  | -20°C/1 year  |
| <b>Synonyms</b>           |   |
| <b>Observed Band</b>      | 104kD   |
| <b>Cell Pathway</b>       | Secreted, extracellular space, extracellular matrix .   |
| <b>Tissue Specificity</b> | Expressed in dermal papilla and dermal fibroblasts (at protein level). Expressed in heart, thymus, placenta, pancreas, colon, epithelium, spleen and osteoblasts.   |
| <b>Function</b>           | function:Promotes cell adhesion and matrix assembly.,induction:Down-regulated in cancer and after osteoblastic differentiation. Up-regulated by dihydrotestosterone (DHT).,PTM:Phosphorylated.,similarity:Belongs to the CCDC80 family.,subunit:Binds to various extracellular matrix proteins.,tissue specificity:Expressed in dermal papilla and dermal fibroblasts (at protein level). Expressed in heart, thymus, placenta, pancreas, colon, epithelium, spleen and osteoblasts., |
| <b>Background</b>         | function:Promotes cell adhesion and matrix assembly.,induction:Down-regulated in cancer and after osteoblastic differentiation. Up-regulated by dihydrotestosterone (DHT).,PTM:Phosphorylated.,similarity:Belongs to the CCDC80 family.,subunit:Binds to various extracellular matrix proteins.,tissue  |

**Nanjing BYabscience technology Co.,Ltd**



specificity:Expressed in dermal papilla and dermal fibroblasts (at protein level).  
Expressed in heart, thymus, placenta, pancreas, colon, epithelium, spleen and  
osteoblasts.,

**matters needing  
attention**

Avoid repeated freezing and thawing!

**Usage suggestions**

This product can be used in immunological reaction related experiments. For  
more information, please consult technical personnel.

## Products Images

Nanjing BYabscience technology Co.,Ltd

网址: [www.njbybio.com](http://www.njbybio.com)

官方热线: 025-5229-8998

监督电话: 15950492658