




FLRT1 Monoclonal Antibody

Catalog No	BYmab-07000
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	WB
Gene Name	FLRT1 UNQ752/PRO1483
Protein Name	Leucine-rich repeat transmembrane protein FLRT1 (Fibronectin-like domain-containing leucine-rich transmembrane protein 1)
Immunogen	Synthesized peptide derived from part region of human protein
Specificity	FLRT1 Monoclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	71kD
Cell Pathway	Call manufacture . Circula mana manufacture . Fundanta maticulum
com ruamuy	Cell membrane; Single-pass membrane protein. Endoplasmic reticulum membrane. Cytoplasmic vesicle membrane. Cytoplasm, perinuclear region. Cell junction, focal adhesion. Secreted. Cell projection, neuron projection. Cell junction. In addition to its location at the cell membrane, colocalizes with FGFR1 in punctate perinuclear cytoplasmic vesicles. Detected along neurites and at contacts between neurite termini and other cells. Proteolytic cleavage gives rise to a shedded ectodomain.
Tissue Specificity	membrane . Cytoplasmic vesicle membrane . Cytoplasm, perinuclear region . Cell junction, focal adhesion . Secreted . Cell projection, neuron projection . Cell junction . In addition to its location at the cell membrane, colocalizes with FGFR1 in punctate perinuclear cytoplasmic vesicles. Detected along neurites and at contacts between neurite termini and other cells. Proteolytic cleavage gives rise to
	membrane . Cytoplasmic vesicle membrane . Cytoplasm, perinuclear region . Cell junction, focal adhesion . Secreted . Cell projection, neuron projection . Cell junction . In addition to its location at the cell membrane, colocalizes with FGFR1 in punctate perinuclear cytoplasmic vesicles. Detected along neurites and at contacts between neurite termini and other cells. Proteolytic cleavage gives rise to a shedded ectodomain.
Tissue Specificity	membrane . Cytoplasmic vesicle membrane . Cytoplasm, perinuclear region . Cell junction, focal adhesion . Secreted . Cell projection, neuron projection . Cell junction . In addition to its location at the cell membrane, colocalizes with FGFR1 in punctate perinuclear cytoplasmic vesicles. Detected along neurites and at contacts between neurite termini and other cells. Proteolytic cleavage gives rise to a shedded ectodomain Expressed in kidney and brain. function:May have a function in cell adhesion and/or receptor signaling.,PTM:N-glycosylated.,similarity:Contains 1 fibronectin type-III domain.,similarity:Contains 10 LRR (leucine-rich) repeats.,tissue

Nanjing BYabscience technology Co.,Ltd







	(FLRT) family. The family members may function in cell adhesion and/or receptor signalling. Their protein structures resemble small leucine-rich proteoglycans found in the extracellular matrix. The encoded protein shares sequence similarity with two other family members, FLRT2 and FLRT3. This gene is expressed in kidney and brain. [provided by RefSeq, Jul 2008],
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658