



# VWCE Monoclonal Antibody

Catalog No	BYmab-07006
Isotype	IgG
Reactivity	Human;Mouse
Applications	WB
Gene Name	VWCE URG11
Protein Name	von Willebrand factor C and EGF domain-containing protein (HBV X protein up-regulated gene 11 protein) (HBxAg up-regulated gene 11 protein)
Immunogen	Synthesized peptide derived from part region of human protein
Specificity	VWCE Monoclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	105kD
Cell Pathway	Secreted . According to PubMed:16496348 is localized in the cytoplasm. .
Tissue Specificity	Expressed in liver.
Function	function:May be a regulatory element in the beta-catenin signaling pathway and a target for chemoprevention of hapatocellular carcinoma.,induction:By HBxAg.,similarity:Contains 4 EGF-like domains.,similarity:Contains 6 VWFC domains.,subcellular location:According to PubMed:16496348 is localized in the cytoplasm.,tissue specificity:Expressed in liver.,
Background	function:May be a regulatory element in the beta-catenin signaling pathway and a target for chemoprevention of hapatocellular carcinoma.,induction:By HBxAg.,similarity:Contains 4 EGF-like domains.,similarity:Contains 6 VWFC domains.,subcellular location:According to PubMed:16496348 is localized in the cytoplasm.,tissue specificity:Expressed in liver.,

Nanjing BYabscience technology Co.,Ltd

**matters needing attention**

Avoid repeated freezing and thawing!

**Usage suggestions**

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## Products Images